


F. Christian Thompson & Adrian C. Pont

Systematic Database of *Musca* Names  
(Diptera)



KOENIGSTEIN  
KOELTZ SCIENTIFIC BOOKS  
1993



Associated Copy  
DO NOT REMOVE FROM FILES  
USDA, ARS, IIBIII  
Systematic Entomology Lab.

J C W Thompson

Date of publication

11 MARCH 94

\* 2641

Printed in Germany!



F. CHRISTIAN THOMPSON & ADRIAN C. PONT

SYSTEMATIC DATABASE OF *MUSCA* NAMES (DIPTERA)



# THESES ZOOLOGICAE

VOLUME 20  
EDITOR: RONALD FRICKE

---

## Systematic Database of *Musca* Names (Diptera)

A catalog of names associated with the genus-group name *Musca* Linnaeus, with information on their classification, distribution, and documentation

by

F. CHRISTIAN THOMPSON & ADRIAN C. PONT



KOENIGSTEIN  
KOELTZ SCIENTIFIC BOOKS  
1993

*Authors' Addresses*

F. Christian Thompson  
Systematic Entomology Laboratory, USDA  
NHB-168 Smithsonian Institution  
Washington, D. C. 20560  
U.S.A.

Adrian C. Pont  
Hope Entomological Collections, University Museum  
Parks Road  
Oxford OX1 3PW  
England

© Koeltz Scientific Books, D-6240 Koenigstein  
Printed in the United States of America  
Published 12.15.1993  
ISBN Germany 3-87429-338-6  
USA 1-878762-37-0  
ISSN 0934-8956

## Contents

Introduction . . . . .	2
Systematic History of <i>Musca</i> . . . . .	3
Earliest times to Linnaeus (1758) . . . . .	3
From Linnaeus to Meigen (1758—1838) . . . . .	10
Meigen to Stein (1838—1921) . . . . .	22
The Modern Era (1921—1992) . . . . .	28
The type species of the genus <i>Musca</i> Linnaeus . . . . .	32
The genus <i>Musca</i> Linnaeus . . . . .	32
Workers: Their collections and publications . . . . .	33
Systematic database of names	
Scope . . . . .	38
Format . . . . .	39
Cut-off date . . . . .	40
List of abbreviations used . . . . .	40
Status of names . . . . .	43
Alphabetical listing of names . . . . .	48
Indices	
By biogeographical realm . . . . .	136
By taxonomic position . . . . .	141
Summary of nomenclatural changes . . . . .	161
Acknowledgements . . . . .	164
References cited . . . . .	166
Postscript . . . . .	220

## Introduction

*There is nothing so disastrous in Science as the arrogant dogmatism which despises the Past and admires nothing but the Present.*

Hoefer

Flies are important to man. They carry diseases, suck the blood of man and his animals, and destroy crops and food, but some are useful as pollinators, predators and parasites of man's enemies. In order to deal effectively with these pests and beneficials, man must have a name for each kind of fly. A kind of fly is called a species, and species are concepts. Names are critical, for without them communication about concepts would be impossible. To be useful, names must be unique: one and only one name for each concept. While names for species have existed since the beginning of recorded time, a formalized system for these names is a recent innovation. The gradual shift to this nomenclatural system has left a confusing array of names, there being the same name for different species and different names for the same species. As a contribution to eliminating this confusion, we have developed a systematic database of the names for flies.

Information does not become obsolete over time. The quality and relative quantity of data may be diminished, but useful information can still be found in old literature. Two examples will suffice to illustrate this point. Despite its antiquity, De Geer's *Mémoires ...* (1773) still contains much useful information about flies and their biology. Vockeroth (1969: 74-76, 139) recognized 4 species groups of the genus *Dasyphorus* based on characters of the male genitalia and abdominal pattern. In a postscript, Vockeroth noted that his species groups agreed moderately well with the subgenera proposed by Dušek and Láska (1967). In addition to adult characters, Dušek and Láska used larval ones, which prompted Vockeroth to conclude "Further study, especially of many currently unknown larvae, may however support the recognition of these subgenera." At that time, Dušek and Láska knew larval examples of 3 of Vockeroth's 4 species groups. Unfortunately due to previous authors' disregard for the literature and the principle of priority, Dušek and Láska did not know that the larva of the 4th species group had also been described by De Geer (*Musca pinastri*). A quick inspection of De Geer's figures reveals that each species group of Vockeroth (or subgenera of Dušek and Láska) is easily recognized in the larval stage by the shape and number of fleshy protuberances.

A second example concerns screwworms. Some screwworms (*Cochliomyia*) are endemic to the New World and others (*Chrysomya*) to the Old World. All screwworms cause great injury to livestock and humans.

Hence, the introduction of Old World screwworms into the New World and vice versa is a phenomenon of great interest to agriculture. The introduction of Old World screwworms into the New World is generally thought to be a modern phenomenon (Gagné, 1981, Baumgartner & Greenberg, 1984), but it may not be so. De Geer described an Old World screwworm (*Musca erythrocephala*) from Surinam more than 200 years ago, but this fact was forgotten by subsequent workers (Thompson, 1973). These workers assumed that the specimens of *Chrysomya megacephala* found in De Geer's collection under the name *Musca erythrocephala*, while agreeing with De Geer's description, could not have been authentic types due to the then known distribution of the species. It is now obvious that De Geer did, in fact, describe an earlier introduction of Old World screwworms into the New World.

To access the older literature, one needs to know that it exists and one must have an index to it. The first step in preparing such an index is determining the meaning of the names under which the information was recorded (that is, developing a list of synonyms). Our database provides this first step for some 2,000 names of flies.

## A Systematic History of *Musca*

*It is an arduous task to give novelty to what is ancient, authority to what is new, interest to what is obsolete, light to what is obscure, charm to what is loathsome, and credit to what is dubious.*

Pliny: Natural History

The word "fly", like the French "mouche" and German "Fliege", has always been a generalized, typological term, and so it was originally with the words for "fly" in Classical Greek ("μύια") and Latin ("musca") which we have now come to accept in our scientific vocabulary as terms with a precise evolutionary meaning. The purpose of this chapter is to trace the metamorphosis and refinement of the concept "musca," from its vernacular origin down to its contemporary and very much more restricted meaning.

### Earliest times to Linnaeus (1758)<sup>1</sup>

Flies, and particularly domestic flies, have been man's constant companions during the entire course of human evolution, and beyond.

<sup>1</sup> For fuller details of the entomology of this period, see Bodenheimer (1928, 1929), Harpaz (1973), Morge (1973), Beier (1973) and Tuxen (1973); and for a detailed consideration of the identity of the insects discussed by classical Greek and Latin authors, see Beavis (1988).

The ancient civilizations of the Mediterranean, our intellectual and cultural forbears, had little precise knowledge of flies. Without the economic resources, social motivation, intense intellectual curiosity, and optical equipment that subsequently became available in 19th and 20th century Europe, they noted only those flies that impinged directly upon their lives, usually anthropomorphising them or endowing them with a numinous aura. For example, the Bible (2 Kings, chapter 1) reports on the worship of Baal-zebub at the Philistine town of Ekron (in modern Israel). Baal-zebub means “lord of the flies”, and by punning with Baal-zebul, “lord of the dung”, the Jews not only included a sneer at polytheism and the worship of idols but implicitly recognized the connection between flies and dung. In the heat of the Middle East, Baal-zebub was worshipped because he gave protection against flies, and even drove them away. The ancient Greeks gave Zeus and certain other gods the cognomina μυιώδης (myiodes: god of flies), απομυιος (apomyios: expeller of flies) and μυιαγρος (myiagros: hunter of flies) because these divinities were able to clear their temples of these insects.

During the period of Classical antiquity, from circa 1000 BC until circa 100 AD, the Greek “μύια” and Latin “*musca*” were used to cover synanthropic flies corresponding to the current families Muscidae and Calliphoridae (Beavis, 1988). These were recognized as distinct from smaller gnats and midges, and usually also from mosquitoes, horse flies, and cattle flies (bots, warbles) for which other terms were available: οιστρος, μυωψ, χωνοψ, εμπις, tabanus, asilus, culex, empis. The greatest zoologist of the period was Aristotle (BC 384-322), whose work was not surpassed for 2000 years. He classified insects firstly according to their wings and secondly according to their mouth-parts. He recognized a group of small insects with two wings, and also reported that the common house fly (*Musca domestica*) developed in heaped manure. His accounts of μύια (e.g. in *The History of Animals* IV, V) and those of other authors contain a curious mixture of sound factual observation and pure superstition and hearsay. For example, different stages of the life-history were known but they were not integrated into a life cycle. Aristotle mentions flies that extrude live maggots, yet he also refers to the spontaneous generation of maggots in dung and other putrefying matter, despite the fact that, at a far earlier period, Homer’s epic poem *Iliad* referred to maggots being produced by flies in the corpses of men slain in battle. The feeding behaviour of μύια was known: Aristotle described them as having a strong “tongue” (χειντπον) with which they could draw blood: he may have had the stable fly (*Stomoxys calcitrans*) in mind at this point, or even certain Tabanidae. Elsewhere he classifies μύια among those insects that feed primarily upon liquids and juices.

In Roman times, the most significant figure was Pliny the Elder (AD 23-79) whose *Historia Naturalis* (AD 77) was founded on Aristotle. This was an encyclopaedic condensation of all that was known at the time, with few original observations. Pliny took up Aristotle's distinction between flies with a fixed biting proboscis and flies with a soft sucking proboscis: “[*muscae*] reliquorum quibusdam aculeus ... quibusdam muscis, omnibus autem his in ore et pro lingua” [“some of the rest have a sting ... and some flies, but with all of these it is in the mouth and serves as a tongue”: interpreted as *Stomoxys calcitrans*]; and “sunt hi aculei quibusdam hebetes, neque ad punctum sed ad suctum, ut muscarum generi, in quo lingua evidens fistula est” [“with some these stings are blunt, and do not serve for pricking but for suction - for instance with a sort of fly, in which the tongue is evidently a tube”: *Musca domestica*].

The Medieval period, up to the close of the 15th century, was one of intellectual regression in scientific knowledge. The temporal and spiritual might of the Roman Catholic church was based upon an authority, tradition, and dogma in which empirical science had no place. Progress was not possible, nor was it desired. What might have been done could not in any case be disseminated, since writing was a monopoly of the monasteries. The dark cloud of ignorance and superstition was gradually dispersed by two events: the invention of printing and the fall of Constantinople, the repository of Classical learning for over 1100 years.

From this immensely long period of stagnation, a few works and authors can be highlighted to illustrate how the concept of “*musca*” was handled. Latin was the language of learning, and so “*musca*” was used for flies in a vernacular rather than a technical sense. Indeed, this continued right down to the time of Linnaeus, with most authors using “*musca*” as we now use “fly,” to include flies, scorpionflies, dragonflies, caddisflies, stoneflies, mayflies, etc.

The *Physiologus*, originating in the 1st century AD, was the most popular zoological book of medieval times but is too overlaid with religious symbolism to be factually reliable: “*musca*,” the fly, is a symbol of the devil and of suffering. Book 12 of *Origines sive Etymologiae* by Isidorus of Seville (560-636) discusses “*musca*” (and also “*culex*,” “*oestrus*,” “*bibio*”), and continues the fiction of the belief in spontaneous generation of maggots in meat. Isidorus mentions biting flies but not house flies. *De Universo* by Rhabanus Maurus (776-856) uses “*muscae*” to include flies, gad flies, and probably *Hippobosca*. *Liber de Natura Rerum* by Thomas Cantipratanus (1201-1263/93) also mentions “*musca*.” Albertus Magnus (1193-1289), in the last book of his *De Animalibus*, discusses “*musca*” and asserts that the

fly has two wings and eight legs. It is thought that both he and Thomas Cantipratanus must have had house flies in mind. Petrus Candidus Decembris (1399-1477) produced a *Codex* of animals that is one of the great exemplars of medieval art. He dealt with a "*Musca proterva*," which is also *Musca domestica*, and with a "*Musca cypri*," which is a fabulous creature addicted, like the salamander, to living in fire ("*in medio ignium impune volant*" — they fly with impunity in the middle of fire).

One bizarre aspect of medieval man's relationship with insects was in the field of pest control. Insects were actually tried before secular and ecclesiastical courts, prosecuted, and sentenced to banishment. In some cases, they were even physically arraigned before the courts and then executed.

Amidst this morasse of scholastic compilation, superstition, and hearsay, the liberating influence first of the Renaissance and then of the Reformation stimulated individuals to look at insects afresh, to make original observations without preconceived notions, and through a genuinely scientific method to bring about an advance in knowledge.

A characteristic work of the mid 16th century was that of the Englishman Edward Wotton (1492-1555), published in 1552. He divided the insects into six groups in a neo-Aristotelian fashion, using mouth-part conformation as his criterion. His group 4 is defined as "*insectae, quae promuscidem seu aculeum in ore gerunt: muscae, culices, cicadae, locustae, traxales*." His chapter ccxiii deals with flies, where he defines two groups: "*culex*" for the nematocerous and biting flies, and "*musca*" for the rest.

Entomological science was considerably advanced by the Italian Ulysse Aldrovandus (1522-1605), a pioneer of pure research. In book 3 of his *De animalibus insectis* (1602) he defined his major groups. A preliminary table included "*musca, tabanus, culex, ephemera*" as a group of two-winged insects. In his text he further defined this group as having no elytra and two wings ("*de anelytris, bipennibus*"). He discussed "*musca*," of which he illustrated 72 species (pp. 345-351), but also placed in the same group certain other "similar" four-winged insects: mayflies were grouped with "*musca*," whilst scorpionflies, lacewings, and some ruby-tailed wasps were included with "*culex*." His "*culex*" group differed from "*musca*" by the harder proboscis and more painful bite.

Similar advances were made in England by Thomas Moufet (1553-1604), whose major work was published posthumously in Latin in 1634 (also in English in Topsel 1658). Three chapters (10-12) deal with "*musca*." Chapter 10 (Topsel version, 1658: 931) contains a delightful anthropomorphism about the fly: "The light, like Truth, he doth exceedingly rejoice in, and doth

behave himself honestly therein and civilly". Chapter 11 "de muscorum differentiis" is concerned with the different kinds of flies known to him. He appears not to discuss the house fly, but his "*musca*" is used in a very broad, almost vernacular, sense and is not confined to two-winged insects. His illustrations of "*musca*" show various Diptera (flesh fly, dog fly, bee fly, etc), caddisflies, scorpionflies, mayflies, parasitic Hymenoptera, alderflies, dragonflies and damselflies, sawflies, and even a cockroach. His text mentions "*Musca lupus*," "*carnivora*," "*canum*," "*equina*," "*bucularia*," though these are used descriptively, and also the related groups "*Asilus*," "*Tabanus*," "*Oestrus*."

The invention of the microscope in Holland at the end of the 16th century opened up new areas for investigation, and the foundation of several academies and scientific societies encouraged dissemination of the results. The Italian F. Redi (1626-1698) was the first entomologist to exploit the microscope. He demonstrated that insects do not arise from decaying matter, but from eggs deposited on suitable substrates by the adult females (Redi, 1668): the era of spontaneous generation was finally over! He illustrated the adult, larva and pupa of a "*Musca cerasorum*" attacking cherries [Tephritidae]. The section of his work dealing with "*moscas*" (Redi, 1778 edition) discussed oestrids and mosquitoes, and also the true flies that produce the maggots in meat and corpses.

Two Dutchmen, J. J. Swammerdam (1637-1680) and A. van Leeuwenhoek (1632-1723) further exploited the power of the microscope. Swammerdam (1685) classified the insects into four classes, according to their metamorphosis, and this reflects real advances made in the acquisition of biological knowledge. His "ordo IV, nympho-vermiformis" comprises "*musca latinarum, asili, tabani, equina, stercoraria, acarus*." The text of his work discusses many different kinds of "*musca*," including mayflies and scorpionflies, so it is clear that he too was using "*musca*" in a vernacular sense here. He knew the flies that come from larvae in rotting meat and that live as parasites in other insects, but did not mention the house fly. He described (1685: 189, plate 11) the egg, larva, and puparium of "*Musca vulgaris latrinicola*" (in index as: "*Musca latrinicola vulgaris*"), which is clearly a species of *Fannia* and probably *scalaris* (Fabricius) ("colorisque est nigricantibus, sed quadam canitie temperat"). The great microscopist Leeuwenhoek (1719: 104-119) discussed "*muscas domesticas*" and other "*muscae*".

A further advance came with the work of the Englishman John Ray (1628-1705), who followed Swammerdam in erecting a system based on morphological and biological facts. Ray is credited with being the first

zoologist to recognize the species and genus as systematic units, defining the species as a collective idea for the offspring from the same or identical progenitors and the genus as a group of species that resemble each other. His great work was published posthumously (Ray, 1710). Flies, “*muscae*,” are divided into those with two wings and those with four. The section of “*muscae bipennes*” (p. 266) describes 39 species. These include “*musca carnaria vulgaris*,” the common flesh fly (Sarcophagidae), and “*musca bombylium referens, humeris rufis, cauda albicante*,” a bee fly (Bombyliidae) that occurs in early spring. On p. 274 the section “*muscae quadripennes*” begins, e.g. “*musca quadripennis, alis longis angustis Papilionum in modum variegatis*.” A number of parasitic “*muscae*” are described in the “*Vespa ichneumon*” group (p. 256-262).

Alongside the work of these system builders — the architects — other work, less ambitious in its aims but equally worthy, was being carried out by the builders of the individual bricks in the edifice of knowledge — the artisans. These further illustrate just how broad the concept of “*musca*” still was in the 17th and early 18th centuries. C. Schwenckfeld (1603) described the insects of Silesia, illustrating a series of species of “*musca*.” Whilst his “*musca*” was defined as having two wings (“*insectum volatile, bipenne*”), many of the insects he included were four-winged: “*Musca infectoria, eine Schmeissfliege*” (Sarcophagidae), “*Musca viridis, eine grüne Fliege*” (Caliphoridae), “*Musca versicolor, eine schöne bundte Fliege*” (?Syrphidae), “*Musca ligniperda, eine Holzfliege/Borfliege*” (Hymenoptera), “*Musca grandior oculata*” and its many forms (Odonata), “*Musca caudata, eine Schwanzfliege*” (Plecoptera). The Dutchman J. Goedart (1620-1668) described 144 numbered experiments in insect breeding (1662a, b, c; English translation by Lister, 1682). He used no names: the description of the immature stages and adults were accompanied by illustrations, and he referred simply to “the fly opposite” or to the fly in the plate. One experiment (1682: 123-124, no. 130) concerned the little house fly, *Fannia canicularis*, and is worth quoting in full:

I had a mind to try, what wou'd become of the putrid and corrupted Urine of a man; I made a Funnell of paper, and so folded it, that no Fly or other Animall cou'd get into it: having infused into it oft times humane Urine; I found some some Worms to be bred in the folds, where the feces stayed. I reserved one of them, for the experiment sake, which composed it self for change, the first of March: and the 14th. of the same moneth, it had the forme of a Fly, depicted in the Table: its head was Red, the body Black, and the hinder-parts Yellowish.

Another Dutchman, S. Blankaart (1688) published a delightful little book with coloured plates and a text in Dutch, dealing with the immature stages of various insects. No. XXIII, page 81, plate 10 (of larva, puparium, adult)

concerns “*van de Worm met het Vliegjen uit de Hoender mest gewassen*” and is obviously *Fannia scalaris*. No. XXII, page 78, plate 10 (of larva, puparium, adult) deals with a rat-tailed maggot, “*van de Swijn made mit zijn Tonnetjen en Vlieg*,” and this has the only Latin epithet of the book, “*Musca apiformis*.” Finally, a German work by J. L. Frisch (1660-1743), published between 1720 and 1738, described biologies of various insects. The house fly appears not to have been described, but there is, for example, a Sarcophagid with illustrations of the larva, spiracles and adult, “*von den Schmeiss-Fliegen mit grau- und schwartzwürflichen Hinterleib*” (1728: 21, plate 14). Another, described as “*Die Fliege ist gläntzend-schwarz, so herauskriecht*,” could be *Hydrotaea* (1732: 8, plate 6, “*von einer kleinen Art der Aas-Fliegen*”).

The last great figure before Linnaeus was the Frenchman R. A. F. de Réaumur (1683-1757), a prolific and enormously influential scientist. He stressed, above all else, the importance of personal observation and personal experience. His *Mémoires* consist of a series of morphological and biological observations, written in French and with a vernacular nomenclature (“*Mouche*,” “*Mouches*,” “*genre des Tipules*”). In his treatment of “*mouches*” (1738) he divided them into a number of groups based on the mouth-parts. The two-winged flies that he recognized (e.g. blue bottles, house flies, mosquitoes) were assigned to the first two classes: (1) “*mouches qui ont une trompe, & qui n'ont point de dents ou de ferres*,” and (2) “*mouches qui ont une bouche sans dents sensibles*.” The other groups are filled with four-winged flies such as scorpionflies, caddisflies, etc. The full range of his “*mouches*” can be seen on plates 8-11. Additional characters that he considered were the antennae, the mode of crossing the wings, and the body-length. He also discussed the biology of flies. Réaumur noted that there were viviparous and oviparous higher flies, observed the method of adult eclosion using the ptilinum, and reared blue bottles, green bottles, flesh flies, and house flies from meat. An interpretation of Réaumur’s species, which included *Musca domestica* and a number of new species, was given by Vallot (1802).

Shortly before the appearance of the 10th edition of Linnaeus’ *Systema Naturae*, another Frenchman, L. Joblot (1754), published a series of microscopical studies of insects, written in French, which illustrate the continuing, broad, multi-ordinal concept of “fly.” Four species of “*mouches*” were described: “*la mouche commune*” (p. 9, pl. 5, Calliphoridae); “*...mouche qui se voit ordinairement sur la surface du vin et sur celle du vinaigre*” (p. 12, pl. 6, Drosophilidae); “*...mouche aquatique et terrestre qui nage sur le dos, qui saute et marche étant hors de l'eau*” (p. 22, pl. 11, Notonectidae); “*...une autre espèce de mouche aquatique et terrestre*” (p. 26, pl. 7, Corixidae).

### From Linnaeus to Meigen (1758-1838)<sup>2</sup>

In following the history of *Musca* during this period, one can trace a direct line of descent from Linnaeus to Fabricius to Meigen. Most authors followed the lead given by these three authors, sometimes adding small details to the system.

With hindsight, it is easy to see that the 10th edition of the *Systema Naturae* was a quantum-leap forward in systematic zoology, an example of "paradigm forming science" (Kuhn, 1962). Yet, at the time, its impact was neither instantaneous nor universal, and there is no caesura in the scientific literature at 1758. Carl von Linnaeus (1707-1778) only arrived gradually at his binominal nomenclature: originally using one word for the genus and up to 12 for the species, he refined this until he was consistently binomial in 1749 for plants (*Pan Svecicus*) and 1754 for animals (*Museum S:ae R:ae M:itis Adolphi Friderici*). In 1758, he gave a well-defined view of orders, genera and species, and a means for distinguishing the species. These conspicuous merits, together with his clarity of thought and limpidity of style, helped to promulgate both his system and his method.

Yet the 1st edition of *Systema Naturae* (1735) was not a great advance on contemporary knowledge and gave little indication of what would eventually follow. There was very scant attention to species. Linnaeus recognized four insect orders, defined by wing form: Coleoptera, Angioptera, Hemiptera, Aptera. Within each of these, he listed "genera," with a diagnosis and a note of the characteristic taxa. Nine "genera" are listed in the Angioptera, for example:

<i>Ephemera</i>	<i>Cauda setosa, alae 4 erectae.</i>	<i>Musca ephemera.</i>
<i>Panorpa</i>	<i>Cauda cheliformis, alae 4, rostrum comeum.</i>	<i>Musca scoriurus.</i>

The last genus is *Musca* (1735: 77):

<i>Musca</i>	<i>Stylus sub alis capitatus. Aiae 2.</i>	<i>Musca diversarum species Oestrum vet. ----- lapporum Tabanus</i>
		<i>Culex teredo navium Tipula. Fomicaleo</i>

It is curious to see the ant lion (Neuroptera: *Myrmeleon*) and the ship worm (Mollusca: *Teredo*) in this list!

---

<sup>2</sup> For more general discussion of this period, see Tuxen (1973) and Lindroth (1973).

There was little change in editions 2 and 3, but in edition 4 (1744: 95) Linnaeus used the term Diptera for his ordo 6 and thus arrived at a conception of two-winged flies that is not much different from our own. He listed seven genera, but clearly had not yet clarified his ideas on nomenclature, using “*Musca*” at the species-level in a vernacular sense:

Genus 172.	<i>Asilus</i>	
	<i>Musca rapax</i>	<i>l'A sille</i>
Genus 173.	<i>Hippobosca</i>	
	<i>Musca equina</i>	<i>la mouche arraignée</i>
Genus 175.	<i>Musca</i>	
	<i>Musca</i>	<i>Os proboscide absque dentibus.</i>
		<i>la mouche</i>

And even in other orders:

Genus 164.	<i>Ephemera</i>	
	<i>Musca ephemera</i>	<i>l'ephémère</i>
Genus 166.	<i>Panorpa</i>	
	<i>Musca scoriurus</i>	<i>la mouche-scorpion</i>

The first edition of *Fauna Svecica* (1746) marked a very striking advance on this. Of course, it was primarily a work on the Swedish fauna rather than a work on classification, and so the species are treated in great detail. There are 7 genera of Diptera, as in the preceding edition of the *Systema*, and the genus *Musca* is dealt with on page 313. Linnaeus describes 64 species in *Musca*, assigned to six groups: *Alis variegatis*; *Hirsutae dictae*; *Variegatae*; *Aphidivora*; *Auratae*; and *Vulgares*. Each species is given a description, with references to earlier works, biology, distribution, etc. The nomenclature is not binomial in the formal sense, the Latin being purely descriptive, e.g.: *Musca fusca*, *oblonga*, *nigra* or *Musca nigra*; *abdomine nitido tessellato*, *thorace lineis pallidioribus longitudinalibus*; *minor*.

In Edition 6 of the *Systema Naturae* (1748) the treatment was more expanded when compared with the previous editions. Linnaeus listed several species in each genus, but still used *Musca* in a partly vernacular sense, e.g. *Musca triplis* in *Ichneumon* (p. 64) and *Musca aestivalis* in *Phryganea* (p. 62). The same seven genera were recognized in the Diptera, and Linnaeus was able to refer back to their descriptions in the *Fauna Svecica* (1746). *Musca* (page 66) is defined as follows: *Os cum proboscide absque dentibus*. The eleven characteristic species that are mentioned are arranged into the six groups (a, wing patterned; b, hairy; c, body patterned; d, aphidophagous; e, gilded; f, common) introduced in *Fauna Svecica*:

a. <i>Alis variegatis</i>	<i>Musca aquatilis</i>
b. <i>Hirsutae</i>	<i>Musca solstitialis</i>
	<i>Musca cerasorum</i>
	<i>Musca stercoraria</i>

<i>c. Corpore variegato</i>	<i>Musca mirabilis</i>
	<i>Musca putrida</i>
	<i>Musca latrinaria</i>
<i>d. Aphidivora</i>	<i>Proboscis elephantis</i> [= <i>Scaeva pyrastris</i> ]
<i>e. Auratae</i>	<i>Musca carnaria</i>
<i>f. Vulgares</i>	<i>Musca domestica</i>
	<i>Vermiculus casei</i> [= <i>Piophila casei</i> ]

Editions 7, 8, and 9 (1748, 1753, 1756) were essentially identical with edition 6 so far as the treatment of *Musca* is concerned.

Linnaeus had used the wings to delimit the orders, and by the 10th edition (1758) he had come to use the mouth-parts to define his genera. His diagnosis for *Musca* (1758: 589), repeated with little change for the next half century, reads: “*Os proboscide carnosa: labiis 2 lateralibus: Palpi nulli*” (mouth with a fleshy proboscis: with two lateral lips: palpi absent). The division of the 100 species of *Musca* into sections (1758: 589, footnote) uses characters of the antenna and arista, and the body vestiture:

<i>Muscae distingvuntur antennis inermibus 1. armatis</i>
<i>Filatae antennis absque seta aut pluma laterali</i>
<i>Armatae antennis cum seta aut pluma laterali</i>
— <i>Tomentosae corpore pubescentes, saepe vix manifeste</i>
— — <i>Plumatis antennis pectinatae</i>
— — <i>Setariis antennis simplici seta</i>
— <i>Pilosae raris setis, imprimis in thorace conspicuis</i>
— — <i>Plumatis antennis</i>
— — <i>Setariis antennis</i>

Broadly speaking, species 1-13 are modern Stratiomyidae and Rhagionidae; 14-46 are Syrphidae; 47-55 are calyprates, including *Musca domestica*; 56-100 are calyprates and acalyprates. It is worth examining the entry for species no. 54, *Musca domestica* (1758: 596) to illustrate Linnaeus' comprehensive yet succinct method of dealing with diagnosis, bibliography, distribution, and biology.

The second edition of his *Fauna Svecica* (1761) also adopted the new nomenclature. It was a descriptive work, not a classificatory work, and *Musca* was used in the sense of 1758. It is interesting to note that Linnaeus has been accused of failing to adopt his own binomial nomenclature consistently in this work, because an appendix (p. 545-556) contains a number of descrip-

<sup>3</sup> The gilded fly (e) is evidently not the flesh fly, our *Sarcophaga carnaria*, but a greenbottle, genus *Lucilia* (cf. Shakespeare's "King Lear" (IV, 6, 115), "the small gilded fly does lecher in my sight.")

this work, because an appendix (p. 545-556) contains a number of descriptions for which the new nomenclature was not used. In fact, even a cursory examination shows that these descriptions are taken from earlier works by himself and De Geer. For example, the description of "*Musca alis albis*" (no. 2311, p. 553) is identical with that of no. 1062, p. 315, in the 1st edition of *Fauna Svecica*. In fact, the title of the appendix states quite explicitly why this is done: "an appendix in which I have set apart animals that I have collected in former times, but which I have subsequently not found for further examination." In other words, Linnaeus, the conscientious taxonomist, did not care to offer diagnoses and names for species that he had not studied for years and that were no longer available to him. [Unfortunately, some subsequent workers were not so conscientious and provided names for these species.]

Several editions and translations of *Systema Naturae* followed, all of which attempted to maintain the Linnaean system whilst incorporating the advances and additional species described by other authors. The 12th edition, by Linnaeus himself (1767), offered no change in the definition of *Musca*, and described 129 species. The de Villers edition (1789) also followed the 1758 diagnoses and divisions, incorporated the works of De Geer, Fabricius, Scopoli, and Schrank, and described 398 species including new species and new names. The Gmelin edition (1790), the 13th edition, followed Scopoli's (1763) modification of the generic description, in particular "*palpi nulli*" became "*palpis duobus brevibus*" ("with two short palpi"). Gmelin also included new species, some new names, and some collection names from "Mus. Lesk.", the species numbered but not named by Zschach (1788). Manuel (1792, 1811) continued the Gmelin tradition with a French version of the species diagnoses. Turton (1801) published an English translation of the Gmelin edition but followed the classification of Fabricius (1794) for insects. Turton maintained the increasingly inadequate definition of *Musca* by its mouth-parts and included 521 species. He marked the Linnaean species with \*, but otherwise gave almost no references to preceding authors. Thus, when an unfamiliar *Musca*-name is encountered, it is extremely difficult and time-consuming to determine if Turton is describing a new species, proposing a new name, or transferring into *Musca* a species already described in another genus, such as *Syrphus* (see discussion on page 43).

Linnaeus' foremost pupil was J. C. Fabricius (1745-1808). He worked hard to develop classificatory ideas based on mouth-parts, and was unwilling to admit that the use of other characters would contribute either to clarity or to a "natural" system: he considered that the greater the number of characters used, the greater would be the ensuing chaos. In his first work (1775) he

divided the genera of Diptera into two sections: “*os proboscide atque haustello*” (10 genera, including *Musca*) and “*os haustello absque proboscide*” (9 genera, including a few Arachnida). Apart from the proliferation of generic names, this is really scarcely different from Pliny’s or even Aristotle’s division of the Diptera! His diagnosis of *Musca* (1775: 773) marks a slight advance on Linnaeus: “*Os proboscide atque haustello. Haustellum seta unica absque vagina. Palpi duo porrecti, extrorsum crassiores. Antennae setariae.*” *Musca* is divided into two groups: “*antennis seta plumata*” (including *domestica*) and “*antennis seta nuda*” [arista plumose or arista bare]. Very broadly, this divides calyptates from acalyptates, as almost all the modern Nematocera, Brachycera, and Aschiza originally placed in *Musca* were recognized as very different and were transferred to other genera (*Bibio*, *Rhagio*, *Rhingia*, *Stratiomys*, *Syrphus*). This framework was more or less followed in his subsequent works (1777, 1781, 1787), but in 1794 (and 1798) he offered several modifications to make allowance for a number of additional genera separated from *Musca*: “*Haustellum seta unica absque vagina. Palpi duo porrecti, extrorsum crassiores. Antennae breves, clavatae, seta instructae.*” [Proboscis with a single bristle, without a sheath. Two palpi, porrect, thickened towards tips. Antennae short, club-shaped, with a bristle.]

His final work on Diptera (1805) was a monograph of the entire order. 81 genera were dealt with, including 5 now in Arachnida, and there is further refinement of the concept of *Musca*. Two generic diagnoses (1805: xii & 283) embrace the characters: “*Os haustello absque vagina. Palpi porrecti, compressi, ciliati, biarticulati, extrorsum crassiores, obtusi. Antennae breves, incumbentes, triarticulatae: articulo tertio elongato, compresso, oblongo.*” A footnote on p. 283 gives a detailed description of *Musca*. The genus is placed after *Stomoxys* and before *Tachina* and *Ocyptera*. His old division within the genus, of species with plumose arista or with bare arista, is still kept, but on this occasion there are 76 species (including *domestica*) in the “plumose” section and 45 in the “bare” section. These sections are close to the longstanding 19th century division between Muscidae and Anthomyiidae, though considerable more refinement, with the removal of many acalyptrates, was carried through by Meigen.

Contemporaneously with Linnaeus and Fabricius, entomologists were beginning to investigate in more detail their local faunas and the biology of insects. Few had a grasp of insect systematics in any way comparable with that of Linnaeus and Fabricius, yet several of them produced noteworthy insights. As always, there were those who welcomed innovative and constructive change and those who clung stubbornly to familiar certainties.

Charles De Geer (1720-1778), who saw himself as the successor of Réaumur, published a series of observations in 7 volumes of *Mémoires* (1752-1778). He gave diagnoses in Latin in a pre-Linnaean style, and volume 3 (1773) is accepted as the first to use binominal nomenclature though it is evident that De Geer was not fully happy with it, or, more probably, was simply indifferent. His method was to give a French and Latin diagnosis, as for *Musca pinastri*:

*MOUCHE velue à poil simple ...*  
*Musca (pinastri) ...*

He sometimes gave replacement names for those Linnaean names that he felt to be inappropriate (see also the discussion in Persson, Pont & Michelsen, 1984):

*Mouche très-velue ...* = [French diagnosis]  
*Musca (bombylius) ...* = [New name & Latin diagnosis]  
*Musca (mystacea) ...* = [Linnaeus' name & diagnosis]

Occasionally he lapsed into trinomials, e.g. *Musca domestica major* [i.e. *Musca domestica* Linnaeus] contrasted with *Musca minor domestica* [i.e. *Fannia scalaris* Fabricius] (see discussion below under workers). His names were tabulated in Linnaean style by Retzius (1783). Volume 6 of the *Mémoires* (1776) dealt with *Musca*, and De Geer followed Geoffroy (1762) in his diagnosis of the genus: “*Antennes à palettes avec un poil latéral. Trompe à lèvres charnues.*” He accepted four subgroups based, as in Linnaeus, on aristal and body vestiture. The German translation of this volume (Goeze, 1782) made no change to the nomenclature or presentation.

Among the many enumerations of local faunas, the diagnosis of *Musca* by Linnaeus (1758) was followed closely by Poda (1761) working in the Graz area of Austria; Sulzer (1761) in Switzerland, who also gave fine illustrations of members of each of the 5 subgroups that he recognized; Houttuyn (1761-1763) in the Netherlands; Brünnich (1764) in Denmark, in a remarkably comprehensive and succinct textbook of entomology that included parallel keys in Latin and Danish to insect genera and their subgroups; Müller (1764, 1776) and Pontoppidan (1765), also in Denmark; Füesslin (1775), Sulzer (1776) and de Razoumowsky (1789) in Switzerland; Schrank (1781) in Austria; Schellenberg (1803) in Switzerland, in a series of beautiful colour plates of species belonging to *Musca*. Even Panzer (1794-1806) took an essentially Linnaean view of *Musca*, though the later volumes of paintings followed Meigen's genera (see the list of Panzer species compiled by Saunders, 1888). Shaw (1806) still used *Musca* exactly as diagnosed by Linnaeus almost 50 years earlier.

The Englishman Moses Harris (1776-1780) followed the Linnaean system and concept of *Musca*. However, he departed from Linnaeus in his classification of the components of *Musca*. Harris was the first entomologist to recognize the importance of wing venation as a character system for classification. Whereas Linnaeus divided *Musca* into 7 groups based on antennal (arista) structure and body pilosity, Harris divided the genus into 9 groups based largely on wing venation. He divided the genus into orders and the orders into sections, and for each he gave an enlarged figure of the wing venation. He did not describe the venation, merely referring to the figure. In the text, he frequently gave additional characters for his groups.

### Harris' subgroups of *Musca*

Order I	Calyprate flies with anteriorly curved M
Order II	Genus <i>Volucella</i> (Syrphidae)
Order III, Section 1	Syrphidae (Eristaliini) with arista pilose
Order III, Section 2	Syrphidae, with arista bare
Order IV, Section 1	Stratiomyidae with scutellar spines
Order IV, Section 2	Stratiomyidae (Beridinae) without scutellar spines
Order V, Section 1	Acalyprates (complete anal vein, parallel wings)
Order V, Section 2	Calyprates (incomplete anal vein, divergent wings)
Order V, Section 3	Dolichopodidae

Unfortunately his work was ignored by Fabricius and subsequently forgotten.<sup>4</sup> Fallén and Meigen rediscovered the importance of wing venation. Harris used binomial nomenclature, that is, he used single words for species names. However, he gave genus group names in the plural (*Muscae*, *Asili*, *Tabani*, *Tipulae*) and his species names included not just nouns and adjectives but also verbal forms (*restituo*, *timeo*, *recurro*) and adverbs (*delicate*, *scitule*, *semulator*) which renders them unavailable according to the current rules of nomenclature. Harris also provided some interesting biological observations. Under his section 1 of *Musca* (which included mostly Tachinidae), he concluded that the large compound eyes in flies were not for vision but that the simple ocelli were: he put opaque white paint on the eyes of a blue-bottle fly and noted that the fly was still able to fly directly to the window, whereas this did not happen when the ocelli were covered.

The Linnaean definition of *Musca* was substantially modified by Geoffroy and Scopoli. Geoffroy (1762) separated several groups (*Stomoxys*, *Stratiomys*, *Volucella*) from *Musca* and introduced new characters into his definition of the genus, stating that it was to be recognized by a combination of characters:

<sup>4</sup> "Moses Harris never ought to have been ignored in the Diptera, in which he was more than an equal with his contemporaries in nomenclature while he was almost 50 years in advance of any other dipterologist in venation" (Verrall, 1901a: 571)

*Antennes formées par une palette plate et solide avec une soie ou poil latéral. Bouche avec une trompe sans dents. Trois petits yeux lisses ... Il dépend de deux parties, savoir des antennes et de la bouche: c'est la configuration de ces deux parties qui constitue le caractère de la mouche."*

Geoffroy divided *Musca* into families, but these subdivisions were essentially the same as those used by Linnaeus (1746, 1748). He did not use a binomial Latin nomenclature, so that his 88 *Musca* species were only validated when subsequent authors happened to mention them and give them Latin or latinized names (principally Geoffroy himself in Fourcroy, 1785; but also Schrank, 1781, 1803; de Villers, 1789; Müller, 1776; even Macquart, 1834<sup>5</sup>). For example: *La mouche à l'étoile* of Geoffroy (1762: 494, no. 3 *Musca cinerea, alis albis apice macula rotunda radiata*) is 1) *M. stellata* of Fourcroy (1785: 470, no. 3) and 2) *M. stellata* of de Villers (1789: 535, no. 333).

Scopoli (1763) advocated using all available structures for classification and modified the Linnaean diagnosis of *Musca*, pointing out that palpi are actually present: "Os vagina carnosa, retractili: apice labiata: basi palpigeria" and (p. 325) "Os armatum rostro retractili: apice dilatato, basi palpigerio: palpis clavatis." Scopoli (1763: 350, footnote) admitted that he was, unfortunately, unable to detect the mouthpart characters necessary to implement the Linnaean classification. He gave essentially the same diagnosis in a later paper (1777). The subdivisions of *Musca* adopted by Scopoli were very different from Linnaeus (1758). He utilized a wide range of structural features (position of wing at rest, abdominal shape & antennal posture) and made many valuable insights.

Schaeffer (1766b) followed both Geoffroy and Scopoli, for example including the presence of palpi and of three ocelli in his description. Yeats (1773) gave a collation of the Geoffroy, Schaeffer, Scopoli, and Linnaeus systems, and Müller (1775) gave a similar review incorporating the results of Houttuyn (1761-1763). Even in the late 1770's, works were appearing in which binomial nomenclature was not accepted, e.g. Schaeffer (1776a: plate 17, fig. 4 "MUSCA spathula antennarum truncata PRIMA. Erste Hausfliege mit abgestutztem Spadel."). Even Barbut (1781), in an exposition of the Linnaean system, listed one of his three cited *Musca* as "*Musca vespae similis*," which is actually *Musca vespariformis* Linnaeus. Grossinger (1794), in a manual of entomology covering systematics, biology, practical importance, ecology, etc., used *Musca* in the Linnaean sense but mentioned neither Linnaeus nor Fabricius: His sources were Aldrovandus, Swammerdam,

<sup>5</sup> See *Onesia fulvipennis* sp. n. (Macquart, 1834: 175), "Musca no. 60 Geoff. En automne sur le lierre"; but already given the name *Musca hederacea* by Geoffroy (1785: 491).

Schwenckfeld, Goedart, Frisch, and, amongst the “classical” systematists, Poda.

The Fabrician era, as already mentioned, saw a proliferation in the number of genera and a progressive refinement in the definition of *Musca*, culminating in Fabricius (1805). Gradually, Fabricius’ more restricted concept of *Musca* supplanted that of Linnaeus. Herbst (1787) gave a German diagnosis of *Musca*, using characters of the antennae and mouth-parts, and pointed out that *Musca* of Linnaeus was the equivalent in Fabricius (1777) of *Bibio*, *Stratiomys*, *Rhagio*, *Syphus* and *Musca*. Forster (1788, 1794) and Roemer (1789) gave an identical diagnosis of *Musca*: this is a very pragmatic version, combining elements of Linnaeus, Fabricius, and Geoffroy. Fabricius (1787) was followed by Rossi (1790, 1794) and Illiger (1807), Petagna (1792), Fabricius himself (1794) and Cederhielm (1798). Schrank (1803) added several characters to the diagnosis of *Musca*, such as the conformation of the abdomen and halteres, but these were not discussed by Fabricius (1805). Nor could the elderly Fabricius accept the much more radical proposals put forward by Meigen (1804), though these two discussed their views and their differences in the course of personal meetings.

Zoologists working in France during the late 18th and early 19th centuries were system-builders, and by using characters from all parts of the insect anatomy were moving towards a more “integrated” system. Lamarck (1801, 1816) and especially Latreille (1796, 1802, 1809, 1810, 1818, 1825, 1829) were the most prolific and influential figures. Latreille was the first to introduce the concept of the “family,” and was highly adaptable, modifying his views from work to work as knowledge of the groups advanced. The definition of *Musca* was based on Geoffroy and Fabricius, and typically runs as follows: “*Antennes à palette, composées de trois articles, dont le dernier porte une soie latérale. Trompe membraneuse, à orifice bilabié, rétractile en entier. Sucoir d'une ou deux soies. Deux antennules insérées sur la trompe.*” (Lamarck, 1801: 310). Within this Lamarck recognized two sections of *Musca*, one with a plumose arista (citing *Musca domestica*) and one with a bare arista. Latreille (1796: 170; 1802: 452) was the first to use the size of the calypters as a feature in classification, and this is still reflected in the “Calyprate” and “Acalyprate” sections of the Muscomorpha (Schizophora). At this time he was using *Musca* in an early Fabrician sense, but as the number of genera multiplied during the first decades of the 19th century he adapted his own classification to accommodate them (Latreille, 1809, 1818, 1825). Latreille generally cited *vomitoria sensu* Fabricius (often with other species) as the typical species of *Musca*.

In a curious work, Walckenaer (1802; see also Bezzi, 1907) transferred 7 species of "Musca," including *domestica* and *vomitoria*, to Latreille's genus *Lispe*, retaining 25 species (including modern Tachinidae, Muscidae, Scathophagidae, Acalyptrates) in the old genus *Musca*. The old-fashioned work of Samouelle (1819) also had its roots in this era.

Carl F. Fallén (1764-1830), who specialized in Swedish Diptera, recognized *Ocyptera* and *Tachina* as distinct from *Musca* (1810, 1817, 1820-1825) but was otherwise rather conservative in his approach. He eventually described 133 species of *Musca*, dividing them into major groups according to the course of vein M (Fallén, 1817, 1820): M with an angular bend forward towards vein R<sub>4+5</sub>, M with a rounded forward curvature, or M running straight to the wing-margin. This is, of course, a very useful division, but Fallén's distinction between *Tachina* and *Musca* was very woolly so that his *Musca* still included many modern Tachinidae, Sarcophagidae, and Calliphoridae as well as *Musca domestica* and its allies; he placed *Musca tempestiva* in the group with more roundedly curved M.

The great J. W. Meigen (1764-1845), the founder of European dipterology, gave a preliminary review of genera (Meigen, 1803) in which he separated *Musca* from *Miltogramma*, *Anthomyia*, and *Stomoxys*, but still included in it the species *domestica*, *carnaria* (subsequently removed to his *Sarcophaga*), and *caesar*, etc. Shortly after this he critically reviewed (Meigen, 1804) the classifications of Linnaeus, De Geer, Fabricius, and Latreille and also discussed new characters in the venation, antennae, legs, etc, that he had discovered (Meigen, 1804, 1818). His definition of the family "Muscides" (Meigen, 1824: xi-xii) is equivalent to the genus *Musca* of his predecessors, and he divides the old "Musca" up into a large number of genera (1826: ix). Genera in which vein M is curved forward towards vein R<sub>4+5</sub> (*Idia*, *Mesembrina*, *Sarcophaga*, *Dexia*, *Musca*) form one group, and genera in which vein M is straight (*Anthomyia*, *Coenosia*, etc) form another. Indeed this division, later formalized as "Muscidae" and "Anthomyiidae," lasted until 1894 when Girschner discovered that the presence of meral (hypopleural) bristles would separate the tachinoid families from the muscoid families, and *Musca* (Muscidae) was definitively separated from *Lucilia*, *Calliphora*, and *Pollenia* (Calliphoridae) (see also Collin 1948). *Musca* was defined by Meigen (1826: 49) as follows, a definition not very different from his early (1803) diagnosis:

*Fühler aufliegend, dreigliederig: das dritte Glied verlängert, stumpf, prismatisch zusammengedrückt: an der Wurzel mit einer gefiederten Rückenborste. Hinterleib eirund, borstig. Flügel halb offen, mit einer Querader an der Spizze.*

*Musca* was now a very much more restricted taxon, comprising the modern Calliphoridae (except Rhiniinae) and Muscidae with a forwardly-curved vein M. Meigen recognized two subgroups, one of shining green unmarked flies (A) and one of differently-coloured flies with striped mesonotum (B); each was further subdivided into those with an angular bend to vein M (a) and those with a rounded curve (b). *Musca domestica* and its relatives were placed in group Ba, together with most of the modern Calliphoridae, whilst *Musca tempestiva* was assigned to group Bb, alongside the present genera *Morellia* and *Muscina*.

Meigen's work was enormously influential and was followed by most of his contemporaries and successors. Wiedemann (1830) gave no diagnosis of *Musca*, but followed Meigen's generic arrangement and the groups within *Musca* for his own work on the non-European species. Two important checklists of British insects followed Meigen (Stephens, 1829a, 1829b; Curtis, 1831, 1837), and his was the classification adopted in faunistic lists and reports: Brullé (1832), Haliday (1833), Eversmann (1834), Bouché (1834), Curtis (1835), Wilson (1835). Even Latreille (1829) was modified in English translation to incorporate Meigen's genera as subgenera (Griffith, 1832).

Two other figures were important during this period, both Frenchmen: the brilliantly eccentric A. J. B. Robineau-Desvoidy (1799-1857) and the more pedestrian but conscientious J. Macquart (1778-1855).

Robineau-Desvoidy's youthful and genial *Essai sur les Myodaires* was completed when the author was in his mid-twenties but was not published until it had been submitted to a protracted series of committees. It is a long and complex work, characterized as "impenetrable" by Brullé (1832). Even before its publication, Latreille (1829) criticized it on a number of grounds: the use of imprecise characters, the formation of groups based on habits, colour and other vague features, the re-naming of previously established groups. Others found the proliferation of genera and the huge number of species unacceptable and criticized the description of almost every species as new, without consideration of the preceding work by Fallén and Meigen. The *Essai* is a monograph of the Muscomorpha (Schizophora): of its 813 pages, some 40% are devoted to our modern Tachinidae and 75% to the Calyptrata. What Meigen treated as his Muscidae with vein M curved forward (the 19th century "Muscidae") comes at the end of Robineau-Desvoidy's 1st division Calypteratae (section Sarcobiae & Coprobiae, "families" Theramydidae (in the Vivipara) and Muscidae (in the Oviparae)), and includes 46 genera where Meigen had 5. Meigen's Muscidae with vein M straight (the 19th

century "Anthomyiidae") becomes the entire 2nd division Mesomydae and includes 44 genera where Meigen had 6.

Interpretation of Robineau-Desvoidy's taxa is bedevilled by the almost total destruction of his collection and it was only in the early decades of this century that most of his genera were recognized and either brought into use or assigned to synonymy. With prolonged study it is possible to interpret most of his work in a rational and logical way.

One section of his Muscidae is termed "Muscidae armentariae," defined by features of the antennae, arista, head-profile, proboscis, and venation. This consists of four genera, which are all now included within *Musca*. It has to be admitted that their definition is poor and recognition is only practical when combined with the species descriptions and habits:

<i>Plaxemya</i> .	1 species.	Now recognized as a species-group within <i>Musca</i> .
<i>Byomya</i> .	3 species.	Also recognized as a species-group within <i>Musca</i> .
<i>Sphora</i> .	2 species, from Australia.	Both now synonymized with <i>Musca domestica</i> .
<i>Musca</i> .	13 species, European and exotic.	All now synonymized with <i>Musca autumnalis</i> and <i>Musca domestica</i> .

This is a very restricted view of *Musca*, and one which is not even accepted nowadays, though almost a century later Malloch and Townsend did attempt to restrict *Musca* to *domestica* and to raise the related segregates to generic rank.

Robineau-Desvoidy fell out with his contemporaries, especially his fellow-Dipterist Macquart, and became increasingly isolated and eccentric both scientifically and socially. He died prematurely, worn out by his selfless work as country doctor in an impoverished and unhealthy district and embittered by the rejection of his work by his scientific peers. They were unable to accept or even comprehend his many genera, though the more obvious segregates were picked out and used either as genera or as subgenera (e.g. *Lucilia*, *Pyrellia*, *Pollenia*, *Calliphora*).

Macquart (1834, 1835), by combining the best elements of Meigen and Robineau-Desvoidy together with his own insights, arrived at a concept of *Musca* (though not a definition) that is in no way different from that in use today. He gave a key to the genera of "sous-tribu *Muscines*" (1834: 9) that distinguishes *Musca* by having the arista haired on upper and lower surfaces, 1st posterior cell (cell r<sub>4+5</sub>) almost closed, and antenna long with basoflagellomere 3 times as long as pedicel. He accepted Robineau-Desvoidy's genera *Pollenia*, *Calliphora*, and *Lucilia* (now Calliphoridae), and assigned

“*Musca*” species with a smoothly curved vein M to a new genus *Curtonevra*. Writing of *Musca*, he stated “ce genre ... est arrivé ... à ne contenir que la Mouche domestique et quelques espèces voisines” (1834: 19). He gave a key to Robineau-Desvoidy’s segregates of *Musca*, perceptively noting the haired eyes and unusually hyaline wings of the species *vitripennis* Meigen (*Plaxemya* of Robineau-Desvoidy). He retained the species *tempestiva* Fallén in *Musca* though several later authors were to transfer it to *Curtonevra* because of the less angular forward bend of vein M.

Much of this material was repeated in 1835, though he there gave a slightly more elaborate description of *Musca* to include the head profile (epistoma not or little projecting) and mentioning that the forwardly-curved section of vein M is generally recurved (Macquart, 1835: 235).

In his final work, Meigen (1838: 290-291) discussed the subdivisions of *Musca* proposed by Robineau-Desvoidy and Macquart and went a little way to accomodating some of them. He used the names *Lucilia* and *Pyrellia* for his greenbottle subdivisions (Aa and Ab); he accepted *Curtonevra* (Bb) (with segregates: bare eyes and hairy eyes) and retained *Musca tempestiva* here; and he accepted *Pollenia* as a distinct segregate within *Musca*, which was otherwise retained in the broad sense of his 1826 work (Ba).

During the latter part of this period (1800-1840) there was no awareness that the principal domestic flies — the house fly (*Musca domestica*) and the bluebottles (*Calliphora* spp.) — might belong to separate groups. Although there was no sense of a formal “type-species” as we now understand it for the genus *Musca*, entomologists did have a feeling for a type or typical species, and both *Calliphora vomitoria* and *Musca domestica* were regarded as typical species of *Musca*.

### **Meigen to Stein (1838-1921)**

There followed a very long period during which few advances were made in the systematics of *Musca*. The genus had been refined to *domestica* Linnaeus and its immediate allies, and it was realized that the European fauna was very small. The interest of system-builders and describers alike centered on other groups and on the wealth of tropical forms being brought back to European collections by travellers and colonial settlers.

The most prolific describers of exotic species were J. Macquart, F. Walker (1809-1874) and J. M. F. Bigot (1818-1893). Macquart (1843) gave a tabulation of genera, which shows that he was still satisfied with his earlier definition of the genus, and he did indeed find that all the exotic species submitted to him (about 20; Macquart, 1843, 1846, 1851, 1855) “se rappro-

*chent plus ou moins de la mouche domestique ou de la mouche corvine".* Despite his tendency to describe the same species several times and to create homonyms, it is fair to say that Macquart's work was of a much higher quality, in terms both of his perceptions and his procedures, than that of Walker and Bigot. Walker (1849) used *Musca* in Meigen's very broad sense and incorporated Robineau-Desvoidy's (1830) divisions and genera as sub-groups of *Musca* into his attempt to arrange world material. Thus, within *Musca s. lat.* he had a section "*Muscidae armentariae*" in which he listed the generic names *Plaxemyia*, *Byomyia*, *Sphora*, and *Musca*, describing several new species within this restricted *Musca* but without understanding or offering any interpretation of the other Robineau-Desvoidy names. Walker re-iterated this broad concept of *Musca* in two subsequent papers (Walker, 1853a, 1853b) and followed it in all his descriptive papers. This concept was hardly progressive, since large numbers of *Musca* were described that are now scattered in the families Calliphoridae and Muscidae, and even Sarcophagidae and Tachinidae, and the original combination and description give little indication of the actual affinities of each new taxon. Bigot's work too contributed little to progress. In his major paper on the group (Bigot, 1888a), he discussed the families Stomoxydidae and Muscidae, the latter being the equivalent of Meigen's genus *Musca*. His key to the genera of Muscidae gave a taxonomic definition of *Musca* based entirely on Rondani (1856 - see discussion below), but he failed to clarify the identity of *Byomia* [sic] Robineau-Desvoidy, which he considered most likely a synonym of *Musca*, and he assigned *Plaxemyia* [sic] Robineau-Desvoidy to his Stomoxydidae.

In Europe, the writers of entomology treatises and textbooks followed Meigen, with Macquart's modifications. For example, Westwood (1840) listed 6 genera in his "*Muscies*," *Lucilia*, *Calliphora*, *Musca*, *Pollenia*, *Mesembrina*, *Curtoneura*. Blanchard (1840) had a less clear-cut concept of *Musca*, giving a diagnosis and including the three species *vomitoria* [*Calliphora*], *domestica* [*Musca*], and *meridiana* [*Mesembrina*]. Morris' (1865) list of British insects followed Meigen precisely.

J. W. Zetterstedt (1785-1874) worked for many years on the Diptera of Scandinavia. His first definition of *Musca* (Zetterstedt, 1838: 652) alluded to some 50 Scandinavian species. Of the 31 enumerated in this work, several belong to the modern Tachinidae, Rhinophoridae, and Calliphoridae. His family "Muscariæ" (Zetterstedt, 1842: 5) was equivalent to Meigen's genus *Musca* and he recognized six genera: *Lucilia*, *Pyrellia*, *Musca*, *Mesembrina*, *Cyrtonoeura*, *Cinochira* (Zetterstedt, 1845: 1312). This is based on Macquart. He defined several subdivisions of *Musca*: thoracic dorsum tomentose (*Pollenia*), thoracic dorsum not tomentose and palpi yellow [various Cal-

liphoridae], and thoracic dorsum not tomentose and palpi black [*domestica*, *corvina*, and the Calliphorid *azurea*]. He assigned the species *Musca tempes-tiva* to *Cyrtoneura* because of its more roundedly curved vein M.

The work of Camillo Rondani (1807-1879), which was concerned mainly with Italian Diptera, marked a real step forward in the systematics of the Muscidae and Anthomyiidae. Not only was his method excellent — keys to genera and species, descriptions of all taxa, and type-species listed for the genera — but he discovered new and significant characters on which to base his re-classification of the groups. His treatment of *Musca* illustrates this. His key diagnosis of the genus (Rondani, 1856: 89) gives the following combination of characters: mid tibia without an internal (i.e. ventral) bristle, and 5th longitudinal vein (vein M) with an angular forward curve. This is precisely the definition still given for the genus (e.g. Hennig, 1963: 905; Emden, 1965: 39). In fact, Rondani recognized two genera where we would now recognize one: eyes bare (*Musca*, 4 species) and eyes haired (*Plaxemya*, 2 species). In revising the species (Rondani, 1862: 220) he correctly included *tempestiva* in the genus *Musca* rather than in *Cyrtoneura*.

The Austrian I. R. Schiner (1803-1873) produced a work that became the standard manual on European Diptera until well into our own century (Schiner, 1862). He based his treatment of this group on that of Rondani and defined *Musca* as having the arista densely plumose on both surfaces, vein M with an angular forward bend towards vein R<sub>4+5</sub>, and mid tibia without bristles inside (Schiner, 1862: lxxxvii). He did not accept Rondani's division of the genus into two, but defined two species-groups that were in any case different from Rondani's *Plaxemya* and *Musca*: hind cross-vein (dm-cu) closer to the bend of M than to small cross-vein (r-m) (*domestica*, *corvina*), and hind cross-vein (dm-cu) midway between the bend of M and small cross-vein (r-m) or closer to the latter (*vitripennis*, *phasiaeformis*, *tempes-tiva*). This work was followed by many European workers for their own local faunas, e.g. Kowarz (1873), Verrall (1888, 1901b).

It is worth pointing out here that details of the venation still confused taxonomists, in particular the elbowed or rounded forward bend of vein M. The eremophilic Old World *Musca lucidula* was described by Loew (1856) as a species of *Cyrtoneura*.

In an uncompleted posthumously-published work Robineau-Desvoidy (1863b) returned to the genera of the "higher" Muscidae. He retained his division "*Muscidae armentariae*" with the three genera *Plaxemya*, *Byomya*, and *Musca*, giving diagnoses that are scarcely different from those of his 1830 work. He recognized one species of *Plaxemya* (*vitripennis* Mg.), 3 species of

*Byomya* (all synonyms of *tempestiva* Fallén), and 15 species of *Musca* (all synonyms of *autumnalis* and *domestica*, except for one subsequently transferred to *Pollenia* (Calliphoridae)). His system was largely followed by Lioy (1864), who produced a synthetic work, without originality and apparently in ignorance of most of the work of his fellow-countryman Rondani. Lioy's genera (*Musca*, *Plaxemya*, *Byomya*) and their diagnoses were taken directly from Robineau-Desvoidy's work. When asked for his verdict on Lioy's work (and his reason for ignoring it), Rondani replied that "c'est de la poésie."

Following from Schiner, the Austrians F. M. Brauer and J. E. von Bergenstamm dealt comprehensively with the European Tachinoidea, together with a large number of exotic forms, and their work also covered the groups of Muscidae with a forwardly-curved vein M. Their interpretation of *Musca* and allied genera was naturally based on that of Rondani and Schiner and they accepted Rondani's separation of *Musca* into two genera, though any merit in this division was obfuscated by inaccuracies in their differentiation (Brauer & Bergenstamm, 1889: 156):

Eyes bare, almost touching in ♂, separated in ♀, etc.

*Musca* L. s. str. Rdi., *domestica* L.

Eyes haired in ♂, pubescent in ♀. Frons of ♂ moderately broad, very broad in ♀, etc.

*Placomyia* R.D. emend. (*Plaxemya* R.D.) *vitripennis* Mg.

This distinction was maintained to the end of their work (Brauer & Bergenstamm, 1893: 179) and the authors also included a check list of the species that they studied (Brauer & Bergenstamm, 1891: 419), which was useful insofar as it also indicated some of the early type/syntype material located in the Vienna Museum.

The last great systematic work of the century was by the Frenchman L. Pandellé (1824-1905) and comprised a monograph of the "Muscidae" (i.e. both calyprates and acalyprates) of France. He introduced a number of new characters from the chaetotaxy of the body, wings, and legs into the taxonomy of the Schizophora and also pioneered the study of male genitalia. Independently of his contemporaries, he discovered the value of the chaetotaxy of the body and legs, but the prolixity of his text and absence of any illustrations deprived him of any justly-earned credit for these findings. Pandellé (1898a: 20-23) gave a key to the genera of his division "Muscées" that was quite different from that of his immediate predecessors, using a wealth of characters that were mostly new. His "Muscées" included *Mesembrina*, *Pyrellia* (in a very expanded sense), *Musca*, *Stomoxys*, *Gymnoidea*, *Graphomya*, and *Muscina*. His genus *Musca* (Pandellé, 1898b: 40) is entirely modern in its scope and with many features in the generic description that

were independently rediscovered by the next generation of taxonomists. Four species were included: *vitripennis*, *tempestiva*, *corvina*, and *domestica*.

Pandellé (1894, 1896) and Girschner (1894, 1896, 1897) were both influenced by the work on chaetotaxy of Osten Sacken (1881b, 1884) and independently arrived at the conclusion that muscids with an angular vein M and bare meron belonged to the Anthomyaires/Anthomyidae and not to the Tachinaires/Tachinidae as previously believed. In retrospect, it is apparent that Pandellé's proposals for the classification of the Schizophora were superior in several respects to those of Girschner. Nevertheless, it was Girschner's views, expressed succinctly and accompanied by clear and schematic drawings, that had the greater influence on 20th century dipterology.

The greatest stimulus to the work of the early part of the 20th century came from the biological investigations by the Russian J. A. Portschinsky (1848-1916). In an important pioneering paper published in 1885 (abstracted and discussed in German by Osten Sacken, 1887), Portschinsky drew attention to an apparent dual reproductive strategy in *Musca corvina* [i.e. *autumnalis*]: although normally laying some 24 large eggs, females in the south of Russia (Crimea and Caucasus) incubated a single large egg and laid a 3rd instar maggot. It is now clear, of course, that two species were involved here, as Portschinsky himself subsequently recognized (Porschinsky, 1910). He continued with his investigations of oviparous and viviparous Muscidae and of their coprophagous, saprophagous, and carnivorous larvae (Porschinsky, 1892, 1910, 1911), and his later work was reviewed by Townsend (1911). There is no doubt that this was enormously influential in alerting field entomologists to the biological interest of this group, at a time when the medical and veterinary importance of *Musca*-flies was increasingly being recognized and this gave a tremendous impetus to the fundamental work on the group that followed during the 1910s and 1920s.

Throughout this period (1885-1920) the dominant figure in the taxonomy of the Muscidae was the German Paul Stein (1852-1921), who specialized almost exclusively on this group and who single-handedly brought order to the chaos into which the group had sunk during the course of the 19th century. From the modern standpoint, Stein's concept of genera seems rather limited, but his grasp of species was remarkable. His papers were methodical and comprehensive and almost invariably contained keys; he travelled to the centers of European DipteroLOGY in Germany, England, Sweden, and France, to study the types of earlier authors; and, towards the end of his life, he was able to produce a key to all the European Muscidae (Stein, 1916), a key to the world genera (Stein, 1919), and catalogues of the world species (Stein,

1919; in Bezzi & Stein, 1907). Stein was most interested in the “Anthomyiidae” (in the 19th century sense) and so *Musca* and allied genera were rather peripheral to the main thrust of his work. He followed Schiner (1862) in his concept of the genus *Musca*. Reviewing extensive collections from the Afrotropical and Oriental regions, he was the first to realize how large and diverse the genus might prove to be (and how very slipshod and inadequate for the most part his precursors in tropical Muscidae had been). For the first time (Stein in Bezzi & Stein, 1907) the Palaearctic *Musca* species were catalogued, with an inventory of recognized species, *species incertae*, and synonyms, a total of 46 names.

The first two decades of the century also saw the initiation of several other strands of work on *Musca* which did not come together until the work of Patton and Malloch in the 1920s. The American tachinid specialist C. H. T. Townsend was always interested in the importance of reproductive biology for systematics (e.g. Townsend, 1913) and on several occasions he briefly turned his attention to the genus *Musca*. He was an advocate of the concept of restricted genera and he realized from Portschinsky’s account of *Musca corvina* [*autumnalis*] that the reproductive biology of this species was very different from that of *Musca domestica*. He erected the new genus *Eumusca* for this species (Townsend, 1911). A few years later, after studying published work on the Indian *Musca* species, he erected the genus *Viviparomusca* for the viviparous species *bezzii* Patton & Cragg (Townsend, 1915) and another genus *Awatia* for the species *indica* Awati (= *planiceps* Wiedemann) because it was a larviparous member of the *Philaematomyia*-group and had well-developed presutural dorsocentral setae.

Among blood-sucking Muscidae collected in the first decade of the century were species of *Musca* that could abrade vertebrate skin and suck blood, but at first it was not realized that these were true *Musca* rather than biting Stomoxyinae. Austen (1909) described a new genus *Philaematomyia* (for a species subsequently found to be identical with *crassirostris* Stein) which he characterized as “a remarkable connecting-link between the ordinary non-biting Muscinae and Stomoxydinae.” He noted that the powerful proboscis could not pierce vertebrate skin but that the flies used the enormous prestomal teeth to cut the skin and then sucked up blood. He did not consider it to be particularly related to *Musca*. Brunetti (1910) described a new genus *Pristirhynchomyia* (for a species subsequently found to be identical with *conducens* Walker) which he considered transitional between *Philaematomyia* and *Musca*: he excluded both genera from the Stomoxyinae, though without giving any reasons. He noted that his species was also a “biting fly,” unable to pierce the skin but able to enlarge existing wounds or sores.

It has been gradually realized that these features in biology and structure are progressive and that various evolutionary stages are represented within the genus *Musca*, as they are in other genera of the Muscini. The fragmentation of the genus on the basis of a few autapomorphies results in a series of typological units rather than monophyletic groups and does not reflect the evolutionary history of the genus.

Work in India by Awati (1915-1920) and Patton & Cragg (1912, 1913a, b) increased the number of species and paved the way for advances made in the 1920s. Awati's work was particularly important, since he looked in such detail at a wide range of morphological, anatomical, and biological features, but the lack of any reference material from his work has hampered the recognition of the actual species that he discussed: indeed, it is still not possible to interpret all the names that he proposed. Patton did an enormous amount of fieldwork and breeding of *Musca* species in South India and this bore fruit when he returned to England and began extensive studies on the taxonomy and systematics of *Musca*: he was able to combine his first-hand knowledge of reproductive strategies, biology, adult feeding habits, etc, with his systematic researches. Both authors used *Musca* in the broad sense defined by Stein.

### The Modern Era (1921-1992)

A conflict between generic "lumpers" and "splitters," familiar in most groups of insects, developed during the 1920s, but a careful study of various character-suites among the species of *Musca* and a better understanding of the evolutionary trends within the genus soon gave rise to a delimitation of the genus which has remained stable now for 60 years. The various splinter genera defined during the 1920s are recognized at the level of subgenera or species groups.

The period up to 1937 was dominated by the Briton W. S. Patton (1867-1960). Through his first career in the Indian Medical Service, Patton developed an interest in all the higher flies of medical and veterinary importance (especially Muscidae and Calliphoridae) and these early years of practical work underpinned all his subsequent systematic research. His correspondence and exchanges with other muscoid workers of the day (E. E. Austen, J. Villeneuve, J. M. Aldrich, and especially M. Bezzi) showed him that the same species were often masquerading under several different names and that the names themselves were often applied in a quite haphazard and inconsistent manner. Believing that he understood the species and species limits sufficiently well, he studied the types and other material in European museums and private collections (Patton, 1922a, 1923a, 1923b, 1925). From

this basis he proceeded to publish revisions of the Oriental species (Patton & Senior-White, 1924; Patton, 1937a, b), Afro-tropical species (Patton 1926, 1936), and Palaearctic species (Patton, 1933b, c). He also undertook more fundamental comparative studies of the male genitalia (Patton, 1932, 1933a), the antennae (Patton & MacGill, 1925), the evolution of the bloodsucking habit (Patton, 1934), and supervised a comparative study of the female ovipositor (Ho, 1938).

Patton was convinced that *Musca* as he understood it (and as Stein had understood it, and as the present generation still understands it) was an indivisible monophyletic unit. He put forward characters to support this contention, such as the autapomorphic presence of a large sensorium at the base of the antennal basoflagellomere (Patton, 1932) and also adduced arguments to show that the more striking biological features in *Musca* (such as viviparity and modifications in the female reproductive system, adaptations in the male genitalia, biting and bloodsucking) are progressive and, in the case of the blood-feeding, have arisen more than once within the genus. He recognized only three groups in the genus, the *domestica*-group, the *sorbens*-group, and the *lusoria*-group. In discussion he was able to persuade Bezzi that this was the right course, though Bezzi had earlier joined in the fragmentation process begun by Townsend. He dismissed Townsend's splinter genera with derision. But he could not reach any accord with Malloch and the two remained irreconcilable. It is clear now that even Patton misidentified a great deal of material: throughout his long career he re-identified, mis-identified, and changed his mind about species limits and names and his keys are generally poor, based on rather weak characters and difficult to use. Malloch too misidentified species. It is obvious that much of their argument over detail can be resolved once it is known what species they actually meant by particular names: for instance, both had dogmatic views over where "*planiceps* Wiedemann" should be placed, yet neither correctly identified this species and, as they were probably not even discussing the same species, their discord appears completely futile. On a higher systematic level, the argument resolves itself over the ranking of hierarchies, whether as species groups (Patton) or genera (Malloch). The present system, largely worked out by Emden (1939, 1965), represents a synthesis of the work of Patton and Malloch.

The Italian Mario Bezzi (1868-1927) worked on *Musca* at various stages of his career, mainly describing new species. He gave a key to all the Mediterranean and African species, and recognized a number of subgroups (Bezzi, 1911). Turning to the blood-sucking forms (Bezzi, 1921) he showed that the species of *Musca sensu* Stein were now distributed among two

subfamilies, *Philaematomyiinae* (*Philaematomyia*, *Pristirhyncomyia*, *Ptilolepis*) and *Muscinae* (5 genera). He subsequently (Bezzi, 1923) described another subgenus, *Lissosterna*, for the desert species *Musca albina* Wiedemann.

The work of the Scottish-American John R. Malloch (1875-1963) advanced knowledge of the family Muscidae to a quite remarkable degree and in two studies of the genus *Musca* (1924, 1925; further notes in 1929b, 1932) he showed the innovative approach and remarkable ability to perceive new characters that are evident in all his work. He discovered several valuable features for identifying and classifying species of *Musca*, some of which were adopted, albeit reluctantly, by Patton. He too was uncertain how to rank the groups that he could define: "It is difficult to assign a definite status to the segregates herein accepted, for, while some workers would consider them as valid genera, others would just as certainly rank them at most as subgenera, or even refuse to accept them as named divisions" (Malloch, 1924: 132). He proceeded to list them as genera (1924), then as subgenera (1925) but with *Philaematomyia* and *Ptilolepis* as separate genera, and then finally as full genera again (1929b), recognising a total of 8 genera (*Musca*, *Byomyia*, *Lissosterna*, *Plaxemya*, *Philaematomyia*, *Eumusca*, *Viviparomusca*, *Ptilolepis*). This work was rejected by Patton, who regarded it as a falsification of the true relationships within *Musca*. Since both were handling in part misidentified material, it was not possible for them to find any common ground at all, though subsequent work has acknowledged merit in both approaches.

Two contrasting views of *Musca* appeared in 1937. Townsend summarized his views on the genera of the Muscini and divided *Musca* into eleven genera: *Awatia*, *Byomyia*, *Eumusca*, *Lissosterna*, *Musca*, *Philaematomyia*, *Plaxemya*, *Pristirhynchomyia*, *Ptilolepis*, *Sphora*, and *Viviparomusca* (Townsend, 1937). Since these segregates were based on the study of a very few species and were founded often on biological or anatomical features, they have not been accepted by anyone. In the same year, Séguay (1937) produced a volume for *Genera Insectorum* that gave a catalogue of World Muscidae. He followed Malloch's (1925) concept of *Musca*, but ranked the blood-sucking *Ptilolepis* and *Philaematomyia* as distinct genera. For these three genera he catalogued a total of 92 species, with 164 names (valid and synonymous). The 92 species include a number that are no longer recognized in *Musca*, *species incertae*, misidentifications, and names that are now synonyms, whilst several names that do in fact belong to *Musca* were overlooked by Séguay.

The most significant work since that time has been done in England by F. I. van Emden (1898-1958). He gave a most sympathetic account of the work by Patton, Malloch, and Ho (Emden, 1939: 73-76) together with a comprehensive key to the Afrotropical species and pointed out that there was indeed no serious discrepancy between Patton's, Ho's, and Malloch's classifications of *Musca*. Apart from a very few anomalous, perhaps misidentified, species and the ranking of the categories, the three systems agreed very well. Emden was thus able to maintain *Musca* in the traditional sense and, by elaborating the characters discovered by Malloch and other authors, to define a series of groups within the genus, several of which are monophyletic. Where existing names were available for these groups, Emden used them as subgenera, thus recognising 7 named categories. In his treatment of the Oriental species (Emden, 1965) he adopted the same arrangement and recognized the same number of groups (subgenera) as had Malloch, though his definitions are far more comprehensive and sophisticated. It was only with the work of Emden that species-limits and species-nomenclature reached a level generally felt to be satisfactory though there are still areas (for example, the Oriental species of the *Viviparomusca*-group) where further work is required.

Most of the other published work over the last 40 years has been descriptive (more closely defining the known species and describing new species) and clarificatory (studying the types of older authors and elaborating distributions). Authors have adopted the same consensus view of *Musca* and, for the most part, of its subgroups. A different view of the subgroups was put forward by Zielke (1971c), who recognized only two groups, the *lusoria*-group and the *sorbens*-group (the latter including Patton's *domestica*-group). However, these groups were not defined, except in the most general way, nor did Zielke give any indication as to which species he assigned to each group. Hennig (1964: 978-982) sought to establish which of the subgroups of *Musca* were monophyletic and he used the results of Patton and Ho for his discussion of character-transformations. He too reduced the number of subgroups that should be recognized in the genus.

As a postscript, reference should be made to the contribution of larval studies to *Musca* systematics. The pioneer work by Thomson (1937, 1947) and Zimin (1948) was summarized by Emden (1965), who was able to list a larval autapomorphy to support the conclusion that *Musca* is a monophyletic genus: mouth-hooks asymmetrical, the left one shorter than the right one. A detailed treatment of the immatures has recently been given by Skidmore (1985), and a comparative account of the family by Ferrar (1987).

### The type species of the genus *Musca* Linnaeus

As mentioned above at the conclusion of section 2, it was generally considered in the early part of the 19th century that the domestic flies *Calliphora vomitoria* and *Musca domestica* were the typical species of the genus *Musca*. Several examples have already been cited. The following interesting cases deserve to be quoted here.

The first formal statement of a species acting as type of *Musca* ("espèce qui ... sert de type") was by Latreille (1810), who named it as *Musca vomitoria* sensu Fabricius (i.e. *Calliphora vomitoria* Linnaeus: Calliphoridae). Curtis (1835) gave *Musca caesar* as the type (i.e. *Lucilia caesar* Linnaeus: Calliphoridae). Macquart (1834) had effectively restricted the genus to *domestica* and its immediate allies, but Westwood (1840) was the first formally to designate *domestica* L. as the type of *Musca*.

Now, one can argue indefinitely about what these authors meant by "type" and also what their interpretations of the various cited taxa may have been, but it remains a fact, as Townsend (1915) pointed out, that the first valid citation of a type species for *Musca* is of a species now placed in the family Calliphoridae. Townsend (1915) followed the logical consequences of this, prescribed by the existing rules of nomenclature: he transferred the generic name *Musca* to the Calliphoridae, to replace *Calliphora*, and proposed the new genus *Promusca* for *domestica* and its allies. It took 10 years of discussion and an ICZN Opinion (ICZN, Opinion 82, 1925) to uphold the common-sense view that *domestica* should be type species of *Musca* and thus to ratify a binomen that is almost as well known as that of *Homo sapiens*.

### The genus *Musca* Linnaeus

*Musca* in current status consists of 66 recognized species and 7 subspecies, for which 221 species-group names, valid and synonymous, have been created. Key works on the genus are: New World (Vockeroth, 1953); Palaearctic (Hennig, 1964); Japan (Shinonaga & Kano, 1971); China (Fan, 1965); Oriental (Emden, 1965); Thailand (Tumrasvin & Shinonaga, 1977); Philippines (Magpayo, Shinonaga & Kano, 1987); Afrotropical (Emden, 1939, Zielke, 1971c); Australian (Pont, 1973); Micronesia (Snyder, 1965); immatures and biology (Skidmore, 1965); bibliography (West & Peters, 1973); and ecology, classification and biotic associations (Greenberg, 1971). Most of the above need to be updated by reference to the various regional taxonomic catalogues: Nearctic (Pont (in press)); Neotropical (Pont, 1972); Palaearctic (Pont, 1986a); Afrotropical (Pont, 1980); Oriental (Pont, 1977b); Australasian/ Oceanian (Pont, 1989).

## Workers: Their collections and publications

The preparation of a work such as this requires knowledge of the workers and their collections and publications. Some of this knowledge can be derived from the publications themselves, but most of it is scattered elsewhere. To aid future workers, we have identified our sources of ancillary information below and in the references cited. The standard entomological bibliographic sources (Hagen, 1862-63; Horn & Schenkling, 1928-9 (corrections, Gaedike & Smetana, 1978 & 1984); Derksen & Scheiding, 1963-75) were used along with other special sources such as Stafleu & Cowan (1979-88), which was extremely useful for dating and verifying various editions of the earlier general natural history works. Two other old bibliographies were useful for identifying cryptic citations in the earlier literature (Cobres, 1782 & Dryander, 1796-1800). The collection information in Horn and Kahle (1935-37) and its supplements (Sachtleben, 1961, Horn *et al.*, 1990) is not repeated below.

### **Bigot, Jacques Marie Frangile (1818-1893)**

Gilbert (1977: 33, 13 citations), Brauer (1899, collection notes); Maa (1963, notes on labels); Ackland & Taylor (1972, notes on types at Oxford); Crosskey (1971, collection, label & type notes), Thompson (1989, collection, label & type notes).

### **Bouché, Peter Friedrich (1784-1856)**

Gilbert (1977: 43, 1 citation). Remnants of his Diptera were acquired by DEI in 1937; Sachtleben (1944, biographical sketch).

### **Curtis, John (1791-1862)**

Gilbert (1977: 81, 9 citations), Blackwelder (1947, on British Entomology); Neboiss (1963, notes on Curtis Collection, now in Melbourne (MVM)).

### **De Geer, Carl (1720-1778)**

Gilbert (1977: 88, 11 citations); Clausen (1977: 514, data on curation, style of pins, and labels used by De Geer); Gurney (1956, last word on proper form of De Geer's surname); Persson, Pont & Michelsen (1984, notes on collection and nomenclature). De Geer's work was abstracted by Retzius (1783) following the Linnaean system.

De Geer began his work before Linnaeus formally elaborated his system of binomial nomenclature and consistently applied it to animals. Hence,

the names in the early volumes of the *Mémoires* are not binominal. Traditionally, however, workers have accepted the names in volume 3 onward as conforming to the Linnaean system (Persson et al. 1984: 91). Unfortunately, under the current rules of nomenclature (ICZN, 1985), De Geer does not consistently follow the principle of binominal nomenclature even in the subsequent volumes. De Geer used binary nomenclature, a single word for the genus, and, while he usually used one word for the species, sometimes a two-word phrase was used for the species name. In the genus *Musca*, he used the adjectives *major* and *minor* to distinguish between large and small varieties (*Musca minor domestica* and *M. domestica major*; *M. major larvarum* & *M. minor larvarum*; *M. vivipara major* & *M. vivipara minor*), but sometimes color and habitat descriptors were used (*Tipula nigra aquatica*) or habitat and structural descriptors (*T. Agarici seticornis*) or structural and pattern descriptors (*T. clavata maculata*). These two-word combinations are phrases, not separate words that together refer to a single entity as permitted under the rules (ICZN, Art. 11(h)v). However, for pragmatic reasons, we have followed tradition and accepted all two-word names as binominal and available, but have rejected as polynominal and unavailable all names of three or more words.

**de Villers, Charles Joseph (1724–1810)**

Gilbert (1977: 396, 3 citations). Collections stated to belong to “Tissier jr. (Lyon)” with duplicates in “Städt. Mus., Lyon” (Horn & Kahle 1936: 289), but we have been unable to verify this. No de Villers material is known to exist in the Musée Guimet d’Histoire Naturelle, Lyon, but, on the other hand, nothing is known of the collection’s fate nor is it known definitely to have been destroyed (J. Clary, *in litt.*, 30 Oct. 90).

**Doleschall, Carl Ludwig (1827–1859)**

Gilbert (1977, 3 citations); some material collected by Doleschall is present in NMW and may represent syntypes of some of his species (Osten Sacken, 1881a).

**Donovan, Edward (1768–1837)**

Gilbert (1977, 2 citations); Weiss (1927, fate of collection). Horn & Kahle (1935: 60) state that Donovan’s collection was auctioned by J. C. Stevens in 1818. Details about the auction can be found in Chalmers-Hunt (1976a: 5, 1976b: 77), and a copy of the sale catalog exists in BMNH where some of the specimens ended up, including material used for Donovan (1805). Other Donovan material ended up in the Macleay Museum, Sydney (Horning, 1988).

**Drury, Dru (1725–1803)**

Gilbert (1977, 9 citations); Weiss (1927, notes on fate of collection). Some Drury material was acquired by Alexander Macleay and is now in Sydney (Horning, 1988). Many specimens were purchased by F. W. Hope and should still be in UMO (*Hope marginalia*). Hayek (1985) provided additional information on the fate of Drury specimens.

**Fabricius, Johann Christian (1745–1808)**

Gilbert (1977, 21 citations); Zimsen (1964, type material); Tuxen (1967a & b, biography & bibliography); Michelsen (1979a, muscoid spp.); Pape (1986, sarcophagid spp.); Evenhuis (1988, Labillardière material).

**Fallén, Carl Fredrik (1764–1830)**

Gilbert [no data]; Ander (1952, collection notes); Persson 1964 (notes on types); Michelsen (1983, anthomyiid spp.); Pont (1984, muscid spp.); Pape 1986 (sarcophagid spp.).

**Forster, Johann Reinhold (1729–1798)**

Gilbert (1977, 5 citations), Day & Fitton (1977, type material).

**Geoffroy, Etienne Louis (1727–1810)**

Gilbert (1977: 135, 3 citations). His principal work (1762) was ruled as non-binomial by ICZN (1954). The Diptera genus-group names were subsequently validated by ICZN (1957), and his species are available from their appearance in Fourcroy (1785). His collections were stated to be deposited in "Mus. Autun" "via de Laplanche" (Horn & Kahle 1935: 88), but several inquiries to the Musée d'Histoire Naturelle, Autun, have remained unanswered.

**Gimmerthal, Benjamin August (1779–1848)**

Gilbert (1977: 140, 1 citation). Types in Riga.

**Gravenhorst, Jean Louis Charles (1777–1857)**

Gilbert (1977, 2 citations). His collection was deposited at Wroclaw, but the Diptera types are now lost among the general material or destroyed (M. Kak, *in litt.*; also Kinel, 1957).

**Gmelin, Johann Friedrich (1748–1804)**

Gilbert (1977: 141, 2 citations); Kabat & Petit (1988, editions of 1790); Spilman (1967, nomenclatural importance of 1790 work); Vane-Wright (1975, definitive treatment of 1790 work and collections). Insect types from the Leskean Museum have been identified in NMI, but none of the Diptera survive (J. P. O'Connor, *in litt.*, 28 Sept 90).

**Haliday, Alexander Henry (1807–1870)**

Gilbert (1977: 156, 9 citations); Nash & Chandler (1978, general notes on material & types).

**Harris, Moses (1730-between 1787–1789)**

Gilbert (1977: 160, 2 citations); Goffe (1946, 1951, syrphid names, notes on works); Lisney (1960, biographic and bibliographic notes); Weiss (1926, biography); Hagen (1861, bibliographic note); Pont & Michelsen (1982, muscoid names, general notes).

**Harris, Thaddeus William (1795–1856)**

Gilbert (1977, 21 citations); Johnson (1925, notes on collection); Wilkinson (1980, notes on papers). The Harris collection is now in MCZ, where it is kept as a separate collection along with Harris' notebooks.

**Linnaeus, Carl von (1707–1778)**

Gilbert (1977: 227, 41 citations); Jackson (1890, history of collection); Bryk (1949, types in De Geer Collection); Lindroth (1957, work, collections); Day & Fitton (1978, re-curation of collection); Pont (1981, muscoid spp.); Thompson *et al.* (1982, syrphid spp.); Mikkola (1983, Linnaean insect pins); White (1987, tephritid spp.).

**Macquart, Justin Pierre Marie (1778–1855)**

Gilbert (1977: 241, 6 citations); Macquart (1850, list of material in Lille); Brauer (1899, collection notes); Maa (1963, notes on labels); Pont (1967, Australian muscid spp.); Ackland & Taylor (1972, notes on types at Oxford); Crosskey (1971, collection, labels & type notes); Lopes (1975, type notes); Thompson (1989, collection, label & type notes).

**Meigen, Johann Wilhelm (1764–1845)**

Gilbert (1977: 251, 11 citations); Steyskal (1974, translation of biographic materials); Herting (1972, tachinid types); Pont (1986b, extensive notes on Meigen's collection and collectors); Morge (1975–76, Meigen's color plates).

**Müller, Otto Friedrich (1730–1784)**

Gilbert (1977: 268, 4 citations); Henriksen (1922, 1940; biographic & bibliographic notes); Anker (1950, on *Zoologia Danica*). According to Hagen (1844: 131; also Horn & Kahle, 1936: 184), the collection of Müller was destroyed when the English fleet under Admiral Nelson bombarded Copenhagen in 1801. However, Anker in her review of Müller makes no mention of this destruction, merely stating that "posthumous scientific belongings were scattered by being sold at a public auction" (Anker, 1950: 86).

**Panzer, Georg Wolfgang Franz (1755–1829)**

Gilbert (1977: 287, 7 citations); Saunders (1888, index). Panzer's collection was given to ZMHU and the material was integrated into the general collection. We do not know of any modern worker who has been able to identify authentic Panzer type material in ZMHU. Some of the species illustrated in his *Favnae insectorum germanicae* came from other collections and, hence, may be found in other places.

**Poda von Neuhaus, Nicolaus (1723–1798)**

Gilbert (1977: 300, 1 citation); Schiner (1856, critical revision of Diptera species in 1763 work). Nothing is known of the collections of Poda. We wrote to the Steiermärkisches Landesmuseum, Graz, and were informed that despite extensive searches for Poda's collection no information about its existence had ever been discovered and the collection is assumed to have been destroyed (J. Madler *in litt.*, 9 July 92).

**Preyssler, Johann Daniel Edvard (17??–post 1838)**

Gilbert [no data]. Rozkosny et al. (1982, notes on Diptera).

**Robineau-Desvoidy, André Jean Baptiste (1799–1857)**

Gilbert (1977: 319, 7 citations); Royer (1931, notes on collection); Sabrosky (1974, defence of). What remains of the Robineau-Desvoidy collection is now at MNHNP, but most of it is known to have been destroyed. A check has been made in the remnants of his collection for the species enumerated in this paper, but nothing survives.

**Rondani, Camillo (1807–1879)**

Gilbert (1977: 322, 16 citations); Guigla (1957, types in Genoa); Sabrosky (1961, bibliographic notes); Hennig (1961, muscid types); Pape (1988, notes on collection & sarcophagid types).

**Roser, Karl Ludwig Friedrich von (1787–1861)**

Gilbert (1977: 324, 1 citation). Roser material is now in the museum at Ludwigsburg (SMN).

**Rossi, Pietro (1738?–1804)**

Gilbert (1977: 324, 3 citations); Baccetti (1962, biographic & bibliographic notes). Horn and Kahle (1936: 231) stated that the Rossi collection passed to Frédéric de Sanvitale of Parma and part thereof (or all) later passed via J. Hellwig to ZMHU in Berlin. Baccetti (1962: 20), however, noted that no Rossi types have been found in Berlin.

**Schrank, Franz von Paula (1747–1835)**

Gilbert (1977: 342, 2 citations). See Gistl, 1837b.

**Scopoli, Johann Anton (1723–1788)**

Gilbert (1977: 344, 7 citations); Higgins (1963, fate of collection and notes on plates for 1763 work); Schiner (1856, critical revision of Diptera species in 1763 work); Voss (1881, detailed information on life and collections); Guglia (1972 in Scopoli 1972, biography).

**Thomson, Carl Gustaf (1824–1899)**

Gilbert (1977: 381, 8 citations); Persson (1964, general notes on types; 1972, notes on "Eugenies Resa" material).

**Thunberg, Carl Peter (1743–1828)**

Gilbert (1977: 382, 8 citations); Karsholt & Schmidt-Nielsen (1986, collection notes).

**Walker, Francis (1809–1874)**

Gilbert (1977: 400, 14 citations); Smith & Taylor (1964, types at Oxford); Pont (1966, type notes); Lopes (1975, type notes); Graham (1979, biography); Walker (1984, material in Melbourne); Thompson (1989, collection, label & type notes).

**Wiedemann, Christian Rudolph Wilhelm (1770–1840)**

Gilbert (1977: 414, 3 citations); Zimsen (1954, types in Copenhagen); Thompson (1989, collection, label & type notes).

**Zetterstedt, Johann Wilhelm (1785–1874)**

Gilbert (1977: 425, 5 citations); Ander (1952, collection notes); Persson (1964, notes on types); Michelsen (1985, anthomyiid spp.); Pape (1986, sarcophagid spp.).

## Systematic Database of *Musca* Names

### Scope

All names proposed in combination with the genus-group name *Musca* or now included in the genus *Musca* are reviewed here. This definition is broad because fly maggots were described as ascarid worms and fossil cockroaches were mistaken for flies! So some names proposed in combination with *Ascaris* (Phylum Nematoda) now are to be found as synonyms of *Musca domestica* and some names proposed in combination with *Musca* are now found assigned to species in the order Blattodea. A few other names associated with these *Musca* names have been included. When a name was originally proposed in another genus but subsequently placed in *Musca* and

there renamed because of homonymy, the original name is included despite its current placement. For example, Gmelin (1790: 2865) transferred *Rhagio maculatus* Fabricius to *Musca* and renamed it as *Musca adspersus* because the Fabrician name was a junior secondary homonym of *Musca maculata* Scopoli (1763: 326). *Rhagio maculatus* is included so that information on its type-locality and type-depository is available. Likewise, data for all valid senior synonyms of names proposed in *Musca* are included. A number of subsequent usages, usually representing new combinations, are also included because they were listed in Sherborn's *Index Animalium*. In Sherborn, there is usually no indication of the status of included names. We have thus included all names listed for *Musca* in Sherborn and have identified their status. These subsequent usages of *Musca* names can be confusing to modern workers. For example, Schumann (1986: 18) and Kurahashi (1989: 703) treated Scopoli's misidentification of *Musca carnaria* Linnaeus (a sarcophagid, but Scopoli applied the name to a calliphorid) as an available name. Likewise, Hennig (1963: 857) rejected the name *Musca laeta* Fallén because Sherborn (1922: 3355) listed *Musca laeta* Turton but did not indicate that the Turton name was only a subsequent use of *Syrphus laetus* Fabricius.

Altogether, the database includes 2,077 names, of which 1,709 are available, 904 are valid, 305 are junior homonyms, and 368 are unavailable. A number of new names (3), new combinations (64) and new synonyms (57) are given.

## Format

The general format of the entries in the alphabetical list is given below. Square brackets here denote information only included when relevant. Valid names are given in **BOLD** face type; available, but invalid names are given in *italics*; and unavailable names are in Roman.

**name** Author, publication year: page of indication and/or description  
(Original genus). Type locality or status of name. Kind of type Sex of type Depository of type. Family, current valid name including author to the extent known. [Comments and further information.]

Type locality data are given in full for all available names except replacement names and emendations. These data were taken from the original literature unless more detailed data were available from type specimens or from subsequent studies. Locality names have been corrected to conform to current usage and political boundaries where possible. When this is not possible, the original data are given in double quotes ("..."). For replacement names and emendations, the status of the name is given, such as "emend. [=emendation of] of *x-us* Author" or "n. n. [=new name for] *x-us* Author".

For *nomina nuda*, locality data are given, if known. For all other unavailable names, the status of the name is given, such as "incosp. [=incorrect original spelling of] *x-us* Author," or "missp. [=misspelling of] *x-us* Author." Data are also provided on the types of names or authentic material (of *nomina nuda*) where such are available. Two indexes are provided: One lists the names by biogeographical realm and the other by the current taxonomical position. The traditional division of the world into biotic regions is used and our definition of those regions conforms to the ones used by the various regional Diptera catalogs. The taxonomic index lists the names by family, genus, and species to which they currently belong. In both indexes, the index term is first, separated by a colon and followed by the list of the names that are associated with the term.

#### Cut-off date

All "*Musca*" names found in the literature before 1 January 1992 have been included in the database.

#### Lists of abbreviations used

For the status of a name, the following abbreviations are used:

N. n. = New name proposed for ...  
Incosp. = Incorrect original spelling of ...  
Missp. = [Subsequent] misspelling of ...  
Emend. = Emendation of ...

For the kind of type specimen(s), the following abbreviations are used:

A = adult,  
♂ ♀ = male(s) and female(s),  
♀ = female,  
L = larva,  
♂ = male,  
X = fossil

For the kind of type, the following abbreviations are used:

HT = holotype,  
LT = lectotype,  
NT = neotype,  
ST = syntypes, and  
T = type, status unknown or typical specimen(s).

*Holotype* is used when one specimen was designated as the type or holotype or equivalent term in the original description or when the original description

was clearly based on a unique specimen (see Article 73a). Under the *International Code of Zoological Nomenclature*, a holotype can only be designated in the original description. However, many modern authors have retrospectively designated "holotypes," usually because only one specimen remains extant. When the original description does not indicate the number of specimens upon which the name was based and this can not be determined from the description itself, then these subsequent "holotype" designations are lectotype designations (Article 74b). For all published declarations of "holotypes" for names which did not have valid holotype designations, we have included the citation to the "holotype" designation.

*Lectotype* is used when one of the syntypes has been subsequently designated as the type in accordance with article 74 of the *International Code of Zoological Nomenclature*. A citation is given to the source of the designation.

*Neotype* is used when a new specimen, not previously a type, has been designated in accordance with article 75 of *International Code of Zoological Nomenclature*. A citation is given to the source of the designation.

*Syntypes* is used when the original description clearly applies to multiple specimens or multiple specimens were listed as examined AND when no lectotype has been designated.

*Type* is used when the status of the type specimens is unknown. "T" is also used in the sense of typical with *nomina nuda* to indicate the presence of voucher specimens associated with the unavailable name.

For the depositories of type specimens, museum acronyms as used by Griffiths for the *Flies of the Nearctic Region* series (1980) are given. These and the other abbreviations used are as follows.

AMNH	American Museum of Natural History, New York
AMS	Australian Museum, Sydney
ANSP	Academy of Natural Sciences, Philadelphia
ASL	Academy of Sciences, St. Petersburg
Auction	Types sold at auction, current location unknown.
Author	Collection of the author of the name. See above under the section on workers for further information.
BMNH	The Natural History Museum, [formerly British Museum (Natural History)], London
BMNH B	Banks Collection, now deposited in BMNH
CAS	California Academy of Sciences, San Francisco
CNC	Canadian National Collection, Biosystematics Research Centre, Agriculture Canada, Ottawa

CRIK	Central Research Institute, Kasauli, India (Awati's material is probably lost)
DEI	Deutsches Entomologisches Institut, Eberswalde
DESTROY	Type known to be destroyed. The source of this information is either given in the comments for the entry or under the author in the section on workers.
HIPHA	High Institute of Public Health, Alexandria
IMZ	Istituto e Museo di Zoologia, University of Turin, Italy
IRSNB	Institut Royal de Sciences Naturelles de Belgique, Brussels
ISS	Instituto Superiore de Sanita, Rome
ITZA	Instituut voor Taxonomische Zoölogie, Amsterdam
LMNR	Latvian Museum of Nature, Riga
Lost	Type not found in appropriate collection, presumed to be lost
LSL	Linnaean Society, London
MCSNG	Museo Civico di Storia Naturale, Genoa
MCSNM	Museo Civico di Storia Naturale, Milan
MCZ	Museum of Comparative Zoology, Harvard University, Cambridge
MHNL	Musée d'Histoire Naturelle, Lille
MNHNP	Muséum National d'Histoire Naturelle, Paris
MRAC	Musée Royal de l'Afrique Centrale, Tervuren
MVM	Museum of Victoria, Abbotsford, Melbourne
NCIP	National Collection of Insects, Pretoria
NMI	National Museum of Ireland, Dublin
NMP	Natal Museum, Pietermaritzburg
NMW	Naturhistorisches Museum, Vienna
NRS	Naturhistoriska Riksmuseet, Stockholm
NSMT	National Science Museum, Tokyo
QMB	Queensland Museum, Brisbane
RNH	Nationaal Natuurhistorisch Museum, Leiden (formerly Rijksmuseum van Natuurlijke Historie)
SAIMR	South African Institute of Medical Research, Johannesburg
SAMCT	South African Museum, Cape Town
SIE	Shanghai Institute of Entomology, Shanghai
SMF	Senckenbergisches Museum für Naturkunde, Frankfurt-am-Main
SMN	Staatliches Museum für Naturkunde, Ludwigsburg
UMO	University Museum, Oxford [see Smith (1986) for details]
Unknown	Locality of type material unknown. More specific information may be found under the author in the section on workers.
USB	Universita degli Studi di Bologna, Bologna
USNM	National Museum of Natural History, Smithsonian Institution [formerly United States National Museum], Washington

UZMC	Universitetets Zoologiske Museum, Copenhagen
UZMC F	Fabrician Collection, now deposited at UZMC
Yeats	Yeats Collections, remains in University Museum, Glasgow
ZIL	Zoological Institute, University of Lund, Lund
ZIU	Zoological Institute, University of Uppsala, Uppsala
ZMHU	Institut für Spezielle Zoologie und Zoologisches Museum der Humboldt Universität, Berlin
ZSI	Zoological Survey of India, Calcutta
ZSZMH	Zoologisches Staatsinstitut und Zoologisches Museum, Hamburg

### Status of Names

The treatment of names follows the 3rd edition of the International Code of Zoological Nomenclature (hereafter, simply the CODE). Today, the status of a name is usually clearly indicated by the author with an explicit statement, such as "new species," "new name," etc. However, in the early literature, the status of names is frequently not so clearly indicated. The problem of "new" names in such comprehensive works as those of Gmelin (1790) and Turton (1801) is an example (see below) of the difficulties involved in determining the true status of names. When there is no explicit statement of status, three factors must be considered to determine the status: 1) Name - Is the spelling of the name identical or not to another name? 2) Characterization - Is the characterization given the same or different from that of other taxa? and 3) Attribution - Is the name attributed to another source? Names that are the same and are associated with the "same" characterization are considered to be the same, the later one being a subsequent usage of the earlier one. If a name is attributed to another worker and is identical in spelling to a name of the other worker, then it is a subsequent usage of that name and is not available unless it appears before the attributed usage. Even if the author deliberately mis-applies the name, the name is not available and is only a subsequent mis-identification unless the mis-application is being used as the type-species of a new genus (Art. 11 h(i)). If a name is attributed to another worker but differs in spelling from the name of the other worker, then the name may be an emendation, replacement, or a misspelling. Emendations, replacement names, and misspellings are separated only by the intent of the worker. Such intent is not always clearly documented. For emendations, the CODE defines intent as citation of both an original spelling and an emended spelling with the use of the emended spelling (Art. 33b(i)). The example given in the CODE of treatment of names in a similar way is not precisely true. For a new name, the CODE also implies a deliberate statement of replacement. Thus, incorrect subsequent spellings are defined by exclusion, that is, anything which is not an emendation nor a replacement name is a

subsequent incorrect spelling. Names that differ from prior ones in spelling and have no attribution are considered to be new proposals, especially if accompanied by a characterisation that is distinct. Our interpretation of names is summarized in the following table.

Some names in Gmelin (1790) and Turton (1801) present special problems. As noted above, names that differ from their original spelling may be considered as new names, emendations, or incorrect subsequent spellings. The first two possibilites are available names, but the last is not. A new name is a name expressly proposed as a replacement for an earlier name; an emendation is any demonstrably intentional change in the original spelling of a name; and incorrect subsequent spelling is anything else. The spellings used for some names in Gmelin and Turton are different from the original ones. The intent of Gmelin and Turton in making changes is not clear from their work. Gmelin gave references to the original spelling of all names, but did not give the original spelling nor an explanation for his deviation from it. Turton did not give the authority or the original source for names taken from Fabricius (1794) as he stated in his introduction that he followed “the last edition of the system of Fabricius” and that he omitted the numerous synonyms and references in Fabricius to reduce the bulk of his work. For names not in Fabricius, Turton did give a reference to the original source. In most cases, Turton did not give the original spelling and in no case did he give an explanation for his changes. The lack of an indication of intent would suggest that the changes in Gmelin and Turton should be considered simply as mistakes (incorrect subsequent spellings). And this is their proper status under the current CODE (ICZN, 1985, Art. 12(b)3 defines “indication” as “the proposal of a new replacement name (*nomen novum*) for an available name”). However, the requirement of an indication of intent for names published before 1930 was not part of earlier CODEs. In the prior edition, names published before 1930 need only be accompanied by an indication such as “the substitution of a new name for a previously established name” (ICZN 1964, Art. 16(a)(iii)). In Gmelin and Turton, it is clear that a new name (that is, one of a different spelling) has been substituted for an old name. Thus, while the purpose for the substitution (or lack of purpose, that is, an erroneous misspelling<sup>6</sup>), whether to replace or emend, was not implicitly stated, the result is indisputable. Most workers have accepted at least some of the substitutions as new replacement names. So, to reject all the substitu-

<sup>6</sup> We do recognize that some of the minor changes were probably unintentional mistakes. For example, a type setter may have easily mistaken *meditabunda* in Gmelin’s handwriting as *meditalnenda* which is what was set in type, and Gmelin may have overlooked the mistake in the proof stage. However, the problem of retrospectively deciding what was a mistake and what was an intentional change still remains!

tions in these works as merely incorrect subsequent spellings as the current CODE requires would be to upset current usage. We therefore consider all changed names to be available (as was the case under previous CODEs), but have arbitrarily considered minor changes, such as *linzensis* Gmelin for *lincensis* Schrank, to be emendations and major changes, such as *australis* Gmelin for *testacea* Fabricius, to be new names. The essential point is not whether the name is an emendation or a replacement name, but that the name (= change) is available and, although synonymous, must be considered for the purposes of homonymy.

**Comparison table for determination of the status of names which are not clearly designated as new.**

Class	Names	Characters	Attribution	Kind	Status
1	Same	Same	Absent	Subsequent Usage	Unavailable
2			Present	Subsequent Usage	Unavailable
3		Different	Present	Misidentification EXCEPT **	Unavailable
4			Absent	Homonym	Available
5	Different	Same	Absent	Synonym	Available
6			Present	Emendation, Misspelling & Replacement name ***	Available/ Unavailable
7		Different	Absent	Valid Name	Available
8			Present	Misidentification & Misspelling	Unavailable
9			Present, but ... *	Synonym or Valid Name	Available

NAME: Same name means identical spelling except that for family group names where it should be taken to mean same type genus and the variant spellings deemed identical in art. 58.

CHARACTER is a set of attributes that differ among taxa. Same characters means objectively having the same type; subjectively belonging to the same taxon.

ATTRIBUTION means some indication of prior use of the name. Minimally this is merely the citation of an authority and maximally this is a full citation to a publication. \*Case 9 is the inclusion of earlier names (concepts) in a synonymy for a new proposal, etc. Depending on the status of the earlier names included, the new proposal may be a synonym or a valid name.

KIND: Names are either valid, homonyms, junior synonyms, emendations or the subsequent use of previously published names. Those subsequent usages can be either misspellings misapplications (misidentifications) or correct spellings and applications. \*\*Misidentifications are unavailable except when the misidentification is deliberately used for the type species of a new genus. \*\*\*Emendations, misspellings, and replacement names are separated only by the intent of the worker. Such intent is not always clearly documented. For emendation, the Code defines intent as “when in the work itself, or in an author’s (or publisher’s) corrigenda, the original and the changed spelling are cited and the latter is adopted in the place of the former ...” (art. 33(b)i). For replacement names, the Code requires that the original and the replacement name be given. Hence, misspellings (incorrect subsequent spellings) are only defined by exclusion, that is, anything which is not an emendation or a replacement name is a subsequent incorrect spelling.

STATUS: Names are either available or unavailable for scientific nomenclature. The problems of class 6 are discussed in the text.

## Alphabetical listing of *Musca* Names

- abbreviata** Gravenhorst, 1807: 380 (*Musca*). Unknown, probably Germany. T A Lost. Scathophagidae.
- abdominalis** Turton, 1801: 650 (*Musca*). Subsequent Usage of *Syrphus abdominalis* Fabricius. Syrphidae, *Psarus coalescens* Geoffroy.
- abdominalis** Fabricius, 1805: 294 (*Musca*). East Indies. HT ♀ UZMC. Calliphoridae, *Thoracites abdominalis* Fabricius. See Townsend 1931c: 372.
- ablectus** Harris, 1776: 41, pl. 10, fig. 2 (*Musca*). England. T A Unknown. Syrphidae.
- abrupta** Gmelin, 1790: 2860 (*Musca*). Europe. T A NMI.
- accincta** Wiedemann, 1830: 407 (*Musca*). Unknown. T ♂ NMW. Calliphoridae, *Phormia regina* Meigen.
- acromion** Wiedemann, 1824: 47 (*Musca*). South America. T A UZMC. Tachinidae, *Cholomyia acromion* Wiedemann.
- acuminata** Gmelin, 1790: 2867 (*Musca*). Europe. T A NMI. Rhagionidae?
- adiaphora** Mayer, 1779: 184 (*Musca*). Czechoslovakia, Bohemia, Schuttenhofen Region. T A UNKNOWN.
- adspersus** Gmelin, 1790: 2865 (*Musca*). N. n. *Rhagio maculatus* Fabricius.
- adumbrata** Wiedemann, 1824: 48 (*Musca*). Java. T ♀ UZMC. Muscidae, *Grphomya adumbrata* Wiedemann.
- adunata** Geoffroy, 1785: 488 (*Musca*). France, Paris. T A Author. Syrphidae, *Ceriana* sp.
- aenea** Scopoli, 1763: 347 (*Musca*). Carniola. T A DESTROY. Syrphidae, *Cheilosia* sp.
- aenea** Gmelin, 1790: 2879 (*Musca*). Europe. T A NMI. Syrphidae. Preocc. Scopoli 1763.
- aenea** Fabricius, 1794: 335 (*Musca*). India. T A UZMC. Otitidae, *Physiphora clausa* Macquart. Preocc. Scopoli 1763. N. Comb.
- aenea** Zetterstedt, 1838: 656 (*Musca*). [Sweden, Lapland]. ST A NRS. Muscidae, *Pyrellia rapax* Harris. Preocc. Scopoli 1763.
- aequinoctialis** Linnaeus, 1758a: 598 (*Musca*). South America. T A NRS. Stratiomyidae.
- aerea** Fallén, 1825: 76 (*Musca*). Sweden, Skåne. LT ♂ NRS. Muscidae, *Spilogona aerea* Fallén. Lectotype designated by Pont 1984: 279.
- aestiuia** Müller, 1764: 86 (*Musca*). Denmark, Sjaelland, Frederiksdal. T A DESTROY.
- aestivationis** Linnaeus, 1761: 454 (*Musca*). Sweden, Uppsala. T A LSL.
- aestuans** Linnaeus, 1758a: 600 (*Musca*). Europe. T A LSL.
- aethiops** Gmelin, 1790: 2831 (*Musca*). Subsequent Usage of *Bibio aethiops* Fabricius. Bombyliidae, *Anthrax aethiops* Fabricius.
- aethiops** Stein, 1913: 467 (*Musca*). Tanzania, Moschi and Kilimandjaro. ST A ZMHU. Muscidae, *Musca aethiops* Stein.
- affinis** Turton, 1801: 639 (*Musca*). Subsequent Usage of *Syrphus affinis* Fabricius. Tachinidae, *Allophora hemiptera* Fabricius.
- affinis** Lamarck, 1816: 363 (*Musca*). Subsequent Usage of *Syrphus affinis* Fabricius. Tachinidae, *Allophora hemiptera* Fabricius.

- affinis** Fallén, 1817: 237 (*Musca*). Sweden, Skåne. LT ♂ NRS. Sarcophagidae, *Agria affinis* Fallén. Lectotype designated by Pape 1986: 303. N. Status  
Pape (1986: 303) used *punctata* Robineau-Desvoidy as he believed that *affinis* Fallén was preoccupied by “*Musca affinis*” Turton and “*Musca affinis*” La-marck, both names being merely subsequent usage of *Syrphus affinis* Fabricius.
- affinis** Meigen, 1838: 304 (*Musca*). Subsequent Usage of *Calliphora affinis* Mac-quart. Calliphoridae, *Calliphora vomitoria* Linnaeus.
- affixa** Walker, 1856a: 27 (*Musca*). Malaysia, Mt. Ophir. T ♂ BMNH. Muscidae, *Morellia affixa* Walker.
- afra** Paterson, 1956: 160, fig. 3 (*Musca*). Tanzania, Torina. HT ♂ SMN. Muscidae, *Musca afra* Paterson.
- africa** Wiedemann, 1824: 49 (*Musca*). South Africa, Cape of Good Hope. T ♀ UZMC. Sarcophagidae.
- africana** Beazzi, 1892: 190 (*Synamphoneura*). Somalia, Obbia. T ♀ MCSNM. Muscidae, *Musca lucidula* Loew.
- agilis** Harris, 1780: 86, pl. 25, fig. 19 (*Musca*). England. T A Unknown. Tachinidae.
- agilis** Rossi, 1790: 311 (*Musca*). Italy, Tuscany, Firenze & Pisa. T A ZMHU. Preocc. Harris 1780.
- agilis** Meigen, 1826: 70 (*Musca*). Germany. T A MNHNP. Calliphoridae, *Bellardia vulgaris* Robineau-Desvoidy. Preocc. Harris 1780. See Rognes 1991a: 44 about types. N. Comb.
- agilitas** Harris, 1780: 108, pl. 32, fig. 45 (*Musca*). England. T A Unknown. Syrphidae, *Platycheirus cyaneus* Müller.
- agromyzina** Fallén, 1825: 74 (*Musca*). Sweden, Skåne, near Äsperöd. LT ♀ NRS. Muscidae, *Coenosia agromyzina* Fallén. Lectotype designated by Pont 1984: 280.
- agrorum** Gmelin, 1790: 2869 (*Musca*). Subsequent Usage of *Syrphus agrorum* Fabricius. Syrphidae, *Palpada agrorum* Fabricius.
- alata** Gmelin, 1790: 2842 (*Musca*). Missp. *elata* Fabricius. Sciomyzidae, *Tetanocera elata* Fabricius.
- alba** Malloch, 1929a: 265 (*Byomya*). Sudan, Khartoum. HT ♂ BMNH. Muscidae, *Musca sorbens* Wiedemann.
- albiceps** Wiedemann, 1819a: 38 (*Musca*). South Africa, Cape of Good Hope. HT ♀ NMW. Calliphoridae, *Chrysomya albiceps* Wiedemann. See Townsend 1931c: 374.
- albicincta** Rossi, 1794: 69 (*Musca*). Italy. T A ZMHU. Scenopinidae, *Scenopinus albicinctus* Rossi.
- albicincta** Fallén, 1825: 73 (*Musca*). Sweden, Skåne. LT ♂ NRS. Anthomyiidae, *Calythea nigricans* Robineau-Desvoidy. Preocc. Rossi 1794. Lectotype designated by Michelsen, 1983: 110.
- albifrons** Linnaeus, 1761: 453 (*Musca*). Sweden. T A LSL. Tachinidae. 1 sp. in LSL (Dear MS).
- albimana** Gmelin, 1790: 2876 (*Musca*). Subsequent Usage of *Syrphus albimanus* Fabricius. Syrphidae, *Platycheirus cyaneus* Müller.
- albina** Wiedemann, 1830: 415 (*Musca*). East Indies. ST ♀ UZMC NMW. Muscidae, *Musca albina* Wiedemann.
- albipennis** Meigen, 1826: 58 (*Musca*). Germany, probably Stolberg. HT ♀ MNHNP. Muscidae, *Musca tempestiva* Fallén. See Pont 1986b: 211.

- albipes** de Villers, 1789a: 426 (*Musca*). Subsequent Usage of *Stratiomys albipes* Geoffroy. Stratiomyidae, *Nemotelus pantherinus* Linnaeus. Synonymy from de Villers.
- albiseta** Scopoli, 1763: 341 (*Musca*). Carniola. T A DESTROY. Sciomyzidae, *Ilione albiseta* Scopoli.
- abolineata** Fallén, 1823: 54 (*Musca*). Sweden, Skåne, Äsperöd. LT ♂ NRS. Muscidae, *Polites domitor* Harris. Lectotype designated by Pont 1984: 280.
- albomaculata** Macquart, 1843: 308 (*Musca*). Mauritius. HT ♀ MHNTP. Muscidae, *Musca xanthomelaena* Wiedemann. See Pont & Matile, 1976: 743.
- albomicans** Schummel, 1832: 74 (*Musca*). *Nomen nudum*.
- albula** Fallén, 1825: 74 (*Musca*). Sweden, Västergötland. LT ♂ NRS. Anthomyiidae, *Delia albula* Fallén. Lectotype designated by Michelsen 1983: 110.
- alceae** Preyssler, 1791: 129, pl. 4, fig. 20 (*Musca*). Czechoslovakia, Bohemia. T A DESTROY. Otitidae, *Physiphora alceae* Preyssler. See Rozkosny, Chvála & Pont 1982: 353.
- aldrichi** Townsend, 1931d: 468 (*Willistonia*). Brazil. HT ♂ NMW. Tachinidae, *Belvosia aldrichi* Townsend. As n. n. for "Musca esuriens" BB. Ald. (nec Fab., nec Wied.)."
- alevo** Harris, 1780: index (*Musca*). Incosp. *alevo* Harris. Fanniidae, *Fannia scalaris* Fabricius.
- alevo** Harris, 1780: 147, pl. 43, fig. 67 (*Musca*). England. T A Unknown. Fanniidae, *Fannia scalaris* Fabricius. Name improperly formed (verb).
- allii** Panzer, 1804: 72 (*Musca*). Germany, Regensburg. T A Unknown.
- alni** Scopoli, 1763: 332 (*Musca*). Carniola. T A DESTROY. Calliphoridae, *Eurychaeta muscaria?* Meigen.
- alpesa** Walker, 1849: 901 (*Musca*). Sierra Leone. HT ♂ BMNH. Muscidae, *Musca alpesa* Walker. See Pont 1990: 163.
- alterabilis** Gmelin, 1790: 2843 (*Musca*). Europe. T A NMI.
- alternata** Schrank, 1781: 448 (*Musca*). Austria. T A Unknown. Syrphidae, *Episyrphus cannabina* Scopoli.
- altratus** Harris, 1780: index (*Musca*). Missp. *atratus* Harris. Tachinidae.
- aluta** Walker, 1849: 911 (*Musca*). Lapland? LT ♀ BMNH. Muscidae, *Myospila meditabunda* Fabricius. Lectotype designated by Pont 1970b: 112.
- ambigua** Fallén, 1823: 56 (*Musca*). Sweden, Skåne, Äsperöd. LT ♂ ZIL. Anthomyiidae, *Zaphne ambigua* Fallén. Lectotype designated by Michelsen 1983: 110.
- ambulans** Müller, 1776: 177 (*Musca*). Denmark. T A DESTROY.
- amentaria** Scopoli, 1763: 335 (*Musca*). Carniola. NT ♂ BMNH. Calliphoridae, *Pollenia amentaria* Scopoli. Neotype designated by Rognes 1991a: 218.
- americana** Fabricius, 1775: 774 (*Musca*). South America. HT A UZMC. Cuterebridae, *Cuterebra americana* Fabricius. See Townsend 1931a: 80.
- americana** Swederus, 1787: 288 (*Musca*). North America. T A BMNH B. Syrphidae, *Eristalis barda* Say. Preocc. Fabricius 1775.
- amica** Linnaeus, 1771: 17 (*Musca*). Sweden, Ryby. T A UNKNOWN. Tephritidae. [Linnaeus 1785: 91].
- amica** Krastin, 1950: 549 (*Musca*). *Nomen nudum*. Muscidae, *Musca amita* Hennig.
- amica** Zimin, 1951: 145, fig. 126 (*Musca*). Mongolia & USSR, Siberia. ST ♂ ♀ ASL. Muscidae, *Musca amita* Hennig. Preocc. Linnaeus 1771.

- amicula** Harris, 1780: 158, pl. 47, fig. 7 (*Musca*). England. T A Unknown. Dolichopodidae.
- amita** Hennig, 1964: 986 (*Musca*). N. n. *Musca amica* Zimin. Muscidae, *Musca amita* Hennig.
- amoena** Harris, 1835: 599 (*Musca*). *Nomen nudum*. Massachusetts. T A MCZ. Muscidae, *Neomyia cornicina* Fabricius. Published without diagnosis or indication.
- analis** Macquart, 1843: 311, pl. 21, fig. 2 (*Musca*). Chile. HT ♂ MNHNP. Muscidae, *Musca domestica* Linnaeus. See Pont 1967: 182.
- anceps** Fabricius, 1805: 296 (*Musca*). South America. HT A UZMC. Tachinidae, *Trichodura anceps* Fabricius. See Townsend 1931a: 95.
- angelicae** Scopoli, 1763: 329 (*Musca*). Carniola. T A DESTROY. Muscidae, *Phaonia angelicae* Scopoli.
- angusticornis** Ratzeburg, 1844: 173 (*Musca*). Germany. T A Unknown. Tachinidae, *Baumhaueria goniaeformis* Meigen.
- angustifrons** Thomson, 1868: 546 (*Musca*). Marianas, Ponape. LT ♂ NRS. Muscidae, *Musca vetustissima* Walker. Lectotype designated by Pont 1977a: 128.
- anilis** Linnaeus, 1761: 442 (*Musca*). Sweden, Skåne. T A LSL. Therevidae, *Dialineura anilis* Linnaeus. 3 sp. in LSL (Dear MS).
- annexa** Müller, 1776: 173 (*Musca*). Denmark. T A Author. Syrphidae, *Neoascia annexa* Müller. See Thompson 1981b: 472.
- annonae** Fabricius, 1794: 358 (*Musca*). West Indies. T A UZMC F. Otitidae, *Euxesta annonae* Fabricius.
- annulata** Schrank, 1781: 477 (*Musca*). Austria, Vienna. T A Unknown. Preocc. Harris 1776.
- annulata** Geoffroy, 1785: 492 (*Musca*). France, Paris. T A Author. Preocc. Harris 1776.
- annulata** Fabricius, 1787: 348 (*Musca*). Cayenne. T A UZMC F. Micropezidae, *Taeniacptera meigenii* Perty. Preocc. Harris 1776.
- annulata** de Villers, 1789a: 418 (*Musca*). Subsequent Usage of *Nemotelus annulatus* De Geer. Rhagionidae, *Rhagio annulatus* De Geer.
- annulatus** Harris, 1776: 40, pl. 10, fig. 4 (*Musca*). England. T A Unknown. Syrphidae, *Volucella inanis* Linnaeus.
- anteiambulo** Harris, 1776: 60, pl. 15, fig. 17 (*Musca*). England. T A Unknown. Syrphidae, *Xanthogramma festiva* Linnaeus.
- anteiambulo** Harris, 1776: 60 (*Musca*). Incosp. **anteiambulo** Harris. Syrphidae, *Xanthogramma festiva* Linnaeus. Name revised by Goffe 1946: 75. Name improperly formed (verb).
- anthracina** Meigen, 1838: 301 (*Musca*). Germany & Belgium, Liège. LT ♀ MNHNP. Calliphoridae, *Melinda viridicyanea* Robineau-Desvoidy. Lectotype designated by Rognes 1991a: 203.
- anthrax** Schrank, 1781: 439 (*Musca*). Austria, Vienna, Prater. T A Unknown. Bombyliidae, *Anthrax anthrax* Schrank.
- antiquissima** Walker, 1849: 901 (*Musca*). Australia. HT ♂ BMNH. Muscidae, *Musca domestica* Linnaeus. See Pont 1973: 169 & 1990: 163.
- aphidioides** Scopoli, 1763: 349 (*Musca*). Carniola. T A DESTROY.
- aphidivora** Geoffroy, 1785: 481 (*Musca*). France, Paris. T A Author. Syrphidae.

- apiaria* Gmelin, 1790: 2867 (*Musca*). Subsequent Usage of *Syrphus apiarius* Fabricius. Muscidae, *Mesembrina mystacea* Linnaeus.
- apiformis* De Geer, 1776: 128, pl. 7, fig. 18 (*Musca*). Sweden. T A NRS. Syrphidae, *Microdon apiformis* De Geer.
- apiformis* Schrank, 1781: 459 (*Musca*). Austria, Vienna. T A Unknown. Syrphidae, *Pocota personata* Harris. Preocc. De Geer 1776.
- apiformis* Geoffroy, 1785: 488 (*Musca*). Sweden. T A LSL. Syrphidae, *Eristalis tenax* Linnaeus. Preocc. De Geer 1776.
- apivora* De Geer, 1776: 56, pl. 3, fig. 4 (*Musca*). Sweden. T A NRS. Syrphidae, *Volucella inanis* Linnaeus.
- aquatica* Geoffroy, 1785: 478 (*Musca*). France, Paris. T A Author. Otitidae, *Doryceria aquatica* Geoffroy. N. Comb.
- aratoria* Fabricius, 1794: 344 (*Musca*). Barbaria. T A MNHNP. Sciomyzidae, *Illione albisetosa* Scopoli.
- arbustorum* Linnaeus, 1758a: 591 (*Musca*). Sweden. T A LSL. Syrphidae, *Eristalis arbustorum* Linnaeus. See Thompson *et al.* 1982: 151.
- arctii* De Geer, 1776: 42, pl. 2, fig. 6 (*Musca*). N. n. *Musca solstitialis* Linnaeus. Tephritidae, *Urophora solstitialis* Linnaeus.
- arcuata* Linnaeus, 1758a: 592 (*Musca*). Sweden. LT ♀ LSL. Syrphidae, *Chrysotoxum arcuatum* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 152.
- arcuata* Fabricius, 1781: 451 (*Musca*). Germany, Kiel. T A UZMC F. Tephritidae, *Xyphosia miliaria* Schrank. Preocc. Linnaeus 1758.
- arcuata* Panzer, 1798c: 15 (*Musca*). Austria. T ♀ ZMHU. Syrphidae, *Brachyopa maculipennis* Thompson. Preocc. Linnaeus 1758.
- ardea* Turton, 1801: 633 (*Musca*). Subsequent Usage of *Rhagio ardea* Fabricius. Therevidae, *Cliorismia ardea* Fabricius.
- argentata* Gmelin, 1790: 2843 (*Musca*). Europe. T A NMI.
- argentea* Manuel, 1811: 19 (*Musca*). Emend. *argentata* Gmelin.
- argyrostoma* Schrank, 1781: 460 (*Musca*). Austria, Vienna. T A Unknown. Tachinidae, *Servilia ursina?* Meigen.
- aricioides* Walker, 1864: 216 (*Musca*). Aru Islands. HT ♂ BMNH. Muscidae, *Myiophaea spissa* Walker. See Pont 1966: 88.
- aristata* Panzer, 1801: 24 (*Musca*). Germany. T A ZMHU. Psilidae, *Loxocera ichneumonea* Linnaeus.
- armipes* Fallén, 1825: 75 (*Musca*). Sweden, Skåne. LT ♀ ZIL. Muscidae, *Hydrotaea armipes* Fallén. Lectotype designated by Pont 1984: 281.
- arnicae* Linnaeus, 1758a: 600 (*Musca*). Europe. LT A Lost. Tephritidae, *Tephritis arnicae* Linnaeus. Lectotype designated by White 1987: 101.
- arrogans* Linnaeus, 1761: 457 (*Musca*). Sweden. T A LSL. Empididae, *Tachydromia arrogans* Linnaeus. 1 sp. in LSL (Dear MS).
- artemisiae* Fabricius, 1794: 351 (*Musca*). Denmark. T A UZMC. Tephritidae, *Trypetia artemisiae* Fabricius.
- aruorum* Fabricius, 1787: 335 (*Syrphus*). China. T A UZMC. Syrphidae, *Eristalinus arvorum* Fabricius.
- arvorum* Gmelin, 1790: 2869 (*Musca*). Subsequent Usage of *Syrphus arvorum* Fabricius. Syrphidae, *Eristalinus arvorum* Fabricius.
- ascarides* Scudder, 1877: 756 (*Musca*). Colorado, White River [Oligocene]. T X USNM. Oestridae, *Hypoderma ascarides* Scudder.

- aselliformis** Geoffroy, 1785: 475 (*Musca*). France, Paris. T A Author.
- asiatica** Shinonaga & Kano, 1977: 111, fig. 2 (*Musca*). Thailand, Khao Yai. HT ♂ NSMT. Muscidae, *Musca asiatica* Shinonaga & Kano.
- asiliformis** Gmelin, 1790: 2871 (*Musca*). Subsequent Usage of *Syrphus asiliformis* Fabricius. Syrphidae, *Leucozona lucorum* Linnaeus.
- asiliformis** Preyssler, 1791: 99, pl. 3, fig. 13 (*Musca*). Czechoslovakia, Prague, Stromovke Park. T A DESTROY. Rhagionidae, *Chrysopilus asiliformis* Preyssler. Unrecognized See Rozkosny, Chvála & Pont 1982: 353.
- assimilis** Fallén, 1823: 56 (*Musca*). South Sweden. LT ♀ ZIL. Muscidae, *Muscina levida* Harris. Lectotype designated by Pont 1984: 281.
- ater** Harris, 1776: 58, pl. 15, fig. 11 (*Musca*). England. T A Unknown. Syrphidae, *Eristalinus sepulchralis* Linnaeus.
- aterrima** de Villers, 1789a: 548 (*Musca*). Europe. T A Author. Tachinidae, *Leucostoma aterrima* de Villers. See Townsend 1932: 33 (HT in MNHNP).
- aterrima** Fabricius, 1794: 334 (*Musca*). France. T A MNHNP. Phoridae, *Phora atra* Meigen. Preocc. de Villers 1789. N. Comb.
- atherix** Donovan, 1813: 23 (*Musca*). N. n. *Atherix maculatus* Meigen. Athericidae, *Atherix ibis* Fabricius.
- atra** Meuschen, 1787: 212 (*Musca*). Unknown. T A Auction. Preocc. Harris 1776.
- atra** de Villers, 1789a: 551 (*Musca*). Europe. T A Author. Rhinophoridae. Preocc. Harris 1776.
- atra** Fallén, 1823: 59 (*Musca*). Sweden, Skåne. HT ♀ ZIL. Muscidae, *Thricops longipes* Zetterstedt. Preocc. Harris 1776. See Pont 1984: 281.
- atramentaria** Meigen, 1826: 65 (*Musca*). Austria. T A NMW. Calliphoridae, *Pollenia atramentaria* Meigen.
- atrata** Fabricius, 1794: 333 (*Musca*). Germany. T A UZMC F. Piophilidae, *Piophila casei* Linnaeus. Preocc. Harris 1776.
- atratula** Ratzeburg, 1844: 170 (*Musca*). Germany. T A Unknown. Chamaemyiidae, *Leucopis atratula* Ratzeburg.
- atratus** Harris, 1776: 37, pl. 9, fig. 7 (*Musca*). England. T A Unknown. Tachinidae.
- atricolor** Fallén, 1825: 82 (*Musca*). Sweden, Skåne. LT ♂ NRS. Muscidae, *Helina atricolor* Fallén. Lectotype designated by Pont 1984: 281.
- atrifrons** Wiedemann, 1830: 403 (*Musca*). Unknown. T ♀ NMW. Tachinidae, *Tromodesiopsis atrifrons* Wiedemann. See Aldrich 1929: 4; Townsend 1931a: 96.
- atrifrons** Bigot, 1887: clxxiii (*Musca*). Cuba. LT ♀ BMNH. Muscidae, *Musca domestica* Linnaeus. Preocc. Wiedemann 1830. Lectotype designated by Pont 1990: 164.
- atropos** Schrank, 1776: 94 (*Musca*). Austria, Linz & Vienna. T A Unknown. Syrphidae, *Myathropa florea* Linnaeus.
- aucta** Walker, 1853b: 334 (*Musca*). Madras or Calcutta. T ♀ BMNH. Calliphoridae, *Calliphora vicina* Robineau-Desvoidy. Preocc. Harris 1780.
- auctus** Harris, 1780: 126, pl. 36, fig. 57 (*Musca*). England. T A Unknown. Anthomyiidae, *Pegomya* sp.
- audaculus** Harris, 1780: 121, pl. 35, fig. 41 (*Musca*). England. T A Unknown. Anthomyiidae, *Paregle audacula* Harris.
- augur** Fabricius, 1775: 777 (*Musca*). Australia. T A BMNH B. Calliphoridae, *Calliphora augur* Fabricius.

- aurata* Linnaeus, 1764a: 309 (*Musca*). *Nomen nudum*. [France]. Published in non-binomial work.
- aurata* Müller, 1764: 86 (*Musca*). Denmark, Sjaelland, Frederiksdal. T A Author.
- aurata* Fabricius, 1787: 347 (*Musca*). Germany, Kiel. T A UZMC F. Stratiomyidae, *Chloromyia formosa* Scopoli. Preocc. Müller 1764.
- aurifacies* Robineau-Desvoidy, 1830: 396 (*Musca*). France, Paris. T ♂ DESTROY. Muscidae, *Musca domestica* Linnaeus.
- auriflua* Wiedemann, 1830: 364 (*Musca*). [Brazil] Included under *Musca tessellata* Fabricius. HT A ZMHU. Sarcophagidae, *Oxysarcodexia aurifinis* Walker. Published in synonymy, not subsequently validated by usage. See Townsend (1931a: 74) for HT designation.
- aurifrons* de Villers, 1789a: 549, pl. 10, fig. 10 (*Musca*). Occitania [=France, Languedoc]. T A Author. Tachinidae, *Ectophasia crassipennis* Fabricius.
- aurulans* Robineau-Desvoidy, 1830: 397 (*Musca*). Brazil & Cayenne. T ♂ DESTROY. Muscidae, *Musca domestica* Linnaeus.
- aurulans* Walker, 1849: 893 (*Musca*). Subsequent Usage of *Calliphora aurulans* Robineau-Desvoidy. Calliphoridae, *Cynomyia cadaverina* Robineau-Desvoidy.
- australia* Sherborn, 1922: 607 (*Musca*). Missp. *australia* Macquart. Muscidae, *Musca domestica* Linnaeus.
- australis* Gmelin, 1790: 2833 (*Musca*). N. n. *Stratiomys testacea* Fabricius. Xylophagidae, *Coenomyia ferruginea* Scopoli.
- australis* Robineau-Desvoidy, 1830: 394 (*Sphora*). Australia, W. A., "Baie des Chiens-Marins." T ♀ DESTROY. Muscidae, *Musca domestica* Linnaeus.
- australis* Boisduval, 1835: 669 (*Musca*). Australia, Port King George. T A MNHNP. Calliphoridae, *Calliphora albifrontalis* Malloch. Preocc. Gmelin 1790.
- australis* Macquart, 1843: 309, pl. 20, fig. 10 (*Musca*). Tasmania, Hobart. LT ♂ MNHNP. Muscidae, *Musca domestica* Linnaeus. Preocc. Gmelin 1790. Lectotype designated by Pont 1967: 187.
- ausus* Harris, 1780: 85, pl. 25, fig. 16 (*Musca*). England. T A Unknown. Tachinidae.
- autumnalis* De Geer, 1776: 83 (*Musca*). Sweden. LT ♂ NRS. Muscidae, *Musca autumnalis* De Geer. Lectotype designated by Persson, Pont & Michelsen 1984: 92.
- avarus* Harris, 1780: 152, pl. 45, fig. 80 (*Musca*). England. T A Unknown. Heleomyzidae.
- avenae* Bjerckander, 1781: 172 (*Musca*). Sweden. T A UNKNOWN. Chloropidae, *Oscinella frit* Linnaeus.
- azurea* Fallén, 1817: 245 (*Musca*). Sweden, Västergötland. LT A NRS. Calliphoridae, *Protocalliphora azurea* Fallén. Lectotype designated by IZN 1990: 235.
- azurea* Walker, 1849: 880 (*Musca*). Subsequent Usage of *Lucilia azurea* Robineau-Desvoidy. Calliphoridae, *Lucilia caesar* Linnaeus.
- azurea* Doleschall, 1859: 111 (*Musca*). Amboina. T A Lost. Calliphoridae, *Lucilia papuensis* Macquart. Preocc. Fallén 1817. Types not found in NMW or ZMHU (Pont).
- bakeri* Patton, 1923a: 316 (*Musca*). Philippines. ST ♂ ♀ USNM. Muscidae, *Musca bakeri* Patton. See Pont 1990: 164; syntypes also in BMNH & MCSNM.
- balteata* De Geer, 1776: 116 (*Musca*). Sweden. T A NRS. Syrphidae, *Episyphus cannabina* Scopoli.

- bardus** Harris, 1780: 106, pl. 32, fig. 39 (*Musca*). England. T A Unknown. Syrphidae, *Cheilosia bardus* Harris.
- basalis** Walker, 1853b: 347 (*Musca*). Jamaica. T ♂ BMNH. Muscidae, *Morellia basalis* Walker.
- basilaris** Macquart, 1843: 310 (*Musca*). Brazil. T ♀ MNHNP. Muscidae, *Musca domestica* Linnaeus.
- bata** Walker, 1849: 875 (*Musca*). Unknown. T A BMNH. Calliphoridae, *Chrysomya megacephala* Fabricius.
- beckeri** Schnabl, 1911: 325 (*Plaxemyia*). Israel, Haifa. HT ♀ Lost? Muscidae, *Musca albina* Wiedemann. Type either in PAN (Schnabl collection, destroyed 1945) or ZMHU.
- benedicta** Walker, 1858b: 105 (*Musca*). Aru Islands. LT ♂ BMNH. Muscidae, *Neomyia timorensis* Robineau-Desvoidy. Lectotype designated by Pont 1966: 89.
- bezzii** Patton & Cragg, 1913a: 19, pl. 4 (*Musca*). India, Kodaikanal & Kotagiri. ST ♂ ♀ Lost. Muscidae, *Musca bezzii* Patton & Cragg. Types lost, not now in BMNH (Pont 1990: 164).
- bibosa** Scudder, 1877: 757 (*Musca*). Colorado, White River [Oligocene]. T X USNM.
- bibula** Wiedemann, 1830: 672 (*Musca*). *Nomen nudum*. A UNKNOWN. Calliphoridae, *Chrysomya albiceps* Wiedemann.
- bicincta** Linnaeus, 1758a: 592 (*Musca*). Sweden. LT ♂ LSL. Syrphidae, *Chrysotoxum bicinctum* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 152.
- bicinctata** Turton, 1801: 644 (*Musca*). N. n. *Musca bicincta* Linnaeus. Syrphidae, *Chrysotoxum bicinctum* Linnaeus.
- bicolor** de Villers, 1789a: 511 (*Musca*). N. n. *Musca putris* Geoffroy. T A Author.
- bicolor** Fabricius, 1798: 564 (*Musca*). France. T A MNHNP. Preocc. de Villers 1789.
- bicolor** Fabricius, 1805: 291 (*Musca*). South America. HT ♀ UZMC. Calliphoridae, *Mesembrinella bicolor* Fabricius. Preocc. de Villers 1789. See Townsend 1931a: 69.
- bicolor** Fallén, 1825: 91 (*Musca*). Subsequent Usage of *Anthomyia bicolor* Wiedemann. Anthomyiidae, *Pegomya bicolor* Wiedemann.
- bicolor** Meigen, 1838: 308 (*Musca*). Subsequent Usage of *Pollenia bicolor* Macquart. Calliphoridae, *Pollenia bicolor* Macquart.
- bicolor** Walker, 1853b: 339 (*Musca*). Unknown. T ♂ BMNH. Calliphoridae, *Phormia regina* Meigen. Preocc. de Villers 1789. N. Syn.
- bicolora** Sulzer, 1776: xxv (*Musca*). *Nomen nudum*. Switzerland. T A UNKNOWN. Bombyliidae, *Hemipenthes morio* Linnaeus. Published without diagnosis or indication. Synonymy by Fuesslin (1778: 165).
- bidentata** de Villers, 1789a: 429, pl. 9, fig. 14 (*Musca*). Southern France, Nemausum [= Nemours]. T A Author. Bibionidae, *Bibio* sp.
- bifasciata** Fabricius, 1775: 777 (*Musca*). West Indies, Virgin Islands. HT A UZMC F. Tachinidae, *Belvosia bifasciata* Fabricius. See Townsend 1931b: 176.
- bifasciata** Schrank, 1776: 95, pl. 3, fig. 21 (*Musca*). Austria, Linz. T A Unknown. Syrphidae, *Neoascia annexa* Müller. Preocc. Fabricius 1775. See Thompson 1981b: 472.
- bigemmis** Walker, 1853b: 341 (*Musca*). Cape. T ♂ BMNH. Calliphoridae, *Chrysomya chloropyga* Wiedemann.
- bilineata** de Villers, 1789a: 550 (*Musca*). Europe. T A Author.

- bilineata** Gmelin, 1790: 2830 (*Musca*). Subsequent Usage of *Bibio bilineata* Fabricius. Therevidae, *Anabarynchus bilineatus* Fabricius.
- bilineata** Gmelin, 1790: 2866 (*Musca*). N. n. *Rhagio punctatus* Fabricius. Anisopodidae, *Sylvicola punctatus* Fabricius. Preocc. de Villers 1789.
- bimaculata** Swederus, 1787: 290 (*Musca*). Australia. T A Lost. Type(s) not now in BMNH B (Pont).
- bipuncta** Wiedemann, 1830: 417 (*Musca*). Brazil. LT ♀ NMW. Muscidae, *Morellia bipuncta* Wiedemann. Lectotype designated by Pont in Pamplona (1986: 50).
- bipunctata** Scopoli, 1763: 341 (*Musca*). Carniola. T A DESTROY. Stratiomyidae, *Sargus bipunctatus* Scopoli.
- bipunctata** Müller, 1776: 174 (*Musca*). France, Paris. T A Author. Syrphidae, *Chrysotoxum arcuatum* Linnaeus. Preocc. Scopoli 1763.
- biseta** Hough, 1898: 173 (*Musca*). Somalia, Sheikh Husein. LT ♀ ANSP. Muscidae, *Musca biseta* Hough. Lectotype designated by Pont 1976: 269.
- bivittata** Doleschall, 1859: 111 (*Musca*). Amboina. T A DESTROY. Calliphoridae, *Euphumosia bivittata* Doleschall.
- bivittata** Thomson, 1868: 547 (*Musca*). Philippines, Luzon, Manila. T ♂ NRS. Muscidae, *Musca sorbens* Wiedemann. Preocc. Doleschall 1859.
- blandus** Harris, 1780: 106, pl. 32, fig. 38 (*Musca*). England. T A Unknown. Syrphidae, *Syrphus ribesii* Linnaeus.
- boganidae** Erichson, 1851: 67 (*Musca*). USSR, W. Taymyr, River Boganida. T A UNKNOWN. Calliphoridae, *Protophormia atriceps* Zetterstedt.
- bohemica** Preyssler, 1793: 203 (*Musca*). Czechoslovakia, Roupov in Plzen District. T A DESTROY. Tachinidae. See Rozkosny, Chvála & Pont 1982: 353.
- bombinatrix** Schrank, 1837: 17 (*Musca*). N. n. *Syrphus apriarius* Fabricius. Muscidae, *Mesembrina mystacea* Linnaeus.
- bombylans** Linnaeus, 1758a: 591 (*Musca*). Sweden. LT ♀ LSL. Syrphidae, *Volucella bombylans* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 152.
- bombyliformis** Geoffroy, 1785: 479 (*Musca*). France, Paris. T A Author. Syrphidae, *Merodon equestris* Fabricius.
- bombylina** Retzius, 1783: 181 (*Musca*). Emend. *bombylius* De Geer. Muscidae, *Mesembrina mystacea* Linnaeus.
- bombylius** De Geer, 1776: 133, pl. 8, fig. 1-3 (*Musca*). N. n. *Musca mystacea* Linnaeus. Muscidae, *Mesembrina mystacea* Linnaeus.
- bombyloides** Sulzer, 1776: 216 (*Musca*). Switzerland, Zürich. T ♂ Unknown. Muscidae, *Mesembrina mystacea* Linnaeus.
- borealis** Zetterstedt, 1837: 42 (*Musca*). *Nomen nudum*. [Sweden, Lapland]. T A ZIL. Muscidae, *Muscina levida* Harris.
- borealis** Zetterstedt, 1838: 660 (*Musca*). Sweden, Lapland. T A ZIL. Muscidae, *Muscina levida* Harris.
- bovina** Robineau-Desvoidy, 1830: 398 (*Musca*). France. T A DESTROY. Muscidae, *Musca autumnalis* De Geer.
- brachyptera** Panzer, 1798c: 16 (*Musca*). Germany, Nuremberg. T A ZMHU. Tachinidae, *Phasia crassipennis* Fabricius.
- braesia** Walker, 1849: 891 (*Musca*). South Africa. T ♂ BMNH. Muscidae, *Neomyia peronii* Robineau-Desvoidy.
- brassicaria** Fabricius, 1775: 778 (*Musca*). Europe. T A UZMC F. Tachinidae, *Cylindromyia brassicaria* Fabricius.

- brevis* Rossi, 1790: 306 (*Musca*). Italy, Tuscany, Firenze & Pisa. T A ZMHU.  
 Tachinidae, *Leucostoma aterrima* de Villers.
- brunnea* Thunberg, 1789: 89 (*Musca*). N. n. *Syrphus inanis* Fabricius. Syrphidae,  
*Copestylum brunneum* Thunberg.
- brunnea* Gmelin, 1790: 2859 (*Musca*). Europe. T A NMI. Preocc. Thunberg 1789.
- buccata* Scopoli, 1763: 342 (*Musca*). Carniola. T A DESTROY.
- buccata* Fallén, 1824: 65 (*Musca*). Sweden, Skåne. LT ♀ NRS. Anthomyiidae,  
*Leucophora sericea* Robineau-Desvoidy. Preocc. Scopoli 1763. Lectotype design-  
 nated by Michelsen 1983: 111.
- buphthalmi* Linnaeus, 1762: 471 (*Musca*). Palestine. T A LSL. Suppressed by  
 I.C.Z.N. 1914: 31.
- cadaverina* Linnaeus, 1758a: 595 (*Musca*). Sweden. LT ♀ LSL. Calliphoridae,  
*Lucilia caesar* Linnaeus. Lectotype designated by Pont 1981: 167.
- cadaverum* Kirby, 1837: 316 (*Musca*). "North America, lat. 65". T A Unknown.  
 Calliphoridae.
- caemeteriorum* Turton, 1801: 648 (*Musca*). Emend. *cemiteriorum* Linnaeus. Syrphi-  
 dae, *Chrysogaster cemiteriorum* Linnaeus.
- caerulea* Meigen, 1826: 63 (*Musca*). Germany. LT ♂ MNHNP. Calliphoridae,  
*Melinda gentilis* Robineau-Desvoidy. Preocc. Wiedemann 1819. Lectotype design-  
 nated by Rognes 1991a: 205.
- caerulea* Brullé, 1832: 315 (*Musca*). Greece, Morea. T ♀ UNKNOWN. Calliphori-  
 dae, *Lucilia* sp. Preocc. Wiedemann 1819.
- caeruleophthalmica* Scopoli, 1763: 331 (*Musca*). Carniola. T A DESTROY. Laux-  
 aniidae, *Sapromyza* sp.
- caerulescens* de Villers, 1789a: 464 (*Musca*). Bressia [= Bresse, France]. T A Author.  
 Stratiomyidae, *Sargus iridatus* Scopoli.
- caerulescens* Meigen, 1826: 55 (*Musca*). Austria. T ♀ NMW. Calliphoridae, *Lucilia*  
*sericata* Meigen. Preocc. de Villers 1789.
- caerulescens* Meigen, 1838: 307 (*Musca*). Subsequent Usage of *Pollenia caerules-  
 cens* Macquart. Calliphoridae, *Pollenia rufa* Fabricius.
- caesar* Linnaeus, 1758a: 595 (*Musca*). Europe. T A LSL. Calliphoridae, *Lucilia*  
*caesar* Linnaeus. 3 sp. in LSL (Dear MS).
- caesarina* Scopoli, 1763: 327 (*Musca*). Carniola. T A DESTROY. Calliphoridae,  
*Lucilia* sp.
- caesarion* Scharfenberg, 1805: 999 (*Musca*). Germany. T A Unknown. Calliphori-  
 dae, *Lucilia* sp. Accredited to Hellwig in the index (p. 1041).
- caesarion* Meigen, 1826: 57 (*Musca*). Portugal. HT ♀ NMW. Muscidae, *Neomyia*  
*cornicina* Fabricius. Preocc. Scharfenberg 1805. See Pont 1986b: 215.
- caesia* Rossi, 1790: 310 (*Musca*). Italy, Tuscany, Firenze & Pisa. T A ZMHU.  
 Stratiomyidae, *Microchrysa polita* Linnaeus.
- caesia* Meigen, 1826: 76 (*Musca*). Austria. HT ♀ NMW. Muscidae, *Muscina pro-  
 lapsa* Harris. Preocc. Rossi 1790. See Pont 1986b: 216.
- caesio* Harris, 1780: 75, pl. 21, fig. 8 (*Musca*). England. T A Unknown. Tephritidae,  
*Myoleja caesio* Harris.
- calamitosa* Bjerckander, 1777: 32 (*Musca*). Sweden. T A UNKNOWN. Chloropidae.
- calcarata* Turton, 1801: 648 (*Musca*). Subsequent Usage of *Syrphus calcaratus*  
 Fabricius. Syrphidae, *Merodon calcarata* Fabricius.

- calcitrans** Harris, 1780: 141 (*Musca*). Subsequent Usage of *Conops calcitrans* Linnaeus. Muscidae, *Stomoxys calcitrans* Linnaeus.
- calisia** Walker, 1849: 912 (*Musca*). South Africa. T ♀ DESTROY. Muscidae, *Morellia calisia* Walker. [teste Pont].
- calleva** Walker, 1849: 905 (*Musca*). South Africa. HT ♀ BMNH. Muscidae, *Musca calleva* Walker. See Pont 1990: 164.
- calliphoroides** Walker, 1861b: 245 (*Musca*). New Guinea, Dorey. ST ♀ BMNH. Calliphoridae, *Euphumosia papua* Guérin-Méneville. ST also in UMO (teste Pont).
- callosus** Harris, 1776: 61, pl. 15, fig. 18 (*Musca*). England. T A Unknown. Syrphidae, *Chrysotoxum bicinctum* Linnaeus.
- campestris** Robineau-Desvoidy, 1830: 395 (*Musca*). France, Saint-Sauveur? T ♀ DESTROY. Muscidae, *Musca autumnalis* De Geer.
- campicola** Robineau-Desvoidy, 1863: 627 (*Musca*). France, Nice. T A DESTROY. Muscidae, *Musca domestica* Linnaeus.
- canicularis** Linnaeus, 1761: 454 (*Musca*). N. n. *Musca lateralis* Linnaeus. Fanniidae, *Fannia canicularis* Linnaeus. Conserved by I.C.Z.N. 1969:124.
- canicularis** Houttuyn, 1768: 506 (*Musca*). Subsequent Usage of *Musca canicularis* Linnaeus. Fanniidae, *Fannia canicularis* Linnaeus.
- canina** Fabricius, 1781: 440 (*Musca*). England. T A BMNH B. Tachinidae, *Dexiosoma caninum* Fabricius.
- cannabina** Scopoli, 1763: 344 (*Musca*). Carniola. T A DESTROY. Syrphidae, *Episyphus cannabina* Scopoli.
- canus** Harris, 1780: 114, pl. 33, fig. 28 (*Musca*). England. T A Unknown. Tachinidae.
- capensis** Meuschen, 1787: 212 (*Musca*). *Nomen nudum*. Unknown. T A Auction. Bombyliidae? Published without diagnosis or indication.
- capensis** Zielke, 1971b: 214 (*Musca*). South Africa, S. W. Cape, N of Van Rhynsdorp. HT ♂ NMP. Muscidae, *Musca capensis* Zielke.
- capitata** De Geer, 1776: 23, pl. 1, fig. 300 (*Musca*). Sweden. T A NRS. Tachinidae, *Gonia capitata* De Geer.
- capito** Lichtenstein, 1796: 216 (*Musca*). Sinai. T A Auction.
- capraria** Linnaeus, 1758: 598 (*Musca*). Incosp. *cuprarius* Linnaeus. Stratiomyidae, *Sargus cuprarius* Linnaeus.
- capucina** Gmelin, 1790: 2831 (*Musca*). Subsequent Usage of *Bibio capucina* Fabricius. Bombyliidae, *Exoprosopa capucina* Fabricius.
- characteristica** Geoffroy, 1785: 494 (*Musca*). France, Paris. T A Author.
- carbonaria** Panzer, 1798a: 15 (*Musca*). Austria. T A ZMHU. Tachinidae, *Periscepsia carbonaria* Panzer. "HT in Berlin?, not seen"(Townsend 1932: 34).
- cardui** Linnaeus, 1758a: 600 (*Musca*). France. LT A Lost. Tephritidae, *Urophora cardui* Linnaeus. Lectotype designated by White 1987: 102.
- cardui** Geoffroy, 1785: 471 (*Musca*). Subsequent Usage of *Musca cardui* Linnaeus. Tephritidae, *Urophora cardui* Linnaeus.
- carinata** Meuschen, 1787: 212 (*Musca*). Unknown. T A Auction.
- carinifrons** Fallén, 1817: 243 (*Musca*). Sweden, Skåne. HT A NRS. Tachinidae, *Dinera carinifrons* Fallén. See Townsend 1932: 37.
- carnaria** Linnaeus, 1758a: 596 (*Musca*). Europe & America. LT ♂ LSL. Sarcophagidae, *Sarcophaga carnaria* Linnaeus. Lectotype designated by Richet 1986: 132.

- carnaria*, Scopoli, 1763: 325 (*Musca*). Misid. *carnaria* Linnaeus. Calliphoridae, *Calliphora vomitoria* Linnaeus.
- carnarida* Stephens, 1829a: 60 (*Musca*). *Nomen nudum*. England. T ♀ BMNH. Calliphoridae, *Protocalliphora rognesi* Thompson & Pont. Published without diagnosis or indication.
- carnifex* Robineau-Desvoidy, 1830: 393 (*Byomyia*). France, Saint-Sauveur. T ♂ DESTROY. Muscidae, *Musca tempestiva* Fallén.
- carnifex* Macquart, 1834: 158 (*Musca*). Subsequent Usage of *Byomyia carnifex* Robineau-Desvoidy. Muscidae, *Musca tempestiva* Fallén.
- carnivora* Fabricius, 1794: 313 (*Musca*). Germany. HT A UZMC F. Calliphoridae, *Calliphora vicina* Robineau-Desvoidy. Suppressed by I.C.Z.N. 1992: 90. See Rognes 1991a: 68.
- caruca* Walker, 1849: 877 (*Musca*). Chile. T ♀ BMNH. Calliphoridae, *Compsomyiops fulvicrura* Robineau-Desvoidy.
- casei* Linnaeus, 1758a: 597 (*Musca*). Europe. T A LSL. Piophilidae, *Piophila casei* Linnaeus.
- cassara* Pont, 1973: 160, fig. 47 (*Musca*). Solomon Islands, Guadalcanal, Kukum. HT ♂ BMNH. Muscidae, *Musca cassara* Pont. See Pont 1990: 164.
- castanea* Meuschen, 1787: 212 (*Musca*). Unknown. T A Auction.
- caudata* Geoffroy, 1785: 480 (*Musca*). France, Paris. T A Author.
- caudex* Harris, 1780: 158, pl. 47, fig. 6 (*Musca*). England. T A Unknown. Dolichopodidae.
- cautus* Harris, 1776: 60, pl. 15, fig. 15 (*Musca*). England. T ♀ Unknown. Syrphidae, *Chrysotoxum caustum* Harris.
- cellaris* Linnaeus, 1758a: 597 (*Musca*). Europe. T A LSL. Drosophilidae, *Drosophila funebris?* Fabricius. Type(s) lost (Bächli 1990: 845).
- cellaris* Bloch, 1776: 173 (*Musca*). Subsequent Usage of *Musca cellaris* Linnaeus. Drosophilidae, *Drosophila funebris?* Fabricius.
- cellaris* Geoffroy, 1785: 499 (*Musca*). Subsequent Usage of *Musca cellaris* Linnaeus. Drosophilidae, *Drosophila funebris?* Fabricius.
- celsus* Harris, 1780: 125, pl. 36, fig. 53 (*Musca*). England. T ♂ Unknown. Muscidae, *Helina celsa* Harris.
- cemiteriorum* Linnaeus, 1758a: 597 (*Musca*). Sweden. LT ♂ LSL. Syrphidae, *Chrysogaster cemiteriorum* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 152.
- centaureae* Fabricius, 1794: 360 (*Musca*). Germany, Kiel. T A UZMC F. Tephritidae, *Euleia heraclii* Linnaeus.
- cephalotes* Bosc, 1792: 55 (*Musca*). France, Paris. T A MNHNP. Pipunculidae, *Pipunculus campestris?* Meigen.
- cerasi* Linnaeus, 1758a: 600 (*Musca*). France. LT A Lost. Tephritidae, *Rhagoletis cerasi* Linnaeus. Lectotype designated by White 1987: 102.
- cerasi* Thunberg, 1789: 91 (*Musca*). T A UNKNOWN. Preocc. Linnaeus 1758.
- cerasina* Harris, 1833: 594 (*Musca*). *Nomen nudum*. Massachusetts. T A MCZ. Tachinidae. Published without diagnosis or indication. . A Say manuscript name.
- cerinus* Harris, 1776: 36, pl. 9, fig. 6 (*Musca*). England. T A Unknown. Tachinidae, *Gymnosoma rotundatum* Linnaeus.
- certima* Walker, 1849: 873 (*Musca*). Florida. HT ♀ BMNH. Calliphoridae, *Cochliomyia macellaria* Fabricius. [teste Pont].

- chaerophylli** Fabricius, 1798: 565 (*Musca*). Europe. T A UZMC F. Sciomyzidae, *Euthycera chaerophylli* Fabricius.
- chalybaea** Gravenhorst, 1807: 378 (*Musca*). Unknown, probably Germany. T A Lost. Calliphoridae.
- chalybata** Forster, 1770: 15 (*Musca*). *Nomen nudum*. England. Stratiomyidae, *Beris chalybata* Forster. Published without diagnosis or indication.
- chalybata** Forster, 1771: 95 (*Musca*). England. T A LSL. Stratiomyidae, *Beris chalybata* Forster.
- chalybea** Turton, 1801: 632 (*Musca*). N. n. *Musca chalybata* Forster. Stratiomyidae, *Beris chalybata* Forster.
- chalybea** Wiedemann, 1830: 402 (*Musca*). Java. T A RNH. Muscidae, *Neomyia gavisa* Walker. Preocc. Turton 1801.
- chalybeata** Gmelin, 1790: 2837 (*Musca*). Missp. *chalybata* Forster. Stratiomyidae, *Beris chalybata* Forster.
- chamaeleon** Linnaeus, 1758a: 589 (*Musca*). Europe. T A LSL. Stratiomyidae, *Stratiomys chamaeleon* Linnaeus. 4 sp. in LSL (Dear MS).
- chilensis** Walker, 1836: 354 (*Musca*). Chile. HT ♀ BMNH. Calliphoridae, *Neta chilensis* Walker. See Dear 1979:150.
- chilensis** Macquart, 1843: 311, pl. 20, fig. 6 (*Musca*). Chile. ST ♂ MNHNP. Muscidae, *Musca domestica* Linnaeus. Preocc. Walker 1836.
- chloris** Haliday, 1833: 165 (*Musca*). Ireland, Down, Holywood. T A NMI. Muscidae, *Neomyia cornicina* Fabricius.
- chloropyga** Wiedemann, 1818: 44 (*Musca*). South Africa, Cape of Good Hope. T A UZMC. Calliphoridae, *Chrysomya chloropyga* Wiedemann.
- chorea** Fabricius, 1781: 444 (*Musca*). Germany. T A UZMC F. Lonchaeidae, *Lonchaea chorea* Fabricius.
- chrysocephala** De Geer, 1776: 60 (*Musca*). N. n. *Musca mortuorum* Linnaeus. Calliphoridae, *Cynomya mortuorum* Linnaeus.
- chrysocephala** Gmelin, 1790: 2860 (*Musca*). Europe. T A NMI. Preocc. De Geer 1776.
- chrysoides** Walker, 1856a: 23 (*Musca*). Malaysia, Mt. Ophir. T ♂ BMNH. Calliphoridae, *Isomyia chrysoides* Walker.
- chrysoptera** Schrank, 1803: 126 (*Musca*). Germany, Bavaria. T A Unknown. Lauxaniidae, *Lauxania cylindricornis* Fabricius.
- chrysopus** Gmelin, 1790: 2865 (*Musca*). N. n. *Rhagio flavipes* Fabricius. Athericidae, *Atherix flavipes* Fabricius.
- chrysorrhæa** Scopoli, 1763: 347 (*Musca*). Upper Carniola. T A DESTROY. Syrphidae.
- chrysorrhœa** Meigen, 1826: 60 (*Musca*). [Austria] Neotype from Aachen, Germany. NT ♀ MNHNP. Calliphoridae, *Protocalliphora rognesi* Thompson & Pont. Preocc. Scopoli 1763. Neotype designated by Peus 1960: 209.
- chrysostoma** Lichtenstein, 1796: 216 (*Musca*). Osaureum [= Ouche, France]. T A Auction.
- chrysostoma** Stephens, 1829b: 303 (*Musca*). Britain. Tachinidae, *Dexiosoma caninum* Fabricius. Published in synonymy, not subsequently validated by usage. Published in synonymy with *Dexia canina* Fabricius.
- chrysotelus** Walker, 1853b: 346 (*Musca*). South Africa. T ♂ BMNH. Muscidae, *Pyrellina marsya* Walker.

- chrysazonias* Gmelin, 1790: 2879 (*Musca*). Europe. T A NMI. Syrphidae, *Chrysotoxum fasciatum* Müller.
- chui* Fan, 1965: 124 (*Musca*). China, Kwangtung, Nada. HT ♂ SIE. Muscidae, *Musca planiceps* Wiedemann.
- cibaria* Linnaeus, 1758a: 599 (*Musca*). Europe. T A LSL. Micropezidae, *Compsobata cibaria* Linnaeus. 1 sp. in LSL (Dear MS).
- cicur* Harris, 1776: 47, pl. 11, fig. 88 (*Musca*). England. T A Unknown. Stratiomyidae, *Chloromyia formosa* Scopoli.
- ciliata* Müller, 1776: 174 (*Musca*). Denmark. T A DESTROY. Dolichopodidae, *Poecilobothrus nobilitatus* Linnaeus.
- ciliata* de Villers, 1789a: 428, pl. 9, fig. 13 (*Musca*). France, Nemausum [= Nemours]. T A Author. Bibionidae, *Dilophus* sp. Preocc. Müller 1776.
- ciliata* Fabricius, 1794: 333 (*Musca*). Germany. HT ♂ UZMC F. Muscidae, *Hydrotaea diabolus* Harris. Preocc. Müller 1776. See Michelsen 1979a: 182.
- cimeoides* Fabricius, 1779: 253 (*Musca*). Norway. T A UZMC F. Empididae, *Trachydromia arrogans* Linnaeus.
- cimicoides* Fabricius, 1781: 447 (*Musca*). Missp. *cimeoides* Fabricius. Empididae, *Trachydromia arrogans* Linnaeus.
- cincta* Müller, 1766: 198 (*Musca*). Italy, Piedmont, Torino. T A IMZ. Tachinidae, *Mintho praeceps* Scopoli.
- cincta* Drury, 1773: 109 (*Musca*). Jamaica. T A Author. Syrphidae, *Meromacrus pinguis* Fabricius. Preocc. Müller 1766.
- cincta* de Villers, 1789a: 419 (*Musca*). Subsequent Usage of *Nemotelus cinctus* De Geer. Xylophagidae, *Xylophagus cinctus* De Geer.
- cincta* Fabricius, 1794: 347 (*Musca*). Denmark. T A UZMC. Sciomyzidae, *Coremacera fabricii* Rozkosny. Preocc. Müller 1766.
- cinctus* Harris, 1776: 43, pl. 10, fig. 6 (*Musca*). England. T ♂ Unknown. Syrphidae, *Eristalis lineata* Harris. Preocc. Müller 1766.
- cinctus* Fabricius, 1787: 333 (*Rhagio*). Germany, Kiel. T A UZMC F. Anisopodidae, *Sylvicola cinctus* Fabricius.
- cineraria* Rossi, 1794: 72 (*Musca*). Italy. T A ZMHU.
- cinerascens* Geoffroy, 1785: 493 (*Musca*). France, Paris. T A Author.
- cinerascens* Wiedemann, 1817: 79 (*Musca*). Germany, Holstein. T A UNKNOWN. Muscidae, *Muscina stabulans* Fallén. Preocc. Geoffroy 1785.
- cinerea* Linnaeus, 1764a: 301 (*Musca*). *Nomen nudum*. [Sweden]. Tephritidae, *Tephritis hyoscyami* Linnaeus. Published in non-binomial work.
- cinerea* de Villers, 1789a: 510 (*Musca*). Europe. T A Unknown. Preocc. Harris 1780.
- cinerea* Fabricius, 1794: 331 (*Musca*). India Orientali. HT ♂ UZMC. Sarcophagidae, *Leucomyia alba* Schiner. Preocc. Harris 1780. See Pape 1986: 302. N. Syn.
- cinerea* Fallén, 1824: 70 (*Musca*). Sweden, Skåne. LT ♂ NRS. Anthomyiidae, *Botanophilà silvatica* Robineau-Desvoidy. Preocc. Harris 1780. Lectotype designated by Michelsen 1983:111.
- cinerea* von Roser, 1840: 58 (*Musca*). Germany, Württemberg. T A SMN. Calliphoridae, *Pollenia* sp. Preocc. de Villers 1780.
- cinerella* Fallén, 1825: 77 (*Musca*). Sweden, Skåne. LT ♂ NRS. Anthomyiidae, *Adia cinerella* Fallén. Lectotype designated by Michelsen 1983: 111.
- cinereus* Harris, 1780: 75, pl. 21, fig. 11 (*Musca*). England. T A Unknown. Tephritidae, *Paroxyyna punctella* Fallén.

- cinerius* Harris, 1780: 74 (*Musca*). Incosp. *cinereus* Harris. Tephritidae, *Paroxyna punctella* Fallén. Name status by present revision.
- cingalaisina* Bigot, 1888: 606 (*Musca*). Sri Lanka. HT ♀ BMNH. Muscidae, *Musca planiceps* Wiedemann. See Pont 1990: 164.
- cingulata* Geoffroy, 1785: 477 (*Musca*). France, Paris. T A Author.
- cingulata* Fabricius, 1805: 301 (*Musca*). West Indies. HT ♂ UZMC. Tachinidae, *Calocarcelia fasciata* Townsend. Preocc. Geoffroy 1785. See Townsend 1931b: 174. N. Comb.
- cingulata* Donovan, 1808: 67, pl. 465 (*Musca*). England, Surrey, Nutfield. T ♂ UNKNOWN. Rhagionidae, *Chrysopilus cristatus* Fabricius. Preocc. Geoffroy 1785.
- cingulatus* Fabricius, 1775: 767 (*Syrphus*). New Zealand. T A BMNH B. Syrphidae, *Helophilus cingulatus* Fabricius.
- citrofasciata* Retzius, 1783: 179 (*Musca*). Missp. *citrofasciata* De Geer. Syrphidae, *Xanthogramma festiva* Linnaeus.
- citrofasciata* De Geer, 1776: 118 (*Musca*). N. n. *Musca festiva* Linnaeus. Syrphidae, *Xanthogramma festiva* Linnaeus.
- clandestina* Rossi, 1790: 314 (*Musca*). Italy, Tuscany, Firenze & Pisa. T A ZMHU.
- clauata* Fabricius, 1787: 351 (*Musca*). Germany, Kiel. T A UZMC F. Preocc. Drury 1773.
- clauda* de Villers, 1789a: 463 (*Musca*). Bressia [= Bresse, France]. T A Author. Syrphidae, *Merodon* sp.
- clavipes* Fabricius, 1781: 427 (*Syrphus*). Italy. T A UZMC F. Syrphidae, *Merodon clavipes* Fabricius.
- clavata* Drury, 1773: 103 (*Musca*). North America, New York. T A Author. Mydidae, *Mydas clavatus* Drury.
- clavicornis* Forster, 1770: 15 (*Musca*). *Nomen nudum*. England.
- clavicrus* Schrank, 1781: 471 (*Musca*). Austria, Vienna, Prater. T A Unknown. Chloropidae, *Meromyza saltatrix* Linnaeus.
- claviformis* Geoffroy, 1785: 476 (*Musca*). France, Paris. T A Author.
- clavipes* Linnaeus, 1767: 981 (*Musca*). Europe. T A LSL. Stratiomyidae, *Beris clavipes* Linnaeus. 2 sp. in LSL (Dear MS).
- claviventris* Gmelin, 1790: 2843 (*Musca*). Europe. T A NMI.
- clemens* Harris, 1780: 154, pl. 45, fig. 87 (*Musca*). England. T A Unknown. Fanniidae, *Fannia lustrator* Harris.
- cloacaris* Fabricius, Otto, 1780: 204 (*Musca*). Greenland, Umarsoroluk. T A UZMC. Scathophagidae.
- cluvia* Walker, 1849: 885 (*Musca*). West Indies. T ♀ BMNH. Calliphoridae, *Lucilia cluvia* Walker.
- coalescens* Geoffroy, 1785: 486 (*Musca*). France, Paris. T A Author. Syrphidae, *Psarus coalescens* Geoffroy.
- coarctata* Fallén, 1825: 84 (*Musca*). Sweden, Skåne. LT ♂ NRS. Anthomyiidae, *Delia coarctata* Fallén. Lectotype designated by Michelsen 1983: 111.
- coccinea* de Villers, 1789a: 426 (*Musca*). Subsequent Usage of *Stratiomyia coccinea* Geoffroy. Stratiomyidae, *Clitellaria ephippium* Fabricius.
- coeo* Harris, 1780: 143, pl. 42, fig. 51 (*Musca*). England. T ♂ Unknown. Muscidae, *Pyrellia rapax* Harris. Name improperly formed (verb).

- coerulata** Sherborn, 1922: 958 (*Musca*). Missp. *coerulea* Wiedemann. Muscidae, *Neomyia* sp.
- coerulea** De Geer, 1776: 57 (*Musca*, as *carnaria coerulea*). *Nomen nudum*. [Sweden]. Calliphoridae, *Calliphora vomitoria* Linnaeus. Not available; polynominal, not binominal.
- coerulea** Wiedemann, 1819a: 23 (*Musca*). Java. T ♀ DESTROY. Muscidae, *Neomyia* sp. See Pont 1973: 198.
- cognata** Meigen, 1830: 374 (*Musca*). N. n. *Musca caerulea* Meigen. Calliphoridae, *Melinda gentilis* Robineau-Desvoidy. N. Syn.
- coleoptrata** Scopoli, 1763: 338 (*Musca*). Carniola. T A DESTROY. Drosophilidae, *Stegana coleoptrata* Scopoli.
- collecta** Walker, 1859b: 139 (*Musca*). Sulawesi, Makassar. T ♂ BMNH. Calliphoridae, *Rhyncomya setipyga*? Villeneuve.
- colombaschensis** de Villers, 1789a: 555 (*Musca*). Subsequent Usage of *Oestrus columbacensis* Scopoli. Simuliidae, *Simulium* sp.
- colossus** Scopoli, 1763: 327 (*Musca*). Carniola. T A DESTROY. Sarcophagidae, *Sarcophaga carnaria?* Linnaeus.
- combinata** Linnaeus, 1767: 997 (*Musca*). Sweden. T A LSL. Opomyzidae, *Geomysa combinata* Linnaeus. 1 sp. in LSL (Dear MS).
- combrea** Walker, 1849: 876 (*Musca*). Unknown. T A BMNH. Calliphoridae, *Chrysomya megacephala* Fabricius.
- comio** Harris, 1780: 153, pl. 45, fig. 86 (*Musca*). Incosp. *comis* Harris. Anthomyiidae.
- comis** Harris, 1780: index (*Musca*). England. T A Unknown. Anthomyiidae.
- comito** Harris, 1780: 117, pl. 34, fig. 25 (*Musca*). England. T A Unknown. Scathophagidae, *Scathophaga lutaria* Fabricius. Name improperly formed (verb).
- commoror** Harris, 1780: 117, pl. 34, fig. 22 (*Musca*). England. T A Unknown. Acalyptrate.
- comoror** Harris, 1780: 74 (*Musca*). Incosp. *commoror* Harris. Acalyptrate. Name status by present revision.
- compressa** Fabricius, 1787: 346 (*Musca*). Spain. T A UZMC F. Tachinidae, *Mintho compressa* Fabricius.
- compressa** Fabricius, 1794: 337 (*Musca*). Cayenne. T A MNHNP. Tachinidae, *Cordyligaster petiolata* Turton. Preocc. Fabricius 1787. N. Comb.
- compressa** Walker, 1849: 893 (*Musca*). Subsequent Usage of *Calliphora compressa* Robineau-Desvoidy. Calliphoridae, *Cynomya cadaverina* Robineau-Desvoidy.
- compunctor** Sherborn, 1902: 235 (*Musca*). Missp. *compunctus* Harris. Muscidae, *Graphomya maculata* Scopoli.
- compunctus** Harris, 1780: 113, pl. 33, fig. 24 (*Musca*). England. T ♂ Unknown. Muscidae, *Graphomya maculata* Scopoli.
- comitus** Harris, 1780: 108, pl. 32, fig. 47 (*Musca*). England. T A Unknown. Syrphidae, *Xanthandrus comitus* Harris.
- concinna** Panzer, 1804: 195 (*Musca*). Germany, Regensburg. T A Unknown.
- concinna** Wiedemann, 1825: , pl. 21 (*Musca*). “In insula Mezzo Dalmatiae”. T A DEI. Calliphoridae, *Rhyncomya impavida* Rossi. Type(s) originally in collection of E. F. Germar.
- concolor** Walker, 1853b: 333 (*Musca*). South America. T ♂ DESTROY. Muscidae, *Helina concolor* Walker. [teste Pont].

- conducens** Walker, 1859b: 138 (*Musca*). Sulawesi, Makassar. HT ♂ BMNH. Muscidae, *Musca conducens* Walker. See Pont 1990: 164.
- conferta** Geoffroy, 1785: 474 (*Musca*). France, Paris. T A Author.
- confinis** Fallén, 1825: 80 (*Musca*, as *subpuncta* var. β). Sweden. LT ♂ NRS. Muscidae, *Helina confinis* Fallén. Lectotype designated by Pont 1984: 282.
- confiscata** Speiser, 1924: 104 (*Musca*). N. n. *Musca fasciata* Stein. Muscidae, *Musca confiscata* Speiser.
- confixa** Walker, 1856a: 26 (*Musca*). Mt. Ophir. T ♂ BMNH. Calliphoridae, *Isomyia confixa* Walker.
- conflagrans** Walker, 1861c: 261 (*Musca*). Sulawesi, Menado. T ♂ BMNH. Calliphoridae, *Isomyia delectans* Walker.
- confluo** Harris, 1780: 143, pl. 42, fig. 50 (*Musca*). England. T A Unknown. Tachinidae. Name improperly formed (verb).
- conformis** Fallén, 1825: 82 (*Musca*). Sweden, Skåne. LT ♀ NRS. Anthomyiidae, *Pegomya conformis* Fallén. Lectotype designated by Michelsen 1979b: 257.
- confusus** Harris, 1780: 110, pl. 33, fig. 51 (*Musca*). England. T A Unknown. Syrphidae, *Platycheirus granditarsus* Forster.
- congolensis** Villeneuve, 1916: 513 (*Musca*). Zaire. HT ♂ MRAC. Muscidae, *Musca alpese* Walker.
- conica** Gmelin, 1790: 2860 (*Musca*). Europe. T A NMI.
- conica** Panzer, 1798c: 20 (*Musca*). Austria. T ♂ ZMHU. Syrphidae, *Brachyopa panzeri* Goffe. Preocc. Gmelin 1790.
- conica** Fallén, 1825: 83 (*Musca*). Subsequent Usage of *Anthomyia conica* Wiedemann. Anthomyiidae, *Hydrophoria lancifer* Harris.
- connexa** Fabricius, 1794: 350 (*Musca*). France. T A MNHNP. Tephritidae, *Euphranta connexa* Fabricius.
- connexa** Walker, 1853b: 342 (*Musca*). Unknown. T ♀ BMNH. Muscidae, *Neomyia cornicina* Fabricius. Preocc. Fabricius 1794.
- conopsea** Gmelin, 1790: 2868 (*Musca*). Subsequent Usage of *Musca conopsooides* Linnaeus *sensu* Fabricius. Syrphidae, *Doros profuges* Harris.
- conopsooides** Linnaeus, 1758a: 590 (*Musca*). Europe. LT ♂ LSL. Syrphidae, *Ceriana conopsooides* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 153.
- conosoma** Jördens, 1802: 30, pl. 7, fig. 9 (*Ascaris*). Germany, Jena. TL UNKNOWN. Muscidae, *Musca domestica?* Linnaeus.
- consanguinea** Rondani, 1848: 78 (*Musca*). Brazil. T A USB. Muscidae, *Musca domestica* Linnaeus.
- consimilis** Fallén, 1825: 85 (*Musca*). Sweden, Bohuslän. LT ♂ ZIL. Muscidae, *Helina consimilis* Fallén. Lectotype designated by Pont 1984: 282.
- consisto** Harris, 1780: 112, pl. 33, fig. 56 (*Musca*). England. T A Unknown. Syrphidae, *Dasyphorus pinastri* De Geer. Name improperly formed (verb).
- consobrinus** Curtis, 1835: 549, pl. 549 (*Musca*). England, Putney. T ♂ MVM. Calliphoridae.
- conspersus** Harris, 1776: 38, pl. 9, fig. 11 (*Musca*). England. T ♀ Unknown. Tachinidae, *Trixia conspersa* Harris.
- consencis** Harris, 1780: 123, pl. 36, fig. 47 (*Musca*). England. T A Unknown. Sepsidae, *Themira putris* Linnaeus.
- constane** Harris, 1780: index (*Musca*). Incosp. *constans* Harris. Muscidae, *Hydrotaea meteorica* Linnaeus.

- constans* Harris, 1780: 149, pl. 43, fig. 75 (*Musca*). England. T ♀ Unknown. Muscidae, *Hydrotaea meteorica* Linnaeus.
- contigua* Fabricius, 1794: 347 (*Musca*). Brazil. T A UZMC.
- contigua* Walker, 1853b: 344 (*Musca*). United States. HT ♀ BMNH. Muscidae, *Musca domestica* Linnaeus. Preocc. Fabricius 1794. See Pont 1990: 164.
- continua* Panzer, 1798c: 19 (*Musca*). Austria. T A ZMHU. Tachinidae, *Clytiomyia continua* Panzer.
- continua* Robineau-Desvoidy, 1863: 628 (*Musca*). France, Nice. T A DESTROY. Muscidae, *Musca autumnalis* De Geer. Preocc. Panzer 1798.
- contracta* Geoffroy, 1785: 491 (*Musca*). Sweden. T A LSL. Syrphidae, *Xylota segnis* Linnaeus.
- convenio* Harris, 1780: 143, pl. 42, fig. 52 (*Musca*). England. T A Unknown. Tachinidae. Name improperly formed (verb).
- convexifrons* Thomson, 1868: 547 (*Musca*). Hong Kong. HT ♂ NRS. Muscidae, *Musca convexifrons* Thomson.
- convolo* Harris, 1780: 143, pl. 42, fig. 49 (*Musca*). England. T A Unknown. Tachinidae.
- corni* Scopoli, 1763: 328 (*Musca*). Carniola. T A DESTROY. Muscidae, *Mydaea corni* Scopoli.
- cornicena* Fabricius, 1794: 317 (*Musca*). Missp. *cornicina* Fabricius. Muscidae, *Neomyia cornicina* Fabricius.
- cornicina* Fabricius, 1781: 438 (*Musca*). Italy. HT A UZMC F. Muscidae, *Neomyia cornicina* Fabricius. See Michelsen 1979a: 183.
- cornuta* Scopoli, 1772: 123 (*Musca*). Idrija. T A Unknown. Tephritidae, *Vidalia cornuta* Scopoli.
- cornuta* Fabricius, 1794: 357 (*Musca*). Germany. T A UZMC F. Tephritidae, *Terellia ceratocera* Hendel. Preocc. Scopoli 1772.
- cornuta* Fabricius, 1805: 298 (*Musca*). Denmark, Sjaelland. HT ♂ UZMC. Muscidae, *Helina setiventris* Ringdahl. Preocc. Scopoli 1772. See Michelsen 1979a: 183.
- corollae* Turton, 1801: 650 (*Musca*). Subsequent Usage of *Syrphus corollae* Fabricius. Syrphidae, *Metasyrphus corollae* Fabricius.
- corrigiolata* Linnaeus, 1767: 995 (*Musca*). S. Europe. T A LSL. Micropezidae, *Micropeza corrigiolata* Linnaeus.
- coruina* Fabricius, 1781: 440 (*Musca*). N. n. *Musca autumnalis* De Geer. Muscidae, *Musca autumnalis* De Geer. See Michelsen 1979a: 184.
- corvina* Fabricius, 1794: 320 (*Musca*). Missp. *coruina* Fabricius. Muscidae, *Musca autumnalis* De Geer.
- corvinoides* Schnabl & Dziedzicki, 1911: 180 (*Musca*). *Nomen nudum*. Muscidae, *Musca larvipara* Portschinsky. Published in synonymy, not subsequently validated by usage.
- coryden* Harris, 1782: index (*Musca*). Missp. *corydon* Harris. Syrphidae, *Cheilosia corydon* Harris.
- corydon* Harris, 1780: 106, pl. 32, fig. 36 (*Musca*). England. T A Unknown. Syrphidae, *Cheilosia corydon* Harris.
- coryleti* Scopoli, 1763: 336 (*Musca*). Carniola. T A DESTROY. Sciomyzidae, *Pherbina coryleti* Scopoli.
- coryletti* Sherborn, 1902: 251 (*Musca*). Missp. *coryleti* Scopoli. Sciomyzidae, *Pherbina coryleti* Scopoli.

- coryli** Rossi, 1794: 72, pl. 3, fig. 1 (*Musca*). Italy. T A ZMHU. Tephritidae, *Aciura coryli* Rossi.
- cossonico** Harris, 1780: 74 (*Musca*). Incosp. *consencis* Harris. Sepsidae, *Themira putris* Linnaeus. Name status by present revision.
- costalis** Fabricius, 1794: 360 (*Musca*). West Indies. T A UZMC F. Otitidae, *Euxesta costalis* Fabricius.
- costalis** Walker, 1853a: 344 (*Musca*). South America. T ♀ BMNH. Tachinidae, *Myiopharus* sp. Preocc. Fabricius 1794.
- costalis** Walker, 1860: 159 (*Musca*). Moluccas, Amboina. HT ♂ BMNH. Calliphoridae, *Stilbomyella gloriosa* Walker. Preocc. Fabricius 1794. See Crosskey 1965: 95.
- costata** Linnaeus, 1767: 226 (*Musca*). Europe. T A UNKNOWN.
- costata** Panzer, 1800: 23 (*Musca*). Germany. T A ZMHU. Tachinidae, *Gymnosoma intermedium* Loew. Preocc. Linnaeus 1767.
- cothurnata** Panzer, 1798a: 20 (*Musca*). Austria. T ♂ ZMHU. Micropezidae, *Compsobata cibaria* Linnaeus.
- crabroniformis** de Villers, 1789a: 428 (*Musca*). France, Cevennes. T A Author. Syrphidae, *Milesia crabroniformis* de Villers.
- craggi** Patton, 1922: 75 (*Musca*). India, Kerala, Nilambur District. ST A BMNH. Muscidae, *Musca craggi* Patton. See Pont 1990: 164.
- crassa** Gmelin, 1790: 2867 (*Musca*). Subsequent Usage of *Syrphus crassus* Fabricius. Syrphidae, *Phytomia crassa* Fabricius.
- crassa** Wiedemann, 1830: 387 (*Musca*). Brazil. HT ♂ ZMHU. Tachinidae, *Chae-totheresia crassa* Wiedemann. See Townsend 1931a: 102.
- crassicornis** Fabricius, 1794: 328 (*Musca*). West Indies. T A UZMC F. Tachinidae, *Gonia crassicornis* Fabricius.
- crassipennis** Fabricius, 1794: 357 (*Musca*). Germany. T A UZMC F. Otitidae, *Melieria crassipennis* Fabricius.
- crassipennis** Lamarck, 1816: 363 (*Musca*). Subsequent Usage of *Syrphus crassipennis* Fabricius. Tachinidae, *Ectophasia crassipennis* Fabricius.
- crassipes** Schrank, 1781: 450 (*Musca*). Austria, Linz. T A Unknown. Syrphidae, *Merodon crassipes* Schrank.
- crassipes** Gmelin, 1790: 2860 (*Musca*). Europe. T A NMI. Preocc. Schrank 1781.
- crassirostris** Stein, 1903: 99 (*Musca*). Egypt, Luxor, Cairo & Assuan. ST ♂ ♀ ZMHU. Muscidae, *Musca crassirostris* Stein.
- crassisetata** Ratzeburg, 1844: 174 (*Musca*). Germany. T A Unknown. Tachinidae, *Masicera sphingivora* Robineau-Desvoidy.
- crax** Turton, 1801: 621 (*Musca*). N. n. *Musca crux* Fabricius. Tephritidae, *Euphranta crux* Fabricius.
- cibrata** de Villers, 1789a: 547, pl. 10, fig. 7 (*Musca*). Bressia [=Bresse, France]. T A Author. Sciomyzidae, *Coremacera marginata* Fabricius. N. Syn.
- cristata** Fabricius, 1775: 782 (*Musca*). England. T A UZMC F. Rhagionidae, *Chrysopilus cristatus* Fabricius.
- cristata** Fabricius, 1794: 339 (*Musca*). Barbaria. T A MNHNP. Anthomyiidae? Preocc. Fabricius 1775. Type figured by Coquebert (1804: 109).
- crocus** Harris, 1780: 118, pl. 34, fig. 27 (*Musca*). England. T A Unknown. Sciomyzidae, *Iline albisetata* Scopoli.
- cruciata** Geoffroy, 1785: 499 (*Musca*). France, Paris. T A Author.

- crucis** Gmelin, 1790: 2834 (*Musca*). Subsequent Usage of *Stratiomys crucis* Fabricius. Xylophagidae, *Coenomyia ferruginea* Scopoli.
- cruentata** Geoffroy, 1785: 487 (*Musca*). France, Paris. T A Author. Syrphidae, *Paragus cruentatus* Geoffroy.
- crux** Fabricius, 1794: 358 (*Musca*). India Orientali. T A UZMC. Tephritidae, *Euphranta crux* Fabricius.
- cryptarum** Turton, 1801: 640 (*Musca*). Subsequent Usage of *Syrphus cryptarum* Fabricius. Syrphidae, *Eristalis cryptarum* Fabricius.
- cucularia** Linnaeus, 1767: 995 (*Musca*). Sweden. T A LSL. Sciomyzidae, *Elgiva cucularia* Linnaeus. 1 sp. in LSL (Dear MS).
- cupraria** Linnaeus, 1758a: 598 (*Musca*). Europe. T A LSL. Stratiomyidae, *Sargus cuprarius* Linnaeus. 3 sp. in LSL (Dear MS).
- cupraria** Geoffroy, 1785: 491 (*Musca*). Subsequent Usage of *Musca cupraria* Linnaeus. Stratiomyidae, *Sargus cuprarius* Linnaeus.
- cuprea** Geoffroy, 1785: 489 (*Musca*). France, Paris. T A Author.
- cuprea** Macquart, 1835: 268 (*Musca*). France, Bordeaux. T ♀ Lost? Muscidae, *Musca tempestiva* Fallén. Preocc. Geoffroy 1785. Types not found in MHNH nor MHNL (Pont).
- cuprea** Walker, 1853b: 331 (*Musca*). South Australia. HT ♀ BMNH. Calliphoridae, *Metallea incisuralis* Macquart. Preocc. Geoffroy 1785. See Dear 1977: 821.
- cuprina** Wiedemann, 1830: 654 (*Musca*). China. T A UZMC. Calliphoridae, *Lucilia cuprina* Wiedemann.
- cursitans** Fabricius, 1775: 782 (*Musca*). England. T A UZMC F. Empididae, *Platypalpus cursitans* Fabricius.
- cursoria** Scopoli, 1763: 338 (*Musca*). Carniola. T A DESTROY. Empididae, *Tachypeza nubila?* Meigen.
- curviforceps** Saccà & Rivosecchi, 1956a: 497 (*Musca*, as *domestica* spp.). Tanzania, Arusha. ST ♂ ♀ ISS. Muscidae, *Musca curviforceps* Saccà & Rivosecchi.
- curvinervis** Zetterstedt, 1845: 1350 (*Musca*). Muscidae, *Musca tempestiva* Fallén. Published in synonymy, not subsequently validated by usage. A Bouché manuscript name.
- curvipes** Gmelin, 1790: 2871 (*Musca*). N. n. Syrphidae, *Syrphus clavipes* Fabricius, *Merodon clavipes* Fabricius.
- curvipes** Fallén, 1825: 90 (*Musca*). Sweden, Skåne. HT ♂ ZIL. Muscidae, *Hydrotaea meridionalis* Portschinsky. Preocc. Gmelin 1790. See Pont 1984: 282.
- cuthbertsoni** Patton, 1936: 470, fig. 2 (*Musca*). Zimbabwe, Salisbury. LT ♂ BMNH. Muscidae, *Musca calleva* Walker. Lectotype designated by Pont 1990: 164.
- cyanaea** Müller, 1764: 85 (*Musca*). Denmark, Sjaelland, Frederiksdal. T A DESTROY. Syrphidae, *Platycheirus cyaneus* Müller.
- cyanaea** Fabricius, 1781: 439 (*Musca*). South Africa, Cape of Good Hope. T A DESTROY. Muscidae, *Neomyia peronii* Robineau-Desvoidy. Preocc. Müller 1764. See Townsend 1931c: 369; Michelsen 1979a: 184.
- cyanella** Meigen, 1826: 77 (*Musca*). Germany, probably Stolberg. LT ♀ MHNH. Muscidae, *Eudasypnora cyanella* Meigen. Lectotype designated by Pont 1986b: 219.
- cyanophthalma** Gmelin, 1790: 2863 (*Musca*). N. n. *Musca annulata* Schrank.
- cylindracea** Gmelin, 1790: 2873 (*Musca*). N. n. Syrphidae, *Ocyptamus cylindricus* Fabricius.

- cylindrica* De Geer, 1776: 30, pl. 1, fig. 12 (*Musca*). Sweden. T A NRS. Tachinidae, *Cylindromyia brassicaria* Fabricius.
- cylindrica* Gmelin, 1790: 2860 (*Musca*). Europe. T A NMI. Preocc. De Geer 1776.
- cylindrica* Fabricius, 1794: 336 (*Musca*). Germany. T A UZMC F. Sepsidae, *Nemopoda nitidula* Fallén. Preocc. De Geer 1776.
- cylindrica* Fallén, 1817: 242 (*Musca*). Sweden, Skåne. T A NRS. Tachinidae, *Phylomyia volvulus* Fabricius. Preocc. De Geer 1776.
- cylindricornis* Fabricius, 1794: 332 (*Musca*). France. T A MNHNP. Lauxaniidae, *Lauxania cylindricornis* Fabricius.
- cylindricus* Fabricius, 1781: 429 (*Syrphus*). West Indies. T A UZMC F. Syrphidae, *Ocyptamus cylindricus* Fabricius.
- cynipsea* Linnaeus, 1758a: 599 (*Musca*). Europe. T A LSL. Sepsidae, *Sepsis cynipsea* Linnaeus. 2 sp. in LSL (Dear MS).
- cynophila* Panzer, 1794c: 22 (*Musca*). Germany, Mannheim. T A ZMHU. Piophilidae, *Thyreophora cynophila* Panzer.
- dasyops* Stein, 1913: 468 (*Musca*). Tanzania, Kilimanjaro. ST ♀ ZMHU. Muscidae, *Musca dasyops* Stein.
- dauci* Fabricius, 1787: 353 (*Musca*). Sweden. T A UZMC F. Tephritidae, *Urophora solstitialis* Linnaeus.
- dawsoni* Rainbow, 1897: 97 (*Degereria*). Ellice Is., Funafuti Atoll, Tuvalu. LT ♂ AMS. Muscidae, *Musca domestica* Linnaeus. Lectotype designated by Pont 1968: 89.
- deceptoria* Poda, 1761: 116 (*Musca*). Austria, Graz. T A Unknown. Tachinidae, *Clytiomyia pellucens?* Fallén.
- decolor* Fallén, 1824: 68 (*Musca*). Sweden, Skåne. LT ♀ NRS. Muscidae, *Drymeia vicana* Harris. Lectotype designated by Pont 1984: 282.
- decore* Harris, 1780: 146, pl. 43, fig. 63 (*Musca*). England. T A Unknown. Anthomyiidae. Name improperly formed (adverb).
- deduco* Harris, 1780: 125, pl. 36, fig. 54 (*Musca*). England. T ♀ Unknown. Muscidae, *Helina sexmaculata* Preyssler. Name improperly formed (verb).
- deduco* Harris, 1780: 120, pl. 35, fig. 34 (*Musca*). England. T A Unknown. Lauxaniidae. Name improperly formed (verb).
- defixa* Walker, 1856a: 24 (*Musca*). Singapore. T ♂ BMNH. Calliphoridae, *Chrysomya defixa* Walker.
- degener* Haliday, 1833: 165 (*Musca*). Ireland, Down, Holywood. T A NMI. Calliphoridae, *Pollenia degener* Haliday.
- dejectus* Harris, 1780: 120, pl. 35, fig. 38 (*Musca*). England. T A Unknown. Heleomyzidae.
- delectans* Walker, 1859b: 134 (*Musca*). Sulawesi, Makasar. ST ♀ BMNH. Calliphoridae, *Isomyia delectans* Walker. HT not found by Townsend 1931c: 376, but 2 STs are in BMNH.
- delicate* Harris, 1780: 118, pl. 34, fig. 26 (*Musca*). England. T A Unknown. Sciomyzidae.
- demandata* Fabricius, 1798: 564 (*Musca*). Germany, Kiel. T A UZMC F. Otitidae, *Physiphora alceae* Preyssler.
- demano* Harris, 1780: 74 (*Musca*). Incosp. *dimano* Harris. Tachinidae, *Eriothrix rufomaculata* De Geer. Name status by present revision.

- denigrata* Linnaeus, 1767: 981 (*Musca*). Europe. T A LSL. Bombyliidae, *Hemipenthes maurus* Linnaeus. 1 sp. in LSL (Dear MS).
- dentipes* Gmelin, 1790: 2877 (*Musca*). Subsequent Usage of *Syrphus dentipes* Fabricius. Syrphidae, *Senogaster dentipes* Fabricius.
- dentipes** Fabricius, 1805: 303 (*Musca*). Denmark. T A DESTROY. Muscidae, *Hydrotaea dentipes* Fabricius. See Michelsen 1979a: 185.
- depressa* Swederus, 1787: 287 (*Musca*). USSR, Kamchatka. T A BMNH B. Syrphidae, *Leucozona glauca* Linnaeus.
- depressa* de Villers, 1789a: 548 (*Musca*). Bressia [= Bresse, France]. T A Author. Preocc. Swederus 1787.
- depressa* Gmelin, 1790: 2861 (*Musca*). Europe. T A NMI. Preocc. Swederus 1787.
- depressa* Panzer, 1804: 191 (*Musca*). Germany, Regensburg. T A Unknown. Preocc. Swederus 1787.
- depressa* Meigen, 1826: 67 (*Musca*). Austria & Germany, Berlin & Hamburg. LT ♀ MNHNP. Calliphoridae, *Pollenia labialis* Robineau-Desvoidy. Preocc. Swederus 1787. Lectotype designated by Rognes 1991a: 228.
- depuncta** Fallén, 1825: 80 (*Musca*). Sweden. LT ♂ NRS. Muscidae, *Helina depuncta* Fallén. Lectotype designated by Pont 1984: 283.
- despecta* Gmelin, 1790: 2861 (*Musca*). Europe. T A NMI.
- determinata* Walker, 1853b: 345 (*Musca*). East Indies. HT ♀ BMNH. Muscidae, *Musca domestica* Linnaeus. See Pont 1990: 164.
- devia* Linnaeus, 1761: 446 (*Musca*). Sweden. T A Lost. Syrphidae, *Microdon devia* Linnaeus. See Thompson *et al.* 1982: 153.
- dexter* Harris, 1780: 108, pl. 32, fig. 46 (*Musca*). England. T A Unknown. Syrphidae, *Platycheirus dexter* Harris.
- diabolus* Harris, 1780: 126, pl. 36, fig. 58 (*Musca*). England. T ♂ Unknown. Muscidae, *Hydrotaea diabolus* Harris.
- diadema* Linnaeus, 1767: 982 (*Musca*). Europe. T A LSL. Dolichopodidae, *Medetera diadema* Linnaeus. 1 sp. in LSL (Dear MS).
- diaphana** Fabricius, 1775: 783 (*Musca*). Germany, Leipzig. T A UZMC F. Dolichopodidae, *Argyra diaphana* Fabricius.
- diaphana* Fallén, 1825: 92 (*Musca*). Subsequent Usage of *Anthomyia diaphana* Wiedemann. Muscidae, *Thricops diaphanus* Wiedemann.
- diaphana* Townsend, 1931b: 157 (*Musca*). Subsequent Usage of *Tachina diaphana* Fabricius. Tachinidae, *Archytas diaphana* Fabricius.
- dichotoma* Bezzi, 1911: 93 (*Musca*). Mozambique, Umbeluzi. HT ♀ USNM. Muscidae, *Musca sorbens* Wiedemann.
- diffidens* Walker, 1856a: 26 (*Musca*). Singapore. T ♂ BMNH. Muscidae, *Neomyia diffidens* Walker.
- diffusa* Walker, 1861d: 290 (*Musca*). Sulawesi, Batchian. HT ♂ BMNH. Calliphoridae, *Stilbomyella gloriosa* Walker. See Crosskey 1965: 95.
- dilecta* Wiedemann, 1830: 419 (*Musca*). Brazil. HT ♂ NMW. Tachinidae, *Medinophyo dilecta* Wiedemann. See Aldrich 1929: 19; Townsend 1931a: 93.
- dimano* Harris, 1780: 142, pl. 42, fig. 46 (*Musca*). England. T A Unknown. Tachinidae, *Eriothrix rufomaculata* De Geer. Name improperly formed (verb).
- dimidiata* Panzer, 1798c: 17 (*Musca*). Austria. T A ZMHU. Tachinidae, *Phasia crassipennis* Fabricius.

- dioclea** Walker, 1849: 869 (*Musca*). Borneo. T ♀ BMNH. Calliphoridae, *Phumosia dioclea* Walker. **N. Comb.**
- dioptalma** Linnaeus, 1758a: 593 (*Musca*). Sweden. LT ♂ LSL. Syrphidae, *Spi-lomyia dioptalma* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 154.
- dirae** Harris, 1780: 112, pl. 33, fig. 57 (*Musca*). England. T A Unknown. Syrphidae, *Pipiza lugubris* Fabricius.
- discoidea** Fabricius, 1787: 351 (*Musca*). Denmark, Copenhagen. T A UZMC. Tephritidae, *Platyparea discoidea* Fabricius.
- discolor** Linnaeus, 1758a: 592 (*Musca*). *Nomen nudum*. Muscidae, *Mesembrina mystacea* Linnaeus. Published in synonymy, not subsequently validated by usage. Mentioned under *M. intricaria*. See Pont 1981: 172.
- discolor** Fabricius, 1794: 320 (*Musca*). India Orientali. HT ♀ UZMC. Calliphoridae, *Stomorhina discolor* Fabricius. See Dear 1977: 792.
- dispar** Turton, 1801: 651 (*Musca*). Subsequent Usage of *Syrphus dispar* Fabricius. Syrphidae, *Toxomerus dispar* Fabricius.
- dispar** Fallén, 1823: 58 (*Musca*). Sweden. LT ♂ NRS. Muscidae, *Spilogona dispar* Fallén. Lectotype designated by Pont 1984: 283.
- distincta** Walker, 1853b: 346 (*Musca*). Senegal? T ♀ BMNH. Muscidae, *Pyrellina distincta* Walker.
- ditissima** Walker, 1861b: 244 (*Musca*). New Guinea, Dorey. T ♂ BMNH. Calliphoridae, *Isomyia ditissima* Walker.
- divaricata** Geoffroy, 1785: 495 (*Musca*). France, Paris. T A Author.
- divaricata** Awati, 1916b: 138 (*Musca*). India. ST ♀ CRIK. Muscidae, *Musca domes-tica* Linnaeus. Preocc. Geoffroy 1785.
- divisa** Walker, 1853b: 333 (*Musca*). East Indies. T ♀ BMNH. Calliphoridae, *Rhyn-comya divisa* Walker.
- divisa** Meigen, 1975: 419, pl. 119, fig. 11 (*Musca*, as *domestica* var.). *Nomen nudum*. [Germany, probably Stolberg]. Muscidae, *Musca domestica* Linnaeus. Varietal name proposed after 1960.
- dizonias** Gmelin, 1790: 2880 (*Musca*). N. n. *Musca bifasciata* Schrank. Syrphidae, *Neoascia annexa* Müller.
- docelea** Walker, 1849: 904 (*Musca*). Unknown. T ♀ DESTROY.
- dolens** Zetterstedt, 1837: 42 (*Musca*). *Nomen nudum*. [Sweden, Torne Lappmark, Torneträsk]. T ♂ ZIL. Calliphoridae, *Bellardia pubicornis* Zetterstedt.
- dolens** Zetterstedt, 1838: 654 (*Musca*). Sweden, Torne Lappmark, Torneträsk. HT ♂ ZIL. Calliphoridae, *Bellardia pubicornis* Zetterstedt. See Rognes 1991a: 51.
- doliata** Geoffroy, 1785: 487 (*Musca*). France, Paris. T A Author.
- domestica** Linnaeus, 1758a: 596 (*Musca*). Europe & America. T A DESTROY. Muscidae, *Musca domestica* Linnaeus. See Pont 1981: 168.
- domestica major** De Geer, 1776: 71, pl. 4, fig. 1 (*Musca*). *Nomen nudum*. N. n. *Musca domestica* Linnaeus. Muscidae, *Musca domestica* Linnaeus. Not available; poly-nominal, not binomial.
- domitor** Harris, 1780: 148, pl. 43, fig. 71 (*Musca*). England. T ♀ Unknown. Muscidae, *Polites domitor* Harris.
- domysa** Walker, 1849: 886 (*Musca*). Australia. HT ♀ BMNH. Muscidae, *Neomyia australis* Macquart. See Pont 1973: 203.

- doronici** Scopoli, 1763: 333 (*Musca*). Carniola. T A DESTROY. Calliphoridae, *Morinia doronici* Scopoli. N. Status
- dorsalis** Fabricius, 1775: 786 (*Musca*). Czechoslovakia, Bohemia. T A UZMC F. Sciomyzidae, *Hydromya dorsalis* Fabricius.
- dorsalis** Walker, 1849: 907 (*Musca*). Australia. T A BMNH. Calliphoridae, *Caliphora augur* Fabricius. Preocc. Fabricius 1775.
- dorsomaculata** Macquart, 1843: 309 (*Musca*). Mauritius. LT ♂ MNHNP. Muscidae, *Musca xanthomelaena* Wiedemann. Lectotype designated by Pont & Matile 1976: 744.
- dotata** Walker, 1856a: 25 (*Musca*). Singapore. T ♂ BMNH. Calliphoridae, *Isomyia dotata* Walker.
- dubia** Fallén, 1825: 84 (*Musca*). *Nomen nudum*. Sweden. T A NRS. Anthomyiidae, *Delia* sp. Published without diagnosis or indication. Mentioned in comparison with *Delia coarctata*.
- dubia** Walker, 1853b: 344 (*Musca*). South Australia. T ♂ BMNH. Calliphoridae, *Aphyssura dubia* Walker.
- dubitata** Fallén, 1825: 82 (*Musca*). Sweden, Skåne. HT ♀ ZIL. Anthomyiidae, *Hydrophoria lancifer* Harris. See Michelsen 1983: 112.
- duplicata** Geoffroy, 1785: 471 (*Musca*). France, Paris. T A Author.
- dux** Eschscholtz, 1822: 114 (*Musca*). Guam. T A Unknown. Calliphoridae, *Chrysomya megacephala* Fabricius.
- efflatouni** Salem & El-Sherif, 1960: 171, fig. 1 (*Musca*). Egypt, Gebel Elba. HT ♂ HIPHA. Muscidae, *Musca sorbens* Wiedemann.
- elara** Walker, 1849: 870 (*Musca*). South Africa. T ♀ BMNH. Calliphoridae, *Chrysomya albiceps* Wiedemann.
- elata** Fabricius, 1781: 441 (*Musca*). Germany, Kiel. T A UZMC F. Sciomyzidae, *Tetanocera elata* Fabricius.
- elater** Sherborn, 1902: 321 (*Musca*). missp. *elata* Fabricius. Sciomyzidae, *Tetanocera elata* Fabricius.
- elatior** Villeneuve, 1937: 4 (*Musca*). Zaire, Lulenga, Mission Rugari. HT ♀ MRAC. Muscidae, *Musca elatior* Villeneuve.
- electa** Walker, 1859b: 136 (*Musca*). Sulawesi, Makassar. LT ♂ BMNH. Muscidae, *Neomyia sperata* Walker. Lectotype designated by Pont 1966: 92.
- elegans** Harris, 1782: index (*Musca*). Missp. *eligans* Harris. Syrphidae, *Epistrophe eligans* Harris.
- elegans** de Villers, 1789a: 464, pl. 9, fig. 27 (*Musca*). Southern France. T A Author. Syrphidae, *Episyphus cannabina* Scopoli.
- elephantopus** Gmelin, 1790: 2880 (*Musca*). N. n. *Musca crassipes* Schrank. Syrphidae, *Merodon crassipes* Schrank.
- elevata** Fabricius, 1805: 303 (*Musca*). Nova Cambria [= Tasmania, southern tip]. T A Lost. Lauxaniidae, *Ceratolauxania elevata* Fabricius. See Evenhuis 1988: 193.
- eligans** Harris, 1780: 105 (*Musca*). England. T A Unknown. Syrphidae, *Epistrophe eligans* Harris.
- elongata** Müller, 1764: 85 (*Musca*). Denmark, Sjaelland, Frederiksdal. T A Author. Sepsidae, *Sepsis cynipsea* Linnaeus. See Thompson 1981b: 472.
- elongata** Fabricius, 1787: 348 (*Musca*). Sierra Leone. T A UZMC F. Stratiomyidae, *Pteicticus posticus* Wiedemann. Preocc. Müller 1764. N. Syn.
- elongata** de Villers, 1789a: 549 (*Musca*). Europe. T A Author. Preocc. Müller 1764.

- elongata* Fabricius, 1794: 338 (*Musca*). South America. T A UZMC. Stratiomyidae,  
*Ptecticus testaceus* Fabricius. Preocc. Müller 1764.
- elongata* Schrank, 1803: 108 (*Musca*). Subsequent Usage of *Syrphus elongatus* Fabricius. Syrphidae, *Baccha elongata* Fabricius.
- emoda* Walker, 1849: 872 (*Musca*). Egypt. T ♀ BMNH. Calliphoridae, *Chrysomya albiceps* Wiedemann.
- ephippium* Gmelin, 1790: 2834 (*Musca*). Subsequent Usage of *Stratiomys ephippium* Fabricius. Stratiomyidae, *Clitellaria ephippium* Fabricius.
- ephippium* Fabricius, 1794: 338 (*Musca*). France. T A BMNH. Micropezidae, *Neria ephippium* Fabricius.
- ephippium* Walckenaer, 1802: 397 (*Musca*). France, Paris. Micropezidae, *Neria ephippium* Fabricius. Probably the lack of attribution to Fabricius was an error.
- equestris* Fabricius, 1775: 782 (*Musca*). Brazil. T A BMNH B. Dolichopodidae.
- equestris* Meigen, 1826: 57 (*Musca*). Germany. HT ♂ NMW. Calliphoridae, *Lucilia illustris* Meigen. Preocc. Fabricius 1775. See Collin 1926: 259.
- ericetorum* Gmelin, 1790: 2870 (*Musca*). Subsequent Usage of *Syrphus ericetorum* Fabricius. Syrphidae, *Asarkina ericetorum* Fabricius.
- erinacea* Fallén, 1824: 65 (*Musca*). Sweden, Småland. LT ♂ NRS. Muscidae, *Phaonia angulicornis* Zetterstedt. Preocc. Fabricius 1794. Lectotype designated by Pont 1984: 283.
- erinaceus* Fabricius, 1794: 328 (*Musca*). Denmark, Copenhagen. T A UZMC. Tachinidae, *Peteina erinaceus* Fabricius.
- eristalooides* Walker, 1858b: 106 (*Musca*). Moluccas, Aru Islands. T A BMNH. Calliphoridae, *Euphumosia eristalooides* Walker. ST also in UMO (*teste* Pont).
- errans* Gmelin, 1790: 2872 (*Musca*). Subsequent Usage of *Syrphus errans* Fabricius. Syrphidae, *Phytomia errans* Fabricius.
- erratica* Linnaeus, 1758a: 593 (*Musca*). Sweden. LT ♀ LSL. Syrphidae, *Eriozona erratica* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 154.
- erratica* Scopoli, 1763: 345 (*Musca*). Carniola. T A DESTROY. Syrphidae, *Pipiza noctiluca* Linnaeus. Preocc. Linnaeus 1758.
- erratica* Panzer, 1804: 53 (*Musca*). Germany, Regensburg. T A Unknown. Preocc. Linnaeus 1758.
- erratica* Fallén, 1825: 77 (*Musca*). Sweden, Bäckaskog or Äsperöd. LT ♂ NRS. Muscidae, *Phaonia errans* Meigen. Preocc. Linnaeus 1758. Lectotype designated by Pont 1984: 284.
- erro* Harris, 1780: 123, pl. 36, fig. 48 (*Musca*). *Nomen nudum*. England. T A Unknown. Chloropidae. Name improperly formed (verb).
- erucarum* Schrank, 1803: 119 (*Musca*). N. n. *Musca larvarum major* De Geer. Tachinidae, *Tachina larvarum* Linnaeus.
- erythrocephala* De Geer, 1776: 146, pl. 29, fig. 2 (*Musca*). Surinam. LT ♀ NRS. Calliphoridae, *Cochliomyia macellaria* Fabricius. Lectotype designated by Thompson 1973: 356.
- erythrocephala* Fabricius, 1787: 351 (*Musca*). Sweden. T A UZMC F. Preocc. De Geer 1776.
- erythrocephala* de Villers, 1789b: cxxxvii (*Musca*). N. n. *Musca viridescens* de Villers [1789b: 548]. Tephritidae. Preocc. De Geer 1776.

- erythrocephala* Meigen, 1826: 62 (*Musca*). Germany. LT ♂ MNHNP. Calliphoridae, *Calliphora vicina* Robineau-Desvoidy. Preocc. De Geer 1776. Lectotype designated by Rognes 1991a: 68.
- erythrophthalma* Panzer, 1794a: 24 (*Musca*). Germany, Brunswick. T A ZMHU. Drosophilidae, *Drosophila funebris*? Fabricius. Type(s) not found (Bächli, *in litt.*). *esuriens*. Turton, 1801: 638 (*Musca*). Subsequent Usage of *Syrphus esuriens* Fabricius. Syrphidae, *Copestylum brunneum* Thunberg.
- esuriens* Fabricius, 1805: 301 (*Musca*). South America. HT ♂ UZMC. Tachinidae, *Iteuthelaira esuriens* Fabricius. See Townsend 1931b: 170.
- esuriens*, Townsend, 1931b: 176 (*Musca*). Misid. *esuriens* by Brauer & Bergenstamm & Aldrich. Tachinidae, *Belvosia aldrichi* Townsend.
- ethiopica* Zielke, 1973: 365 (*Musca*). Ethiopia, The Divide, NW of Addis Adaba. HT ♀ CNC. Muscidae, *Musca ethiopica* Zielke.
- eustolia* Walker, 1849: 909 (*Musca*). Sierra Leone. T ♂ BMNH. Muscidae, *Gracilimyia eustolia* Walker.
- europaenata* Bigot, 1888: 605 (*Musca*). Vietnam, Saigon. LT ♂ BMNH. Muscidae, *Musca sorbens* Wiedemann. Lectotype designated by Pont 1990: 165.
- evectus* Harris, 1780: 125, pl. 36, fig. 55 (*Musca*). England. T ♀ Unknown. Muscidae, *Helina erecta* Harris.
- evonymellae* Ratzeburg, 1844: 172 (*Musca*). Germany. T A UNKNOWN. Tachinidae, *Bactromyia aurulenta* Meigen.
- exactus* Harris, 1780: 147, pl. 43, fig. 70 (*Musca*). England. T A Unknown. Anthomyiidae.
- exalbida* Stein, 1913: 467 (*Musca*). "Cap". T ♂ NMW. Muscidae, *Musca sorbens* Wiedemann. Published in synonymy, not subsequently validated by usage. A male specimen in NMW.
- excors* Harris, 1780: 154, pl. 45, fig. 90 (*Musca*). England. T A Unknown. Anthomyiidae.
- exempta* Walker, 1856b: 128 (*Musca*). Borneo, Sarawak. T ♀ BMNH. Calliphoridae, *Chlororhinia exempta* Walker.
- exilis* Harris, 1780: 122, pl. 36, fig. 44 (*Musca*). England. T A Unknown. Anthomyiidae. Preocc. Harris 1780: 117.
- exilis* Harris, 1780: 117, pl. 34, fig. 24 (*Musca*). England. T A Unknown. Scathophagidae, *Scathophaga stercoraria* Linnaeus.
- eximia* Wiedemann, 1819b: 53 (*Musca*). Brazil. T A NMW. Calliphoridae, *Lucilia eximia* Wiedemann.
- exinanita* Gmelin, 1790: 2867 (*Musca*). N. n. *Syrphus inanis* Fabricius. Syrphidae, *Copestylum brunneum* Thunberg.
- extenuata* Rossi, 1790: 312 (*Musca*). Italy, Tuscany, Firenze & Pisa. T A ZMHU. Psilidae, *Chyliza extenuata* Rossi.
- extranea* Stephens, 1829b: 306 (*Musca*). England. Muscidae, *Phaonia tuguriorum* Scopoli. Published in synonymy, not subsequently validated by usage. Published in synonymy with *signata* Meigen.
- extrema* Walker, 1853b: 348 (*Musca*). Tasmania. HT ♀ BMNH. Muscidae, *Pyrellia tasmaniae* Macquart. See Pont 1973: 212.
- fabricii* Steyskal, 1968: 9 (*Hoplocheiloma*). N. n. *Musca fasciata* Fabricius. Micropezidae, *Hoplocheiloma totiana* Gmelin. N. Syn.

- facultas* Harris, 1780: 109, pl. 33, fig. 49 (*Musca*). England. T A Unknown. Syrphidae, *Melanostoma mellinum* Linnaeus.
- falcata* Scopoli, 1763: 330 (*Musca*). Carniola. T A DESTROY. Tephritidae, *Orellia falcata* Scopoli.
- fallax* Linnaeus, 1758a: 592 (*Musca*). Sweden. LT ♀ LSL. Syrphidae, *Blera fallax* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 154.
- fallax* Geoffroy, 1785: 496 (*Musca*). France, Paris. T A Author. Preocc. Linnaeus 1758.
- familiaris* Panzer, 1804: 208 (*Musca*). Germany, Regensburg. T A Unknown.
- familiaris* Harris, 1869: 336 (*Musca*). Massachusetts. ST A MCZ. Calliphoridae, *Pollenia rufidus* Fabricius. Preocc. Panzer 1804. See Rognes, 1987b: 498.
- fasciata* Müller, 1764: 85 (*Musca*). Denmark, Sjælland, Frederiksdal. T A Author. Syrphidae, *Chrysotoxum fasciatum* Müller.
- fasciata* Fabricius, 1775: 781 (*Musca*). America, West Indies. T A Auction. Micropezidae, *Hoplocheiloma tortiliana* Gmelin. Preocc. Müller 1764. Type(s) originally in collection of Otto Thott.
- fasciata* de Villers, 1789a: 419 (*Musca*). Subsequent Usage of *Nemotelus fasciata* De Geer. Therevidae, *Thereva plebeja* Linnaeus.
- fasciata* Fabricius, 1805: 299 (*Musca*). Guinea. HT ♂ UZMC. Tachinidae, *Bogosiella pomeroyi* Villeneuve. Preocc. Müller 1764. See Townsend 1931c: 388.
- fasciata* Walker, 1853b: 337 (*Musca*). Brazil. T ♀ BMNH. Calliphoridae, *Cochliomyia macellaria* Fabricius. Preocc. Müller 1764.
- fasciata* Stein, 1910: 149 (*Musca*). Seychelles, Mahé. LT ♂ BMNH. Muscidae, *Musca confiscata* Speiser. Preocc. Müller 1764. Lectotype designated by Pont 1990: 165.
- fasciatus* Fabricius, 1779: 233 (*Rhagio*). Norway. T A UZMC F. Rhagionidae.
- fasciolata* De Geer, 1776: 124, pl. 7, fig. 14 (*Musca*). Sweden. T A NRS. Syrphidae, *Chrysotoxum fasciolatum* De Geer.
- favillacea* Walker, 1859b: 135 (*Musca*). Sulawesi, Makassar. HT ♀ BMNH. Calliphoridae, *Bengalia ieuna* Fabricius. See Townsend 1931c: 371.
- felina* Fabricius, 1794: 317 (*Musca*). New Castilia [= Castilla la Nueva, Spain]. T A UZMC F. Calliphoridae, *Rhyncomya felina* Fabricius.
- felix* Walker, 1853b: 338 (*Musca*). Canary Islands. T ♂ BMNH. Calliphoridae, *Chrysomya albiceps* Wiedemann. N. Syn.
- femorata* Linnaeus, 1758a: 595 (*Musca*). Sweden. LT ♀ LSL. Syrphidae, *Chalcosyrphus femoratus* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 154.
- femorata* Panzer, 1794b: 24 (*Musca*). Germany, Dresden. T ♀ ZMHU. Syrphidae, *Lejops lineatus* Fabricius. Preocc. Linnaeus 1758.
- fenestralis* Linnaeus, 1758a: 597 (*Musca*). Europe. T A LSL. Scenopinidae, *Scenopinus fenestralis* Linnaeus. 1 sp. in LSL (Dear MS).
- fera* Linnaeus, 1761: 453 (*Musca*). Sweden. T A LSL. Tachinidae, *Tachina fera* Linnaeus. 5 sp. in LSL (Dear MS).
- fergusoni* Johnston & Bancroft, 1920a: 201 (*Musca*). Australia, Queensland, Eidsvold, Upper Burnett River. LT ♂ QMB. Muscidae, *Musca fergusoni* Johnston & Bancroft. Lectotype designated by Pont 1973: 140.
- ferina* Fallén, 1817: 242 (*Musca*). Sweden, Västergötland. T A NRS. Tachinidae, *Dinera ferina* Fallén.

- ferruginea** Scopoli, 1763: 340 (*Musca*). Carniola. T A DESTROY. Xylophagidae, *Coenomyia ferruginea* Scopoli.
- ferruginea** Fabricius, 1794: 342 (*Musca*). India Orientali. T A UZMC F. Tephritidae, *Bactrocera dorsalis* Hendel. Preocc. Scopoli 1763.
- fervida** Panzer, 1804: 168 (*Musca*). Germany, Regensburg. T A Unknown.
- festinans** Scopoli, 1763: 349 (*Musca*). Carniola. T A DESTROY. Phoridae, *Phora* sp.
- festiva** Linnaeus, 1758a: 593 (*Musca*). Sweden. LT ♂ LSL. Syrphidae, *Xanthogramma festiva* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 155.
- festiva** Geoffroy, 1785: 479 (*Musca*). France, Paris. T A Author. Syrphidae, *Chrysotoxum arcuatum* Linnaeus. Preocc. Linnaeus 1758.
- festiva** Donovan, 1811: 13, pl. 511 (*Musca*). England. T ♀ UNKNOWN. Dolichopodidae. Preocc. Linnaeus 1758.
- fictor** Harris, 1780: 152, pl. 45, fig. 81 (*Musca*). England. T A Unknown. Muscidae, *Helina fictor* Harris. Unrecognized.
- fieldi** Rainbow, 1897: 98, pl. 1, fig. 4 (*Ebenia*). Ellice Is, Funafuti Atoll, Tuvalu. HT ♀ AMS. Muscidae, *Musca vetustissima* Walker. See Pont 1968: 90.
- filata** Gmelin, 1790: 2830 (*Musca*). Subsequent Usage of *Bibio filata* Fabricius. Mydidae, *Mydas clavatus* Drury.
- filiformis** Fabricius, 1794: 337 (*Musca*). Germany. T A UZMC F. Micropezidae, *Micropeza corrigiolata* Linnaeus.
- fimetaria** Linnaeus, 1758a: 599 (*Musca*). Europe. T A LSL. Psilidae, *Psila fimetaria* Linnaeus. 2 sp. in LSL (Dear MS).
- fimetri** Linnaeus, 1761: 456 (*Musca*). Sweden. T A Lost. Sepsidae, *Themira putris* Linnaeus. Types lost, not in LSL (*teste* Pont MS).
- fimetorum** Geoffroy, 1785: 496 (*Musca*). France, Paris. T A Author.
- flauescens** Fabricius, 1798: 565 (*Musca*). Morocco, Tangiers. T A UZMC. Tephritidae, *Euleia marmorea* Fabricius. Preocc. Gmelin 1790.
- flauicans** Müller, 1764: 85 (*Musca*). Denmark, Sjaelland, Frederiksdal. T A Author.
- flava** Linnaeus, 1758a: 600 (*Musca*). Europe. T A LSL. Chyromyidae, *Chyromya flava* Linnaeus. 1 sp. in LSL (Dear MS).
- flava** Geoffroy, 1785: 499 (*Musca*). Subsequent Usage of *Musca flava* Linnaeus. Chyromyidae, *Chyromya flava* Linnaeus.
- flava** Schellenberg, 1803: 62 (*Musca*). Subsequent Usage of *Musca flava* Linnaeus. Psilidae, *Psila fimetaria* Linnaeus. Misidentification.
- flaveola** Fabricius, 1794: 343 (*Musca*). Denmark, Copenhagen. T A UZMC. Dryomyzidae, *Dryomyza flaveola* Fabricius.
- flavescens** de Villers, 1789a: 433 (*Musca*). Subsequent Usage of *Volucella flavescens* Geoffroy. Syrphidae.
- flavescens** Gmelin, 1790: 2860 (*Musca*). Europe. T A NMI.
- flavicans** Panzer, 1798a: 18 (*Musca*). Austria. T ♀ ZMHU. Preocc. Müller 1764.
- flavicans** Doleschall, 1859: 112 (*Musca*). Amboina. T A NMW. Calliphoridae, *Pollenia* sp. Preocc. Müller 1764. Type(s) not found in Vienna [*teste* Pont].
- flavicans** Sherborn, 1902: 373 (*Musca*). Missp. *flauicans* Müller.
- flaviceps** Ratzeburg, 1844: 172 (*Musca*). Germany. T A Unknown. Tachinidae, *Ceromyia flaviceps* Ratzeburg.
- flaviceps** Walker, 1849: 870 (*Musca*). Subsequent Usage of *Lucilia flaviceps* Macquart. Calliphoridae, *Chrysomya megacephala* Fabricius.

- flavicornis* Gmelin, 1790: 2874 (*Musca*). Subsequent Usage of *Syrphus flavicornis* Fabricius. Syrphidae, *Cheilosia flavicornis* Fabricius.
- flavicornis* Fallén, 1825: 88 (*Musca*). Sweden, Skåne. HT ♂ ZIL. Muscidae, *Coenosia flavicornis* Fallén. See Pont 1984: 284.
- flavifacies* Bigot, 1888: 606 (*Musca*). New Caledonia. HT ♀ BMNH. Muscidae, *Musca domestica* Linnaeus. See Pont 1990: 165.
- flavifrons* Panzer, 1798c: 22 (*Musca*). Austria. T A ZMHU. Sciomyzidae, *Limnia unguicornis* Scopoli.
- flavinervis* Thomson, 1868: 547 (*Musca*). Ross Island [=Cocos-Keeling]. T ♀ NRS. Muscidae, *Musca domestica* Linnaeus.
- flavipalpis* Meigen, 1838: 305 (*Musca*). N. n. *Musca autissiodorensis* Robineau-Desvoidy. Calliphoridae, *Calliphora autissiodorensis* Robineau-Desvoidy.
- flavipennis* Fallén, 1823: 59 (*Musca*). Sweden, Västergötland. LT ♀ ZIL. Anthomyiidae, *Chirosia flavipennis* Fallén. Lectotype designated by Michelsen 1983: 112.
- flavipennis* Bigot, 1887: clxxiii (*Musca*). USA, Rocky Mts. LT ♂ BMNH. Muscidae, *Musca domestica* Linnaeus. Preocc. Fallén 1823. Lectotype designated by Pont 1990: 165.
- flavipes* Fabricius, 1781: 420 (*Rhagio*). Germany. T A UZMC F. Athericidae, *Atherix flavipes* Fabricius.
- flavipes* de Villers, 1789a: 419 (*Musca*). Subsequent Usage of *Nemotelus flavipes* De Geer.
- flavipes* Fabricius, 1794: 254 (*Bibio*). Germany. T A UZMC F. Therevidae, *Dialineura anilis* Linnaeus.
- flavipes* Schrank, 1803: 123 (*Musca*). Germany, Bavaria. T A Unknown. Sciomyzidae, *Sepedon sphegeus* Fabricius.
- flavipes* Fallén, 1825: 90 (*Musca*). Sweden, Skåne. LT ♂ NRS. Anthomyiidae, *Pegomya pulchripes* Loew. Preocc. Schrank 1803. Lectotype designated by Michelsen 1983: 112.
- flavissima* Turton, 1801: 629 (*Musca*). Subsequent Usage of *Stratiomys flavissima* Rossi. Stratiomyidae, *Odontomyia flavissima* Rossi.
- fletcheri* Patton & Senior-White, 1924: 574, pl. 31, fig. 21 (*Musca*). India, Samalkota. LT ♂ BMNH. Muscidae, *Musca fletcheri* Patton & Senior-White. Lectotype designated by Pont 1990: 165.
- floralis* Fallén, 1824: 71 (*Musca*). Sweden, Skåne. LT ♂ ZIL. Anthomyiidae, *Delia floralis* Fallén. Lectotype designated by Michelsen 1983: 112.
- floralis* Robineau-Desvoidy, 1830: 397 (*Musca*). France. T ♀ DESTROY. Muscidae, *Musca autumnalis* De Geer. Preocc. Fallén 1824.
- floralis* Meigen, 1838: 307 (*Musca*). Subsequent Usage of *Pollenia floralis* Robineau-Desvoidy. Calliphoridae, *Pollenia rudis* Fabricius.
- florea* Linnaeus, 1758a: 591 (*Musca*). Sweden. LT ♂ LSL. Syrphidae, *Myathropa florea* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 156.
- florea* Fabricius, 1794: 255 (*Bibio*). Germany. T A UZMC F. Phoridae, *Diplonevra florescens* Turton. Preocc. Linnaeus 1758. N. Syn.
- florescens* Turton, 1801: 636 (*Musca*). N. n. *Bibio florea* Fabricius. Phoridae, *Diplonevra florescens* Turton. N. Status
- florescentiae* Linnaeus, 1758a: 601 (*Musca*). Europe. T A LSL. Tephritidae, *Trypetia florescentiae* Linnaeus.

- florum** Fabricius, 1794: 352 (*Musca*). Italy. T A UZMC F. Opomyzidae, *Opomyza florum* Fabricius.
- formosa** Scopoli, 1763: 339 (*Musca*). Carniola. T A DESTROY. Stratiomyidae, *Chloromyia formosa* Scopoli.
- formosa** Panzer, 1798b: 21 (*Musca*). Austria. T A ZMHU. Otitidae, *Otites elegans* Latreille. Preocc. Scopoli 1763. N. Syn.
- formosa** Donovan, 1830: 320 (*Musca*). [Australia]. Tachinidae, *Rutilia formosa* Robineau-Desvoidy. Published in synonymy, not subsequently validated by usage.
- formosana** Malloch, 1925: 375 (*Musca*). Taiwan, Okaseki. HT ♂ DEI. Muscidae, *Musca formosana* Malloch.
- formosus** Harris, 1780: 107, pl. 32, fig. 43 (*Musca*). England. T A Unknown. Syrphidae, *Epistrophe grossulariae* Meigen. Preocc. Scopoli 1763.
- fortunata** Walker, 1859: 137 (*Musca*). Sulawesi, Makassar. T ♂ BMNH. Calliphoridae, *Hemipyrellia ligurriens* Wiedemann.
- freedmani** Paterson, 1957: 108, fig. 3 (*Musca*). Botswana, near Sekhuma. HT ♂ SAIMR. Muscidae, *Musca freedmani* Paterson.
- frigida** Fabricius, 1805: 307 (*Musca*). Norway, Lapland. T A UZMC. Coelopidae, *Coelopa frigida* Fabricius.
- frit** Linnaeus, 1758a: 598 (*Musca*). Europe. T A LSL. Chloropidae, *Oscinella frit* Linnaeus. 3 sp. in LSL (Dear MS).
- frondescentiae** Linnaeus, 1758a: 601 (*Musca*). Europe. T A LSL. Otitidae, *Herina frondescentiae* Linnaeus. 2 sp. in LSL (Dear MS).
- frontalis** Latreille, 1802: 457 (*Musca*). Unknown (=Europe). T A MNHNP. Sarcophagidae, *Metopia argyrocephala* Meigen.
- frontalis** Gimmerthal, 1834: 115 (*Musca*). Livonia (=Estonia & Latvia). T ♂ LMNR. Muscidae. Preocc. Latreille 1802.
- frontalis** Macquart, 1843: 308, pl. 21, fig. 3 (*Musca*). Algeria, Algiers. T ♂ MHNL. Muscidae, *Musca domestica* Linnaeus. Preocc. Latreille 1802.
- frontalis** Rondani, 1868: 51 (*Musca*). Italy. T A USB. Muscidae, *Musca domestica* Linnaeus. Preocc. Latreille 1802.
- frutetorum** Gmelin, 1790: 2870 (*Musca*). Subsequent Usage of *Syrphus frutetorum* Fabricius. Syrphidae, *Parhelophilus frutetorum* Fabricius.
- fucata** Fabricius, 1794: 359 (*Musca*). West Indies. T A UZMC F. Tephritidae, *Acinia tessariae* Kieffer & Jorgensen. Preocc. Harris 1780.
- fucatus** Harris, 1780: 83, pl. 24, fig. 29 (*Musca*). England. T A Unknown. Syrphidae, *Xylota segnis* Linnaeus.
- fuciformis** Fabricius, 1794: 290 (*Syrphus*). France. T A MNHNP. Syrphidae, *Mallota fuciformis* Fabricius.
- fucina** Walker, 1849: 883 (*Musca*). South Africa. T ♀ BMNH. Calliphoridae, *Lucilia cuprina* Wiedemann.
- fulges** Harris, 1780: 87, pl. 25, fig. 23 (*Musca*). England. T A Unknown. Calliphoridae, *Lucilia caesar* Linnaeus.
- fulva** Gmelin, 1790: 2878 (*Musca*). Europe. T A NMI. Syrphidae, *Cheilosia illustrata* Harris. Preocc. Harris 1780.
- fulvescens** Robineau-Desvoidy, 1830: 397 (*Musca*). Cayenne. T ♂ DESTROY. Muscidae, *Musca domestica* Linnaeus.
- fulvicornis** Meigen, 1838: 307 (*Musca*). Subsequent Usage of *Pollenia fulvicornis* Robineau-Desvoidy. Calliphoridae, *Pollenia rudis* Fabricius.

- fulvicornis* Zetterstedt, 1859: 6186 (*Musca*). Sweden, Skåne. HT ♂ ZIL. Calliphoridae, *Pollenia viatica* Robineau-Desvoidy. Preocc. Robineau-Desvoidy 1830. See Rognes 1991aa: 230, 1991ab: 483.
- fulvimaculata* Manuel, 1811: 20 (*Musca*). N. n. *Musca quadrimaculata* Swederus. Calliphoridae, *Calliphora quadrimaculata* Swederus.
- fulvipalpis* Meigen, 1838: 308 (*Musca*). Subsequent Usage of *Pollenia fulvipalpis* Macquart. Calliphoridae, *Pollenia rufis* Fabricius.
- fulvipes* Geoffroy, 1785: 495 (*Musca*). France, Paris. T A Author.
- fulvithorax* Santos Abreu, 1976: 140 (*Musca*, as *vitripennis* var.). *Nomen nudum*. Canary Is. Muscidae, *Musca osiris* Wiedemann. Varietal name proposed after 1960.
- fulviventris* Schrank, 1781: 469 (*Musca*). Austria, Vienna. T A Unknown. Platystomatidae, *Platystoma seminationis* Fabricius.
- fulvus* Harris, 1780: 80, pl. 24, fig. 22 (*Musca*). England, Essex, East Tilbury. T A Unknown. Syrphidae, *Arctophila superbens* Müller.
- fumigata* Scopoli, 1763: 336 (*Musca*). Carniola. T A DESTROY. Sciomyzidae, *Euthycera fumigata* Scopoli.
- fumipennis* Walker, 1856a: 25 (*Musca*). Singapore. T ♂ BMNH. Calliphoridae, *Hypopygiopsis fumipennis* Walker.
- funebres* Harris, 1780: 106, pl. 32, fig. 37 (*Musca*). England. Syrphidae, *Cheilosia funebres* Harris.
- funebris* Harris, 1782: index (*Musca*). Missp. *funebres* Harris. Syrphidae, *Cheilosia funebres* Harris.
- funebris* Fabricius, 1787: 345 (*Musca*). Denmark, Copenhagen. ST ♀ UZMC. Drosophilidae, *Drosophila funebris* Fabricius. See Bächli 1982: 293.
- funesta* Turton, 1801: 648 (*Musca*). Subsequent Usage of *Syrphus funestus* Fabricius. Syrphidae, *Merodon funesta* Fabricius.
- fungorum* Scopoli, 1763: 337 (*Musca*). Carniola. T A DESTROY.
- fungorum* De Geer, 1776: 89, pl. 5, fig. 2 (*Musca*). Sweden. LT ♂ NRS. Anthomyiidae, *Pegomya steini* Hendel. Preocc. Scopoli 1763. Lectotype designated by Persson, Pont & Michelsen 1984: 92.
- funibres* Harris, 1780: 74 (*Musca*). Incosp. *funebres* Harris. Syrphidae, *Cheilosia funebres* Harris. Name status by present revision.
- furcata* Fabricius, 1794: 343 (*Musca*). France. T A MNHNP. Piophilidae, *Centrophlebomyia furcata* Fabricius.
- furcata* Turton, 1801: 622 (*Musca*). N. n. *Musca fucata* Fabricius. Tephritidae, *Acinia tessariae* Kieffer & Jorgensen. Preocc. Fabricius 1794.
- furta* Linnaeus, 1767: 991 (*Musca*). Sweden, Uppsala. LT ♂ ZIU. Drosophilidae, *Stegana furta* Linnaeus. Lectotype designated by Bächli 1982: 299.
- fusca* de Villers, 1789a: 511 (*Musca*). Sweden. T A Unknown. Preocc. Harris 1776.
- fusca* Gmelin, 1790: 2833 (*Musca*). Non-European. T A NMI. Bombyliidae. Preocc. Harris 1776.
- fuscata* Gmelin, 1790: 2865 (*Musca*). Subsequent Usage of *Tipula fuscata* Fabricius. Anisopodidae, *Sylvicola fuscatus* Fabricius.
- fuscata* Fallén, 1825: 85 (*Musca*). Sweden, Skåne. LT ♂ NRS. Muscidae, *Phaonia fuscata* Fallén. Lectotype designated by Pont 1984: 284.
- fuscipennis* Fabricius, 1805: 291 (*Musca*). Sumatra. HT ♀ UZMC. Calliphoridae, *Silbomyia fuscipennis* Fabricius. See Townsend 1931c: 374; Crosskey 1965:68.

- fuscipennis* von Roser, 1840: 58 (*Musca*). Germany, Württemberg. T A SMN.  
*Muscidae*, *Pyrellia rapax* Harris. Preocc. Fabricius 1805.
- fuscipes* Gmelin, 1790: 2843 (*Musca*). Europe. T A NMI.
- fuscula* Fallén, 1825: 86 (*Musca*). Sweden, Skåne, Äsperöd. LT ♂ ZIL. Fanniidae,  
*Fannia fuscula* Fallén. Lectotype designated by Pont 1984: 285.
- fucus* Scopoli, 1763: 355 (*Conops*). N. n. *Conops interruptus* Poda. Syrphidae,  
*Eristalis interrupta* Poda.
- fucus* Harris, 1776: 42, pl. 10, fig. 3 (*Musca*). England. T A Unknown. Syrphidae,  
*Eristalis intricaria* Linnaeus.
- fusiformis* Turton, 1801: 642 (*Musca*). N. n. *Musca fuciformis* Fabricius. Syrphidae,  
*Mallota fuciformis* Fabricius.
- gabonensis* Macquart, 1855: 135 (*Musca*). Gabon. HT ♂ BMNH. Muscidae, *Musca gabonensis* Macquart. See Pont 1990: 165.
- galeata* Rossi, 1794: 73 (*Musca*). Italy. T A ZMHU.
- galpinus* Harris, 1780: 118, pl. 35, fig. 28 (*Musca*). England. T A Unknown.  
Opomyzidae.
- gamelia* Walker, 1849: 878 (*Musca*). Uruguay, Montevideo. T ♂ BMNH. Calliphoridae, *Compsomyiops fulvicrura* Robineau-Desvoidy.
- gangraenosa* Panzer, 1798b: 22 (*Musca*). Austria. T A ZMHU. Otitidae, *Melieria gangraenosa* Panzer. N. Status
- gastrodes* Rossi, 1794: 67, pl. 3, fig. 11 (*Musca*). Italy. T A ZMHU. Tachinidae.
- gavisa* Walker, 1859: 138 (*Musca*). Sulawesi, Makassar. T ♀ BMNH. Muscidae,  
*Neomyia gavisa* Walker.
- genarum* Zetterstedt, 1837: 42 (*Musca*). *Nomen nudum*. [Sweden, Torne Lappmark, Torneträsk]. T ♂ ZIL. Calliphoridae, *Calliphora genarum* Zetterstedt.
- genarum* Zetterstedt, 1838: 658 (*Musca*). Sweden, Torne Lappmark, Torneträsk. LT ♂ ZIL. Calliphoridae, *Calliphora genarum* Zetterstedt. Lectotype designated by Rognes 1986: 352.
- geniculata* De Geer, 1776: 38, pl. 2, fig. 19 (*Musca*). Not given, probably Sweden.  
T A Lost. Tachinidae, *Siphona geniculata* De Geer. See Andersen 1982: 168.
- geniculata* Fallén, 1825: 89 (*Musca*). Sweden. LT ♂ NRS. Muscidae, *Coenosia femoralis* Robineau-Desvoidy. Preocc. De Geer 1776. Lectotype designated by Pont 1984: 285.
- gentilis* Meigen, 1838: 302 (*Musca*). Germany. T A MNHNP. Calliphoridae, *Melinda gentilis* Robineau-Desvoidy. N. Syn.
- geographica* Geoffroy, 1785: 478 (*Musca*). France, Paris. T A Author.
- germana* Walker, 1849: 880 (*Musca*). Subsequent Usage of *Lucilia germana* Robineau-Desvoidy. Calliphoridae, *Onesia germana* Robineau-Desvoidy.
- germanica* Gmelin, 1790: 2842 (*Musca*). N. n. *Musca plebeja* Fabricius. Tachinidae, *Dexia rustica?* Fabricius.
- germinationis* Linnaeus, 1758a: 600 (*Musca*). Europe. T A LSL. Opomyzidae,  
*Opomyza germinationis* Linnaeus. 2 sp. in LSL (Dear MS).
- germinationis* Rossi, 1790: 316 (*Musca*). Subsequent Usage of *Musca germinationis* Linnaeus. Opomyzidae, *Opomyza germinationis* Linnaeus.
- gibba* Müller, 1776: 175 (*Musca*). Iceland. T A DESTROY. Dolichopodidae,  
*Dolichopus plumipes* Scopoli.
- gibba* Rossi, 1794: 73 (*Musca*). Italy. T A ZMHU. Preocc. Müller 1776.

- gibba** Fabricius, 1805: 297 (*Musca*). South America. T A UZMC. Curtonotidae,  
*Curtonotum taeniatum* Hendel. Preocc. Müller 1776. N. Syn.
- gibbosa** Linnaeus, 1758a: 593 (*Musca*). Sweden. T A LSL. Acroceridae, *Ogcodes gibbosa* Linnaeus. 1 sp. in LSL (Dear MS).
- gibbosa** De Geer, 1776: 35, pl. 2, fig. 1 (*Musca*). Sweden. T A NRS. Chamaemyiidae. Preocc. Linnaeus 1758.
- gibbosa** Thunberg, 1789: 90 (*Musca*). *Nomen nudum*. Proposed as n. n. *Hippobosca cincta* Thunberg. Preocc. Linnaeus 1758.
- gibsoni** Patton & Cragg, 1913a: 14, pl. 1 (*Musca*). India, Kodaikanal. LT ♂ BMNH. Muscidae, *Musca convexifrons* Thomson. Lectotype designated by Pont 1990: 165.
- gigantea** Lichtenstein, 1796: 213 (*Musca*). Unknown. T A Auction.
- glabrata** Fabricius, 1794: 341 (*Musca*). France. T A MNHNP. Lauxaniidae, *Calliopomyia glabrata* Fabricius.
- glabricula** Fallén, 1825: 75 (*Musca*). Sweden. LT ♂ NRS. Muscidae, *Hydrotaea glabricula* Fallén. Lectotype designated by Pont 1984: 285.
- glaucia** Linnaeus, 1758a: 593 (*Musca*). Sweden. LT ♀ LSL. Syrphidae, *Leucozona glaucia* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 156.
- glaucoptera** Gmelin, 1790: 2861 (*Musca*). Europe. T A NMI.
- globosa** Gmelin, 1790: 2875 (*Musca*). Subsequent Usage of *Syrphus globosus* Fabricius. Tachinidae, *Gymnosoma globosum* Fabricius.
- globulipes** Geoffroy, 1785: 495 (*Musca*). France, Paris. T A Author. Empididae, *Hilara globulipes* Geoffroy. N. Comb.
- gloriosa** Walker, 1858b: 104 (*Musca*). Moluccas, Aru Islands. HT ♀ BMNH. Calliphoridae, *Stilbomyella gloriosa* Walker. See Crosskey 1965: 95.
- graeca** Pontoppidan, 1765: 229 (*Musca*). Denmark. T A Unknown. Stratiomyidae, *Oxycera trilineata* Linnaeus. Suppressed by I.C.Z.N. 1987: 149.
- graminum** Fabricius, 1775: 785 (*Musca*). Germany, Leipzig. T A UZMC F. Otitidae.
- graminum** Fabricius, 1794: 346 (*Musca*). France, Paris. T A Author. Otitidae, *Doryceria aquatica* Geoffroy. Preocc. Fabricius 1775. N. Syn.
- grandis** Schrank, 1837: 17 (*Musca*). Germany, Burghusii [= Italy, Burgursio]. T A Unknown. Xylophagidae, *Coenomyia ferruginea* Scopoli.
- granditarsa** Forster, 1770: 15 (*Musca*). *Nomen nudum*. England. Syrphidae, *Platycheirus granditarsus* Forster. Published without diagnosis or indication.
- granditarsa** Forster, 1771: 99 (*Musca*). England. T A LSL? Syrphidae, *Platycheirus granditarsus* Forster. See Day & Fitton 1977.
- greeni** Patton, 1933d: 477, fig. 1 (*Musca*). Malaya, Kuala Lumpur. LT ♂ BMNH. Muscidae, *Musca formosana* Malloch. Lectotype designated by Pont 1990: 166.
- grisea** de Villers, 1789a: 511 (*Musca*). Sweden, Skåne. T A Unknown.
- grisea** Fabricius, 1805: 293 (*Musca*). South America. LT ♀ UZMC. Muscidae, *Cyrtoneurina spiloptera* Wiedemann. Preocc. de Villers 1789. Lectotype designated by Michelsen 1979a: 185.
- grisea** Fallén, 1823: 57 (*Musca*). Sweden, Västergötland. LT ♂ NRS. Anthomyiidae, *Leucophora grisella* Hennig. Preocc. de Villers 1789. Lectotype designated by Michelsen 1983: 113.
- grisella** Robineau-Desvoidy, 1863: 622 (*Musca*). France, Paris. T A DESTROY. Muscidae, *Musca autumnalis* De Geer.

- grisescens** Fallén, 1817: 243 (*Musca*). Sweden. T ♂ NRS. Tachinidae, *Dinera grisescens* Fallén.
- groenlandica** Zetterstedt, 1838: 657 (*Musca*). Norway, Greenland (Lectotype) & Sweden. LT ♂ UZMC. Calliphoridae, *Protophormia terraenovae* Robineau-Desvoidy. Lectotype designated by Rognes 1991a: 136.
- grossa** Linnaeus, 1758a: 596 (*Musca*). Europe. TA LSL. Tachinidae, *Tachina grossa* Linnaeus. 2 sp. in LSL (Dear MS).
- grossificationis** Linnaeus, 1758a: 599 (*Musca*). Europe. HT ♀ LSL. Rhinophoridae, *Melanophora roralis* Linnaeus. See Crosskey 1977: 51.
- grossipes** Linnaeus, 1767: 988 (*Musca*). Europe. T A LSL. Empididae, *Hybos grossipes* Linnaeus. 1 sp. in LSL (Dear MS).
- guineensis** Wiedemann, 1824: 48 (*Musca*). Guinea. T ♂ DESTROY. Muscidae, *Neomyia guineensis* Wiedemann.
- gulo** Fabricius, 1805: 283 (*Musca*). South America. HT ♀ UZMC. Sarcophagidae, *Paraprisopoda gulo* Fabricius. See Townsend 1931a: 74; Pape 1986: 302.
- gurnei** Patton & Cragg, 1912: 513 (*Musca*). India, Kodaikanal. LT ♂ BMNH. Muscidae, *Musca inferior* Stein. Lectotype designated by Pont 1990: 166.
- gurneyi** Patton, 1922: 422 (*Musca*). Missp. *gurnei* Patton & Cragg. Muscidae, *Musca inferior* Stein.
- guttata** Geoffroy, 1785: 495 (*Musca*). France, Paris. T A Author.
- gymnosomea** Rondani, 1862: 222 (*Musca*). Malta. T A USB. Muscidae, *Musca domestica* Linnaeus.
- haemorrhoa** Gravenhorst, 1807: 378 (*Musca*). Unknown, probably Germany. T A Lost. Sarcophagidae.
- haemorrhoidalis** de Villers, 1789a: 548, pl. 10, fig. 8 (*Musca*). France, Nemausum [= Nemours]. T A Author. Tephritidae.
- haemorrhoidalis** Fallén, 1817: 237 (*Musca*). Sweden. LT ♂ NRS. Sarcophagidae, *Ravinia pernix* Harris. Preocc. de Villers 1789. Lectotype designated by Pape 1986: 304.
- halterata** Panzer, 1798a: 13 (*Musca*). Austria. T A ZMHU. Sarcophagidae, *Nyctia halterata* Panzer.
- hamata** Fallén, 1823: 61 (*Musca*). Sweden, Skåne, Äsperöd. LT ♂ NRS. Muscidae, *Drymeia hamata* Fallén. Lectotype designated by Pont 1984: 285.
- harpyia** Harris, 1835: 599 (*Musca*). *Nomen nudum*. Massachusetts. T A MCZ. Muscidae, *Musca domestica* Linnaeus. Published without diagnosis or indication.
- harpyia** Harris, 1841: 413 (*Musca*). New England. T A MCZ. Muscidae, *Musca domestica* Linnaeus.
- hartigii** Ratzeburg, 1844: 172 (*Musca*). Germany. T A Unknown. Tachinidae, *Goniocera versicolor* Fallén.
- hauriens** Wiedemann, 1830: 398 (*Musca*). Egypt. T ♀ SMF. Sarcophagidae.
- hebes** Harris, 1780: 158, pl. 47, fig. 4 (*Musca*). England. T A Unknown. Dolichopodidae.
- hebes** Walker, 1853b: 338 (*Musca*). Senegal? T ♀ BMNH. Calliphoridae, *Chrysomya chloropyga* Wiedemann. Preocc. Harris 1780.
- hederacea** Geoffroy, 1785: 491 (*Musca*). France, Paris. T A Author. Calliphoridae, *Onesia hederacea* Geoffroy. N. Comb.
- heidiae** Zielke, 1974: 5 (*Musca*). Tanzania, Manyara Nat. Park, Res. Camp. HT ♂ CAS. Muscidae, *Musca heidiae* Zielke.

- helenii** de Villers, 1789a: 536 (*Musca*). N. n. *Musca duplicata* Geoffroy.
- helianthi** Rossi, 1794: 73 (*Musca*). Italy. T A ZMHU. Tephritidae, *Acanthiophilus helianthi* Rossi.
- helluo** Fabricius, 1805: 295 (*Musca*). Austria. T A UZMC F. Tachinidae, *Heliozeta helluo* Fabricius. "HT not seen" Townsend 1931c: 387.
- helvola** Meuschen, 1787: 212 (*Musca*). Unknown. T A Auction.
- hemiptera** Turton, 1801: 639 (*Musca*). Subsequent Usage of *Syrphus hemipterus* Fabricius. Tachinidae, *Allophora hemiptera* Fabricius.
- hemipterus** Donovan, 1807: 95 (*Musca*). Subsequent Usage of *Syrphus hemipterus* Fabricius. Tachinidae, *Allophora hemiptera* Fabricius.
- hemisphaerica** Geoffroy, 1785: 480 (*Musca*). France, Paris. T A Author. Tachinidae, *Gymnosoma rotundatum* Linnaeus.
- heraclei** Fabricius, 1794: 354 (*Musca*). Germany, Kiel. T A UZMC F. Tephritidae, *Tephritis postica* Loew.
- heraclii** Linnaeus, 1758a: 600 (*Musca*). Europe. T A LSL. Tephritidae, *Euleia heraclii* Linnaeus. See White 1987: 103.
- heraea** Walker, 1849: 881 (*Musca*). North America? T ♂ BMNH. Muscidae, *Neomyia cornicina* Fabricius.
- hervei** Villeneuve, 1922: 335 (*Musca*). China & Tonkin. ST ♂ ♀ MNHNP. Muscidae, *Musca hervei* Villeneuve.
- hieracii** Fabricius, 1775: 787 (*Musca*). England. T A UZMC F. Sciomyzidae, *Trypetoptera punctulata* Scopoli.
- hilaris** Fallén, 1823: 57 (*Musca*). Sweden, Västergötland. LT ♀ NRS. Anthomyiidae, *Eustalomyia hilaris* Fallén. Lectotype designated by Michelsen 1983: 113.
- hilli** Johnston & Bancroft, 1920e: 35 (*Musca*). Australia, Queensland. LT ♂ QMB. Muscidae, *Musca terraereginae* Johnston & Bancroft. Lectotype designated by Pont 1973: 181.
- hilli** Johnston & Bancroft, 1920b: 23 (*Musca*). *Nomen nudum*. Australia, Queensland. Muscidae, *Musca terraereginae* Johnston & Bancroft. [also 1920c: 52, 1920d: 74].
- himella** Walker, 1849: 876 (*Musca*). Unknown. T ♂ DESTROY. Calliphoridae, *Chrysomya albiceps* Wiedemann.
- hirsuta** de Villers, 1789a: 427, pl. 9, fig. 11 (*Musca*). France, Massilia (=Marseille). T A Author. Bombyliidae, *Hemipenthes maurus* Linnaeus. Synonymy by Evenhuis (in litt.). N. Syn.
- hirsuta** de Villers, 1789a: 509 (*Musca*). Europe. T A Unknown. Acalyptrate. Preocc. de Villers 1789.
- hirsuta** Donovan, 1810: 60, pl. 490, fig. 2 (*Musca*). England. T ♀ UNKNOWN. Tachinidae, *Servillia lurida* Fabricius. Preocc. de Villers 1789.
- hirsutissima** de Razoumowsky, 1789: 230 (*Musca*). Switzerland. T A UNKNOWN. Syrphidae, *Volucella bombylans* Linnaeus.
- hirta** Drury, 1773: 109 (*Musca*). Jamaica. T A UNKNOWN. Tachinidae.
- hirta** de Villers, 1789a: 420 (*Musca*). Subsequent Usage of *Nemotelus hirta* De Geer. Therevidae, *Thereva nobilitata* Fabricius.
- hirta** Gravenhorst, 1807: 379 (*Musca*). Unknown, probably Germany. T A Lost. Tachinidae. Preocc. Drury 1773.
- hirticeps** Fallén, 1824: 67 (*Musca*). Sweden, Skåne. HT ♀ ZIL. Muscidae, *Hydrotaea hirticeps* Fallén. See Pont 1984: 285.

- hirticollis** Harris, 1835: 599 (*Musca*). *Nomen nudum*. Massachusetts. Published without diagnosis or indication.
- histrionica** Fabricius, 1794: 343 (*Musca*). India Orientali. T A UZMC F. Psilidae.
- hoi** Fan, 1965: 119, pl. 18, fig. 381 (*Musca*). China, Shensi, Tsuihuashan. HT ♂ SIE. Muscidae, *Musca pilifacies* Emden.
- hordei** Linnaeus, 1764c: 354 (*Musca*). *Nomen nudum*. Sweden. T A LSL. Chloropidae, *Oscinella frit* Linnaeus. Published in non-binomial work.
- hordei** Bjerkanter, 1777: 34 (*Musca*). Sweden. T A UNKNOWN. Chloropidae, *Oscinella frit* Linnaeus.
- hortensia** Wiedemann, 1824: 49 (*Musca*). Java. LT ♂ UZMC. Muscidae, *Morellia hortensia* Wiedemann. Lectotype designated by Pont 1973: 221.
- hortensis** Geoffroy, 1785: 481 (*Musca*). France, Paris. T A Author.
- horticola** De Geer, 1776: 140, pl. 8, fig. 12 (*Musca*). N. n. *Musca nemorum* Linnaeus. Syrphidae, *Eristalis arbustorum* Linnaeus.
- hortona** Walker, 1849: 894 (*Musca*). New Zealand. T ♀ BMNH. Calliphoridae, *Xenocalliphora hortona* Walker.
- hortorum** Gmelin, 1790: 2869 (*Musca*). Subsequent Usage of *Syrphus hortorum* Fabricius. Syrphidae, *Palpada hortorum* Fabricius.
- hortorum** Fallén, 1817: 252 (*Musca*). Sweden. LT ♂ NRS. Muscidae, *Morellia hortorum* Fallén. Lectotype designated by Pont 1984: 286.
- hortulana** Linnaeus, 1758b: (*Musca*). Subsequent Usage of *Tipula hortulana* Linnaeus. Bibionidae, *Bibio hortulanus* Linnaeus. [also Linnaeus 1760: 244].
- hortulana** Scopoli, 1763: 343 (*Musca*). Carniola. T A DESTROY. Syrphidae, *Cheilosia hortulana* Scopoli.
- hortulana** Rossi, 1790: 313 (*Musca*). Italy, Tuscany, Firenze & Pisa. T A ZMHU. Otitidae, *Ceroxys hyalinata* Panzer. Preocc. Scopoli 1763. N. Syn.
- hortulana** Wiedemann, 1830: 417 (*Musca*). China. ST ♀ DESTROY. Muscidae, *Morellia hortensia* Wiedemann. Preocc. Scopoli 1763. See Emden 1965: 110.
- hottentotta** Robineau-Desvoidy, 1830: 399 (*Musca*). South Africa, Cape of Good Hope. T ♂ DESTROY. Muscidae, *Musca domestica* Linnaeus. Preocc. Linnaeus 1758.
- hottentotta** Linnaeus, 1758a: 590 (*Musca*). Europe. T A LSL. Bombyliidae, *Anthrax hottentotta* Linnaeus. 2 sp. in LSL (Dear MS).
- hugonis** Pont, 1980: 725 (*Musca*). N. n. *Musca longipes* Paterson. Muscidae, *Musca hugonis* Pont.
- humilis** Wiedemann, 1830: 418 (*Musca*). East Indies. LT ♂ NMW. Muscidae, *Musca sorbens* Wiedemann. Lectotype designated by Pont 1973: 287.
- hyalinata** Panzer, 1798c: 24 (*Musca*). Austria. T A ZMHU. Otitidae, *Ceroxys hyalinata* Panzer. N. Status
- hyalinata** Fallén, 1823: 64 (*Musca*). Sweden. LT ♂ NRS. Muscidae, *Thricops semicinereus* Wiedemann. Preocc. Panzer 1798. Lectotype designated by Pont 1984: 286.
- hydroleon** Linnaeus, 1758a: 589 (*Musca*). Europe. T A LSL. Stratiomyidae, *Odonomyia hydroleon* Linnaeus. 1 sp. in LSL (Dear MS).
- hydrophila** Rossi, 1794: 74 (*Musca*). Italy. T A ZMHU.
- hydropica** Scudder, 1877: 757 (*Musca*). Colorado, White River [Oligocene]. T X USNM.

- hyoscyami** Linnaeus, 1758a: 600 (*Musca*). Europe. T A LSL. Tephritidae, *Tephritis hyoscyami* Linnaeus. See White 1987: 103.
- hypoleon** Linnaeus, 1767: 980 (*Musca*). Germany. T A LSL. Stratiomyidae, *Oxycera trilineata* Linnaeus. 3 sp. in LSL (Dear MS).
- hystric** Fabricius, 1775: 777 (*Musca*). N. n. *Musca pilosa* Drury. Tachinidae, *Archytas pilosus* Drury.
- iacobae** Panzer, 1805: 22 (*Musca*). Incosp. *jacobae* Panzer. Tephritidae, *Urophora stylata* Fabricius. Name status by present revision.
- icela** Walker, 1849: 897 (*Musca*). New Zealand. T ♀ BMNH. Calliphoridae, *Xenocalliphora hortona* Walker.
- ichneumonea** Linnaeus, 1758a: 593 (*Musca*). Europe. T A LSL. Psilidae, *Loxocera ichneumonea* Linnaeus.
- ichneumonoides** Geoffroy, 1785: 498 (*Musca*). France, Paris. T A Author.
- idessa** Walker, 1849: 908 (*Musca*). Ontario, St. Martin's Falls. HT ♀ BMNH. Muscidae, *Graphomya idessa* Walker. See Arntfield 1975: 282.
- ieiuna** Fabricius, 1787: 342 (*Musca*). India, Tranquebar. HT ♀ UZMC. Calliphoridae, *Bengalia ieiuna* Fabricius. See Townsend 1931c: 371.
- ignavus** Harris, 1780: 154, pl. 45, fig. 91 (*Musca*). England. T ♀ Unknown. Muscidae, *Hydrotaea ignava* Harris.
- ilerda** Walker, 1849: 895 (*Musca*). (Ontario, St. Martin's Falls). T ♀ BMNH. Calliphoridae, *Cynomya cadaverina* Robineau-Desvoidy.
- iliaca** Ratzeburg, 1840: 125 (*Musca (Tachina)*). Germany. T A UNKNOWN. Tachinidae.
- illingworthi** Patton, 1923b: 323 (*Musca*). Java, Buitenzorg. ST A BBM. Muscidae, *Musca illingworthi* Patton. See Pont 1990: 166 (STs also in Manchester Museum).
- illucens** Linnaeus, 1758a: 589 (*Musca*). South America. T A NRS. Stratiomyidae, *Hermetia illucens* Linnaeus.
- illustratus** Harris, 1780: 104, pl. 32, fig. 32 (*Musca*). England. T A Unknown. Syrphidae, *Cheilosia illustrata* Harris.
- illustris** Meigen, 1826: 54 (*Musca*). Germany. ST ♂ ♀ NMW. Calliphoridae, *Lucilia illustris* Meigen. See Collin 1926: 258.
- imbellis** Harris, 1776: 60, pl. 15, fig. 16 (*Musca*). England. T A Unknown. Syrphidae, *Chrysotoxum arcuatum* Linnaeus.
- impavida** Rossi, 1790: 304 (*Musca*). Italy, Tuscany, Firenze & Pisa. T A ZMHU. Calliphoridae, *Rhyncomya impavida* Rossi.
- importuna** Haliday, 1836: 149 (*Musca*). Ireland, Down, Holywood. T ♂ NMI. Muscidae, *Morellia hortorum* Fallén.
- impuncta** Fallén, 1825: 79 (*Musca*). Sweden. LT ♂ NRS. Muscidae, *Helina impuncta* Fallén. Lectotype designated by Pont 1984: 286.
- inaequalis** Geoffroy, 1785: 483 (*Musca*). France, Paris. T A Author. Syrphidae, *Episyrphus cannabina* Scopoli.
- inanis** Linnaeus, 1758a: 595 (*Musca*). Sweden. LT ♂ LSL. Syrphidae, *Volucella inanis* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 156.
- inanis** Fabricius, 1775: 763 (*Syrphus*). West Indies. T A UZMC F. Syrphidae, *Copestylum brunneum* Thunberg. Preocc. Linnaeus 1758.
- inanis** Fallén, 1825: 91 (*Musca*). Sweden, Skåne. LT ♂ NRS. Muscidae, *Achanthiptera rohrelliformis* Robineau-Desvoidy. Preocc. Linnaeus 1758. Lectotype designated by Pont 1984: 287.

- incerta* Walker, 1853b: 334 (*Musca*). Columbia. HT ♀ BMNH. Calliphoridae, *Sarconesia magellanica* Le Guillou. See Dear 1979: 161.
- incerta* Patton, 1922b: 71 (*Musca*). India, Madras, Saidapet. LT ♂ BMNH. Muscidae, *Musca pattoni* Austen. Preocc. Walker 1853. Lectotype designated by Pont 1990: 166.
- inconclusa* Walker, 1860: 160 (*Musca*). Moluccas, Amboina. T ♂ BMNH. Calliphoridae, *Polleniopsis inconclusa* Walker.
- inconstans* Wiedemann, 1830: 672 (*Musca*). *Nomen nudum*. Muscidae, *Musca crassirostris* Stein. Published without diagnosis or indication.
- incurvata* Bouché, 1834: 68 (*Musca*). Germany. T A UNKNOWN. Muscidae, *Myospila meditabunda* Fabricius.
- inda* Schrank, 1781: 438 (*Musca*). Austria, Vienna & Linz. T A Unknown. Stratiomyidae, *Clitellaria ephippium* Fabricius.
- indica* Walker, 1849: 880 (*Musca*). Subsequent Usage of *Lucilia indica* Robineau-Desvoidy. Muscidae, *Neomyia indica* Robineau-Desvoidy.
- indica* Awati, 1916b: 138 (*Musca*). India, Delhi, Kasauli & S. India. ST ♂ ♀ CRIK. Muscidae, *Musca planiceps* Wiedemann. Preocc. Harris 1776.
- indicus* Harris, 1776: 47, pl. 11, fig. 77 (*Musca*). England. T A Unknown. Stratiomyidae, *Sargus iridatus* Scopoli.
- inducta* Walker, 1853b: 335 (*Musca*). East Indies. T ♀ BMNH. Calliphoridae, *Lucilia papuensis* Macquart.
- inferior* Stein, 1909: 213 (*Musca*). Java, Semarang. LT ♂ ITZA. Muscidae, *Musca inferior* Stein. Lectotype designated by Pont 1970a: 89.
- infixa* Walker, 1856a: 25 (*Musca*). Singapore. T ♀ BMNH. Calliphoridae, *Hypopygiopsis violacea* Macquart.
- inflata* Turton, 1801: 637 (*Musca*). Subsequent Usage of *Syrphus inflatus* Fabricius. Syrphidae, *Volucella inflata* Fabricius.
- ingens* Walker, 1859b: 134 (*Musca*). Sulawesi, Makasar. T ♂ BMNH. Calliphoridae, *Catapicephala ingens* Walker.
- ingredior* Harris, 1780: 140, pl. 41, fig. 33 (*Musca*). England. T ♂ Unknown. Muscidae, *Neomyia cornicina* Fabricius. Name improperly formed (verb).
- ingruo* Harris, 1780: 144, pl. 42, fig. 53 (*Musca*). England. T A Unknown. Dolichopodidae, *Sciapus* sp. Name improperly formed (verb).
- ingrus* Harris, 1780: 143 (*Musca*). Incosp. *ingruo* Harris. Dolichopodidae, *Sciapus ingruo* Harris. Name status by present revision.
- inscribens* Walker, 1859b: 136 (*Musca*). Sulawesi, Makassar. T ♀ BMNH. Calliphoridae, *Lucilia papuensis* Macquart.
- insignis* Austen, 1909: 298, fig. 1 (*Philaematomyia*). India. LT ♂ BMNH. Muscidae, *Musca crassirostris* Stein. Lectotype designated by Pont 1990: 166.
- insularis* Walker, 1853b: 340 (*Musca*). West Indies. T ♀ BMNH. Calliphoridae, *Lucilia eximia* Wiedemann.
- intermedia* Fallén, 1825: 87 (*Musca*). Sweden. LT ♂ NRS. Muscidae, *Coenosia intermedia* Fallén. Lectotype designated by Pont 1984: 287.
- intermedia* Meigen, 1838: 306 (*Musca*). Subsequent Usage of *Pollenia intermedia* Macquart. Calliphoridae, *Pollenia intermedia* Macquart.
- interpuncta* Fallén, 1823: 63 (*Musca*). *Nomen nudum*. Muscidae, *Helina impuncta* Fallén. Published without diagnosis or indication. Mentioned under *quadrimaculata*, error for *impuncta*.

- interpunctionis** Ratzeburg, 1844: 170 (*Musca*). Germany. T A Unknown. Tachinidae.
- interpunctus** Harris, 1776: 59, pl. 15, fig. 13 (*Musca*). England. T ♀ Unknown. Syrphidae, *Lejops interpunctus* Harris.
- interrupta** Geoffroy, 1785: 482 (*Musca*). France, Paris. T A Author. Syrphidae.
- interrupta** Gmelin, 1790: 2879 (*Musca*). Europe. T A NMI. Syrphidae, *Epistrophe eligans* Harris. Preocc. Geoffroy 1785.
- interrupta** Walker, 1853b: 343 (*Musca*). Cape. HT ♂ BMNH. Muscidae, *Musca interrupta*.
- interruptella** Pont. Preocc. Geoffroy 1785. See Pont 1990: 166.
- interruptella** Pont, 1980: 725 (*Musca*). N. n. *Musca interrupta* Walker. Muscidae, *Musca interruptella* Pont.
- interruptus** Poda, 1761: 118 (*Conops*). Austria, Graz. T A Unknown. Syrphidae, *Eristalis interrupta* Poda.
- intersecta** Geoffroy, 1785: 479 (*Musca*). France, Paris. T A Author. Syrphidae.
- interventum** Harris, 1780: 144, pl. 42, fig. 57 (*Musca*). England. T A Unknown. Rhinophoridae, *Melanophora roralis* Linnaeus.
- interventus** Harris, 1780: 145, pl. 42, fig. 60 (*Musca*). England. T A Unknown. Tachinidae.
- intrahens** Walker, 1859: 137 (*Musca*). Sulawesi, Makassar. T ♀ BMNH. Muscidae, *Neomyia timorensis* Robineau-Desvoidy.
- intricaria** Linnaeus, 1758a: 592 (*Musca*). Sweden. LT ♀ LSL. Syrphidae, *Eristalis intricaria* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 156.
- intricata** Turton, 1801: 638 (*Musca*). N. n. *Musca intricaria* Linnaeus. Syrphidae, *Eristalis intricaria* Linnaeus.
- investigator** Harris, 1776: 35, pl. 9, fig. 3 (*Musca*). England. T A Unknown. Tachinidae, *Linnaemya vulpina?* Fallén.
- invisito** Harris, 1780: 83, pl. 24, fig. 31 (*Musca*). England. T A Unknown. Syrphidae, *Sphaerophoria scripta* Linnaeus. Name improperly formed (verb).
- ionia** Meigen, 1975: 435, pl. 199, fig. 9 (*Musca*). *Nomen nudum*. Germany, probably Stolberg. Calliphoridae, *Lucilia* sp. Published after 1930 without a description.
- iridata** Scopoli, 1763: 340 (*Musca*). Carniola. T A DESTROY. Stratiomyidae, *Sargus iridatus* Scopoli.
- irritans** Harris, 1780: 148 (*Musca*). Subsequent Usage of *Conops irritans* Linnaeus. Muscidae, *Haematobia irritans* Linnaeus.
- irritans** Fallén, 1823: 62 (*Musca*). Sweden. LT ♂ NRS. Muscidae, *Hydrotaea irritans* Fallén. Lectotype designated by Pont 1984: 287.
- irwinrossi** Zielke, 1974: 7 (*Musca*). Tanzania, Manyara Nat. Park, Res. Camp, Ndala River. HT ♂ CAS. Muscidae, *Musca xanthomelaena* Wiedemann.
- jacobeae** Panzer, 1805: 22 (*Musca*). Germany. T A UNKNOWN. Tephritidae, *Urophora stylata* Fabricius.
- jacobsoni** Malloch, 1928: 334 (*Bromya*). Sumatra, Fort de Kock. HT ♂ ITZA. Muscidae, *Musca fletcheri* Patton & Senior-White.
- jamaicensis** Gmelin, 1790: 2871 (*Musca*). N. n. *Syrphus nigrita* Fabricius. Syrphidae, *Sterphus jamaicensis* Gmelin.
- japonica** Thunberg, 1789: 90 (*Musca*). Japan. T A ZIU. Stratiomyidae, *Pteicticus japonica* Thunberg. N. Status
- jeiuna** Fabricius, 1794: 312 (*Musca*). emend. *ieiuna* Fabricius. Calliphoridae, *Bengalia ieiuna* Fabricius.

- jejuna* Fabricius, 1805: 283 (*Musca*). emend. *ieiuna* Fabricius. Calliphoridae, *Bengalia ieiuna* Fabricius.
- jejuna* Schrank, 1837: 17 (*Musca*). Austria, Gern. T A Unknown. Stratiomyidae, *Oplodontha viridula* Fabricius. Preocc. Fabricius 1787.
- joco* Harris, 1780: 157, pl. 47, fig. 1 (*Musca*). England. T A Unknown. Dolichopodidae, *Poecilobothrus nobilitatus* Linnaeus.
- kasauliensis* Awati, 1916b: 138 (*Musca*). India, Kasauli. ST ♂ ♀ CRIK. Muscidae, *Musca ventrosa* Wiedemann.
- kihuris* Zielke, 1971c: 127 (*Musca*). Tanzania, Kihurio. ST ♂ ♀ UNKNOWN. Muscidae, *Musca lusoria* Wiedemann.
- kweilinensis* Ôuchi, 1938b: 31 (*Musca*). China, Kwangsi, Kweilin. HT ♂ SIE. Muscidae, *Musca conducens* Walker.
- labiata* Fabricius, 1787: 346 (*Musca*). Denmark, Copenhagen. T A UNKNOWN. Sarcophagidae, *Metopia labiata* Fabricius. See Pape 1986: 302.
- lacustris* Scopoli, 1763: 343 (*Musca*). Carniola. T A DESTROY. Dolichopodidae, *Orthoceratium lacustre* Scopoli.
- laemica* White, 1843: 291 (*Musca*). New Zealand. LT ♀ BMNH. Calliphoridae, *Calliphora stygia* Fabricius. Lectotype designated by Dear 1985a: 25.
- laemica* Walker, 1849: 906 (*Musca*). New Zealand (Lectotype) & Australia. LT ♀ BMNH. Calliphoridae, *Calliphora stygia* Fabricius. Preocc. White 1843. Lectotype designated by us here: LT of *laemica* White = LT of Walker.
- laeta* Turton, 1801: 647 (*Musca*). Subsequent Usage of *Syrphus laetus* Fabricius. Syrphidae, *Xanthogramma laeta* Fabricius.
- laeta* Fallén, 1823: 56 (*Musca*). Sweden, Skåne, Bäckaskog. LT ♀ NRS. Muscidae, *Phaonia laeta* Fallén. Lectotype designated by Pont 1984: 287.
- lagyra* Walker, 1849: 885 (*Musca*). Azores, Fayal. T ♀ BMNH. Calliphoridae, *Lucilia sericata* Meigen.
- lamed* Schrank, 1781: 475 (*Musca*). Austria. T A Unknown. Otitidae, *Otites lamed* Schrank.
- lancifer* Harris, 1780: 126, pl. 36, fig. 59 (*Musca*). England. T A Unknown. Anthomyiidae, *Hydrophoria lancifer* Harris.
- laniaria* Wiedemann, 1830: 683 (*Musca*, as *macellaria* var.). Brazil. T A Lost. Calliphoridae, *Cochliomyia macellaria* Fabricius. ST not found in ZMHU (Dear 1985a: 141).
- lanio* Panzer, 1798a: 11 (*Musca*). Austria. T A ZMHU. Calliphoridae, *Pollenia* sp.
- lappae* Cederhielm, 1798: 319 (*Musca*). USSR, Ingrija [= St. Petersburg Area]. T A UNKNOWN. Tephritidae, *Terellia lappae* Cederhielm.
- lappona* Linnaeus, 1758a: 591 (*Musca*). Sweden. LT ♂ LSL. Syrphidae, *Sericomyia lappona* Linnaeus. Lectotype designated by Thompson et al. 1982: 157.
- lar* Gmelin, 1790: 2832 (*Musca*). Subsequent Usage of *Bibio lar* Fabricius. Bombyliidae, *Litorhina lar* Fabricius.
- lardaria* Fabricius, 1781: 436 (*Musca*). Germany. T A DESTROY. Muscidae, *Polistes lardaria* Fabricius. See Michelsen 1979a: 185.
- larvarum* Linnaeus, 1758a: 596 (*Musca*). Europe. T A LSL. Tachinidae, *Exorista larvarum* Linnaeus.
- larvarum* Scopoli, 1763: 331 (*Musca*). Carniola. ST L DESTROY. Tachinidae. Four varieties described, each from different Lepidopteran host.

- larvata* Lepechin, 1774: 310, pl. 20, fig. 2 (*Musca*). *Nomen nudum*. [Russia]. Published in non-binominal work.
- larvata* Schrank, 1781: 452 (*Musca*). Austria, Upper Austria, Eberspergae [= Ebersberg]. T A Unknown.
- larvata* Geoffroy, 1785: 475 (*Musca*). France, Paris. T A Author. Preocc. Schrank 1781.
- larvata* Gmelin, 1790: 2845 (*Musca*). USSR, Urals. T A Unknown. Preocc. Schrank 1781.
- larvincola* Ratzeburg, 1844: 173 (*Musca*). Germany. Tachinidae, *Exorista larvarum* Linnaeus.
- larvipara* Portschinsky, 1910: 29, fig. 30 (*Musca*). Southern Russia. ST ♂ ♀ ASL. Muscidae, *Musca larvipara* Portschinsky.
- lasciuia* Fabricius, 1798: 564 (*Musca*). Cayenne. T A UZMC F. Micropezidae, *Taeniacaptera lasciva* Fabricius.
- lasiopa* Villeneuve, 1936: 414 (*Musca*). Kenya, Mt Kenya. ST ♀ BMNH. Muscidae, *Musca lasiopa* Villeneuve. Syntypes also at MRAC. See Pont 1990: 166.
- lasiophthalma* Thomson, 1868: 548 (*Musca*). South Africa, Cape of Good Hope. T A NRS. Muscidae, *Musca lasiophthalma* Thomson.
- lateralis* Linnaeus, 1758a: 597 (*Musca*). Europe. LT ♂ LSL. Fanniidae, *Fannia canicularis* Linnaeus. Suppressed by I.C.Z.N. 1969:124. Lectotype designated by Pont 1981: 170.
- lateralis* Fabricius, 1775: 778 (*Musca*). Germany. T A UZMC F. Tachinidae, *Eriothrix rufomaculata* De Geer. Preocc. Linnaeus 1758.
- lateralis* Geoffroy, 1785: 496 (*Musca*). Subsequent Usage of *Musca lateralis* Linnaeus. Fanniidae, *Fannia canicularis* Linnaeus.
- lateralis* Donovan, 1792: 69 (*Musca*). England. T A Author. Tachinidae, *Eriothrix rufomaculata* De Geer. Preocc. Linnaeus 1758. N. Syn.
- lateralis* Fabricius, 1805: 286 (*Musca*). South America. HT ♂ UZMC. Tachinidae, *Opsozelia discalis* Townsend. Preocc. Linnaeus 1758. See Townsend 1931a: 96.
- lateralis* Fallén, 1817: 241 (*Musca*). Sweden, Skåne. T A NRS. Tachinidae, *Thelaira nigrina* Fallén. Preocc. Linnaeus 1758.
- lateralis* Macquart, 1834: 158 (*Musca*). France, Lille. T ♀ MHNL. Muscidae, *Musca domestica* Linnaeus. Preocc. Linnaeus 1758.
- lateritia* Schrank, 1796: 129 (*Musca*). T A UNKNOWN.
- laternaria* Müller, 1776: 174 (*Musca*). Denmark. T A Author. Syrphidae, *Leucozona laternaria* Müller.
- laticornis* Fallén, 1825: 86 (*Musca*). Sweden, Skåne. HT ♀ ZIL. Anthomyiidae, *Pegomya laticornis* Fallén. See Michelsen 1983: 113.
- laticornis* Zetterstedt, 1837: 42 (*Musca*). *Nomen nudum*. [Sweden, Bottnia occidentalis, Muonioniska]. T ♀ ZIL. Calliphoridae, *Calliphora genarum* Zetterstedt.
- laticornis* Zetterstedt, 1838: 658 (*Musca*). Sweden, Bottnia occidentalis, Muonioniska. LT ♀ ZIL. Calliphoridae, *Calliphora genarum* Zetterstedt. Preocc. Fallén 1825. Lectotype designated by Rognes 1986: 352.
- latifrons* Fallén, 1817: 238 (*Musca*). Sweden, Skåne, Äsperöd. LT ♂ NRS. Sarcophagidae, *Sarcophila latifrons* Fallén. Lectotype designated by Pape 1986: 305.
- latifrons* Wiedemann, 1830: 656 (*Musca*). China, Macao. LT ♀ NMW. Muscidae, *Musca vetustissima* Walker. Preocc. Fallén 1817. Lectotype designated by Pont 1973: 148.

- latiparafrons* Awati, 1916b: 138 (*Musca*). India. ST ♀ CRIK. Muscidae, *Musca convexifrons?* Thomson.
- latipes* Gmelin, 1790: 2861 (*Musca*). Europe. T A NMI.
- latrinarum* De Geer, 1776: 31, pl. 1, fig. 15 (*Musca*). N. n. *Musca serrata* Linnaeus. Heleomyzidae, *Heleomyza serrata* Linnaeus.
- latro* Fabricius, 1787: 349 (*Musca*). Europe. T A UZMC F.
- lauta* Wiedemann, 1830: 410 (*Musca*). Java. LT ♀ NMW. Muscidae, *Neomyia lauta* Wiedemann. Lectotype designated by Pont 1973: 192 (Townsend (1931c: 369) HT designation invalid).
- lineatus* Harris, 1780: index (*Musca*). Missp. *lineatus* Harris. Syrphidae, *Eristalis lineata* Harris.
- lenio* Harris, 1780: 147, pl. 43, fig. 66 (*Musca*). England. T A Unknown. Anthomyiidae, *Hydrophoria linogrisea* Meigen. Name improperly formed (verb).
- lenis* Harris, 1780: 154, pl. 45, fig. 88 (*Musca*). England. T A Unknown. Anthomyiidae.
- leonina* Fabricius, 1775: 776 (*Musca*). Australia. HT ♂ BMNH B. Calliphoridae, *Amenia leonina* Fabricius. See Townsend 1931c: 374 (HT in Kiel!); Crosskey 1965: 112.
- leontodontis* De Geer, 1776: 46, pl. 2, fig. 18 (*Musca*). N. n. *Musca hyoscyami* Linnaeus. Tephritidae, *Tephritis hyoscyami* Linnaeus.
- levida* Fallén, 1825: 76 (*Musca*). Subsequent Usage of *Anthomyia lepida* Wiedemann. Fanniidae, *Fannia lepida* Wiedemann.
- lepidus* Harris, 1780: 121, pl. 35, fig. 40 (*Musca*). England. T A Unknown. Chloropidae, *Chlorops lepidus* Harris. N. Status
- leporinus* Harris, 1780: 117, pl. 34, fig. 23 (*Musca*). England. T A Unknown. Acalyptrate.
- leprae* Linnaeus, 1758a: 598 (*Musca*). America. T A LSL.
- leptogaster* Panzer, 1798a: 19 (*Musca*). Austria. T A ZMHU. Psilidae, *Chyliza leptogaster* Panzer.
- leucocephala* de Villers, 1789a: 550, pl. 10, fig. 11 (*Musca*). France, Cevennes Mts. T A Author. Sarcophagidae.
- leucocephala* Gmelin, 1790: 2878 (*Musca*). Europe. T A NMI. Syrphidae, *Eristalinus aeneus* Scopoli. Preocc. de Villers 1789.
- leucocephala* Rossi, 1790: 306 (*Musca*). Italy, Tuscany, Firenze & Pisa. T A ZMHU. Sarcophagidae, *Metopia argyrocephala* Meigen. Preocc. de Villers 1789.
- leucocephala* Walker, 1853b: 339 (*Musca*). Senegal? T ♀ BMNH. Calliphoridae, *Hemipyrellia ferdinica* Macquart. Preocc. de Villers 1789.
- leucomelas* Wiedemann, 1824: 48 (*Musca*). South Africa, Cape of Good Hope. T ♀ UZMC. Muscidae, *Graphomya leucomelaena* Wiedemann.
- leuconyx* Rossi, 1794: 65 (*Musca*). Italy, "Etruria". T A ZMHU.
- leucopa* Linnaeus, 1767: 983 (*Musca*). America. T A LSL. Stratiomyidae, *Hermetia illucens* Linnaeus. 1 sp. in LSL (Dear MS).
- leucophaeus* Harris, 1782: 74 (*Musca*). Missp. *lucophaeus* Harris.
- leucopis* Gmelin, 1790: 2860 (*Musca*). Europe. T A NMI.
- leucoptera* Gmelin, 1790: 2861 (*Musca*). Europe. T A NMI.
- leucoptera* von Roser, 1840: 58 (*Musca*). Germany, Württemberg. T A SMN. Calliphoridae, *Pollenia* sp. Preocc. Gmelin 1790.
- leucosticta* Gmelin, 1790: 2860 (*Musca*). Non-European. T A NMI.

- leucozona** Panzer, 1806: 19 (*Musca*). Germany. T A UNKNOWN. Tachinidae, *Thelaira leucozona* Panzer.
- levidus** Harris, 1780: 124, pl. 36, fig. 52 (*Musca*). England. T A Unknown. Muscidae, *Muscina levida* Harris.
- libatrix** Scopoli, 1763: 346 (*Musca*). Carniola. T A DESTROY. Syrphidae.
- libatrix** Geoffroy, 1785: 498 (*Musca*). France, Paris. T A Author. Preocc. Scopoli 1763.
- libatrix** Panzer, 1798a: 12 (*Musca*). Austria. T A ZMHU. Tachinidae, *Zenillia fulva* Fallén. Preocc. Scopoli 1763. N. Syn.
- liberia** Snyder, 1951: 2 (*Musca*). Liberia, Robertsport, Near Bendu Village. HT ♂ AMNH. Muscidae, *Musca liberia* Snyder.
- ligurriens** Wiedemann, 1830: 655 (*Musca*). China & Java. ST ♂ UZMC NMW. Calliphoridae, *Hemipyrellia ligurriens* Wiedemann.
- lilaea** Walker, 1849: 894 (*Musca*). (Ontario, St. Martin's Falls). T ♀ BMNH. Calliphoridae, *Cynomya cadaverina* Robineau-Desvoidy.
- limatus** Harris, 1780: 122, pl. 36, fig. 45 (*Musca*). England. T A Unknown. Tephritidae.
- limbata** Gmelin, 1790: 2853 (*Musca*). N. n. *Musca marginata* Fabricius. Sciomyzidae, *Coremacera marginata* Fabricius.
- limbata** Manuel, 1811: 31 (*Musca*). Subsequent Usage of *Musca limbata* Gmelin. Sciomyzidae, *Coremacera marginata* Fabricius.
- lincensis** Schrank, 1776: 96, pl. 3, fig. 20 (*Musca*). Austria, Linz. T A Unknown. Tephritidae.
- lindneri** Paterson, 1956: 158, fig. 2 (*Musca*). Tanzania, Ngerengere. HT ♂ SMN. Muscidae, *Musca lindneri* Paterson.
- lineata** Fabricius, 1775: 754 (*Tipula*). England. T A BMNH. Keroplatidae, *Cerotelion striatus* Gmelin. N. Syn.
- lineata** Fabricius, 1781: 453 (*Musca*). Germany. T A UZMC F. Chloropidae, *Chlorops punilionis* Bjerkander. Preocc. Harris 1776.
- lineata** Geoffroy, 1785: 485 (*Musca*). France, Paris. T A Author. Syrphidae, *Eumerus strigatus* Fallén. Preocc. Harris 1776. N. Comb.
- lineata** Turton, 1801: 620 (*Musca*). Subsequent Usage of *Musca lineata* Fabricius. Chloropidae, *Chlorops punilionis* Bjerkander.
- lineata** Fallén, 1817: 238 (*Musca*). Sweden, Skåne, Bäckaskog. LT ♂ NRS. Sarcophagidae, *Blaesoxipha plumicornis* Zetterstedt. Preocc. Harris 1776. Lectotype designated by Pape 1986: 305.
- lineata** Brunetti, 1910: 91, pl. 8, fig. 19 (*Pristirhynchomyia*). India, W Bengal, Calcutta. ST ♀ ZSI. Muscidae, *Musca conducens* Walker. ST also in BMNH (Pont 1990: 167).
- lineatus** Harris, 1776: 42, pl. 10, fig. 4 (*Musca*). England. T A Unknown. Syrphidae, *Eristalis lineata* Harris.
- lineola** Turton, 1801: 634 (*Musca*). Subsequent Usage of *Rhagio lineola* Fabricius. Rhagionidae, *Rhagio lineola* Fabricius.
- lineolae** Harris, 1776: 58, pl. 15, fig. 10 (*Musca*). England. T A Unknown. Syrphidae, *Eristalis interrupta* Poda.
- lineolata** Gmelin, 1790: 2879 (*Musca*). Europe. T A NMI. Syrphidae, *Chrysogaster cemiteriorum* Linnaeus.

- lingens Meigen, 1975: 452, pl. 270, fig. 5 (*Musca*). *Nomen nudum*. Germany, probably Stolberg. Tachinidae, *Strongygaster globula* Meigen. Published after 1930 without a description.
- linzensis* Gmelin, 1790: 2863 (*Musca*). Emend. *lincensis* Schrank. Tephritidae.
- liris* Walker, 1849: 882 (*Musca*). Australia (?Tasmania). HT ♀ BMNH. Tachinidae, *Rutilia liris* Walker. See Crosskey 1973a: 136.
- lithophila* Germar, 1839: 222 (*Musca*). Germany, Bavaria, Solenhofen [Jurassic]. T X UNKNOWN. Blattariae, *Mesoblattina lithophila* Germar.
- litorea* Fallén, 1823: 63 (*Musca*). Sweden, Skåne, Äsperöd. LT ♂ NRS. Muscidae, *Spilogona litorea* Fallén. Lectotype designated by Pont 1984: 288.
- liturariae* Ratzeburg, 1844: 170 (*Musca*). Germany. T A Unknown. Anthomyiidae, *Delia antiqua* Meigen.
- liturata* Gmelin, 1790: 2861 (*Musca*). Europe. T A NMI.
- litus* Harris, 1780: 119, pl. 35, fig. 32 (*Musca*). England. T A Unknown. Anthomyiidae, *Anthomyia pluvialis* Linnaeus.
- livens* Fabricius, 1794: 345 (*Musca*). Denmark. T A UZMC. Scathophagidae, *Hydromyza livens* Fabricius.
- longerro* Harris, 1780: 152, pl. 45, fig. 79 (*Musca*). England. T A Unknown. Psilidae.
- longervo* Harris, 1780: 74 (*Musca*). Incosp. *longerro* Harris. Psilidae. Name status by present revision.
- longicornis* Fabricius, 1775: 783 (*Musca*). America. LT ♂ UZMC F. Dolichopodidae, *Condylostylus longicornis* Fabricius. Lectotype designated by Bickel *in press*.
- longicornis* Fabricius, 1781: 420 (*Rhagio*). Germany. T A UZMC F. Keroplatidae, *Macrocerata macrocerata* Gmelin. **N. Status**
- longicornis* Geoffroy, 1785: 477 (*Musca*). France, Paris. T A Author. Tephritidae. Preocc. Fabricius 1775; synonym of *Musca lincensis* Schrank.
- longicornis* de Villers, 1789a: 538 (*Musca*). N. n. *Musca lincensis* Schrank. Tephritidae, *Genus? lincensis* Schrank. Preocc. Fabricius 1775.
- longipennis* Fabricius, 1794: 323 (*Musca*). Germany. T A UZMC F. Lauxaniidae, *Minettia longipennis* Fabricius.
- longipes* Scopoli, 1763: 336 (*Musca*). Carniola. T A DESTROY.
- longipes* Geoffroy, 1785: 497 (*Musca*). France, Paris. T A Author. Preocc. Scopoli 1763.
- longipes* Fabricius, 1787: 348 (*Musca*). Cayenne. T A UZMC F. Micropezidae, *Cardiacephala guttata* Walker. Preocc. Scopoli 1763.
- longipes* Thunberg, 1791: 127 (*Musca*). N. n. *Musca femorata* Linnaeus. Syrphidae, *Chalcosyrphus femoratus* Linnaeus. Preocc. Scopoli 1763.
- longipes* Fabricius, 1805: 298 (*Musca*). South America. HT ♂ UZMC. Tachinidae, *Cholomyia acromion* Wiedemann. Preocc. Scopoli 1763. See Townsend 1931a: 93.
- longipes* Presl, 1822: 206 (*Musca*). Ligurian Prussian Amber. T A Unknown. Preocc. Scopoli 1763.
- longipes* Stephens, 1829b: 302 (*Musca*). England. Tachinidae. Published in synonymy, not subsequently validated by usage. Published in synonymy with *Dexia rustica* Fabricius.
- longipes* Paterson, 1957: 106, fig. 2 (*Musca*). Botswana, Martin's Drift. HT ♂ SAIMR. Muscidae, *Musca hugonis* Pont. Preocc. Scopoli 1763.

- longisco Harris, 1780: 83, pl. 24, fig. 30 (*Musca*). England. T A Unknown. Syrphidae, *Xylota sylvarum* Linnaeus. Name improperly formed (verb).
- longula** Fallén, 1824: 72 (*Musca*). Sweden, Skåne. LT ♂ NRS. Anthomyiidae, *Subhylemyia longula* Fallén. Lectotype designated by Michelsen 1983: 114.
- lothari** Zielke, 1974: 9 (*Musca*). Kenya, 26 Miles SW Nairobi. HT ♂ CAS. Muscidae, *Musca xanthomelaena* Wiedemann.
- lucens** Villeneuve, 1922: 336 (*Pristirynchomyia*). Ceylon, Kandy. ST ♂ UNKNOWN. Muscidae, *Musca conducens* Walker.
- lucida** Scopoli, 1763: 347 (*Musca*). Carniola. T A DESTROY. Syrphidae, *Chrysogaster lucida* Scopoli.
- lucidula** Loew, 1856: 48 (*Cyrtoneura*). Egypt, El Tûr. T ♀ NMW. Muscidae, *Musca lucidula* Loew.
- lucifer** Gmelin, 1790: 2832 (*Musca*). Subsequent Usage of *Bibio lucifer* Fabricius. Bombyliidae, *Poecilanthrax lucifer* Fabricius.
- lucophaeus** Harris, 1780: 74, pl. 21, fig. 3 (*Musca*). England. T A Unknown. Scathophagidae, *Scathophaga scybalaria* Linnaeus.
- lucopheus** Harris, 1780: 74, pl. 21, fig. 3 (*Musca*). Incosp. *lucophaeus* Harris. Scathophagidae, *Scathophaga scybalaria* Linnaeus. Name status by present revision.
- lucorum** Linnaeus, 1758a: 592 (*Musca*). Sweden. LT ♂ LSL. Syrphidae, *Leucozona lucorum* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 157.
- lucorum** Houttuyn, 1768: 488 (*Musca*). Subsequent Usage of *Musca lucorum* Linnaeus. Syrphidae, *Leucozona lucorum* Linnaeus.
- lucorum** Fallén, 1823: 55 (*Musca*). Sweden. LT ♂ NRS. Muscidae, *Helina erecta* Harris. Preocc. Linnaeus 1758. Lectotype designated by Pont 1984: 288.
- lucozona** Sherborn, 1922: 3688 (*Musca*). Missp. *leucozona* Panzer. Tachinidae, *Thelaira leucozona* Panzer.
- luctuosa** Schrank, 1803: 122 (*Musca*). Germany, Bavaria, Ingolstadt. T A Unknown.
- ludeus** Harris, 1780: 157, pl. 47, fig. 3 (*Musca*). England. T A Unknown. Dolichopodidae, *Argyra diaphana* Fabricius.
- ludicrous** Harris, 1780: index (*Musca*). Incosp. *ludicus* Harris. Dolichopodidae, *Poecilobothrus nobilitatus* Linnaeus. Name status by present revision.
- ludicus** Harris, 1780: 157, pl. 47, fig. 2 (*Musca*). England. T A Unknown. Dolichopodidae, *Poecilobothrus nobilitatus* Linnaeus.
- ludifacies** Robineau-Desvoidy, 1863: 621 (*Musca*). France, Paris. T A DESTROY. Muscidae, *Musca autumnalis* De Geer.
- ludifica** Fabricius, 1794: 323 (*Musca*). Germany, Kiel. LT ♂ UZMC F. Muscidae, *Musca autumnalis* De Geer. Lectotype designated by Michelsen 1979a: 186.
- lugens** Fabricius, 1787: 350 (*Musca*). Sierra Leone. T A UZMC F. Platystomatidae, *Engistoneura lugens* Fabricius.
- lugubris** Fabricius, 1775: 770 (*Syrphus*). Denmark. T A UZMC F. Syrphidae, *Pipiza lugubris* Fabricius.
- lugubris** Gmelin, 1790: 2832 (*Musca*). Subsequent Usage of *Bibio lugubris* Fabricius. Therevidae, *Thereva lugubris* Fabricius.
- lunata** Turton, 1801: 645 (*Musca*). Subsequent Usage of *Syrphus lunatus* Fabricius. Syrphidae, *Eumerus lunatus* Fabricius.

- lunata** Fabricius, 1805: 292 (*Musca*). Madeira. LT ♂ UZMC. Calliphoridae, *Stomorrhina lunata* Fabricius. See Townsend 1931c: 372. Lectotype designated by Rognes 1991a: 245.
- lunula** de Villers, 1789a: 433, pl. 9, fig. 16 (*Musca*). Europe. T A Author. Syrphidae, *Eristalis pertinax* Scopoli.
- lupina** Swederus, 1787: 289 (*Musca*). New Zealand. T A BMNH B. Tachinidae, *Bothrophora lupina* Swederus.
- lupulina** Fabricius, 1787: 344 (*Musca*). Denmark, Zealand. T A UZMC. Lauxaniidae, *Minettia lupulina* Fabricius.
- lurida** Fabricius, 1781: 441 (*Musca*). Germany. T A UZMC F. Tachinidae, *Servillia lurida* Fabricius.
- lusoria** Wiedemann, 1824: 47 (*Musca*). South Africa, Cape of Good Hope. T ♂ UZMC. Muscidae, *Musca lusoria* Wiedemann.
- lustrator** Harris, 1780: 148, pl. 43, fig. 72 (*Musca*). England. T A Unknown. Fanniidae, *Fannia lustrator* Harris.
- lutaria** Fabricius, 1794: 344 (*Musca*). Germany, Kiel. T A UZMC F. Scathophagidae, *Scathophaga lutaria* Fabricius.
- lutea** Gmelin, 1790: 2843 (*Musca*). Europe. T A NMI. Preocc. Harris 1780.
- luteipennis** Zetterstedt, 1845: 1325 (*Musca*). *Nomen nudum*. Muscidae, *Pyrellia rapax* Harris. Published in synonymy, not subsequently validated by usage. Proposed in synonymy with *aenea* Zetterstedt.
- luteola** Scopoli, 1763: 349 (*Musca*). Carniola. T A DESTROY. Chyromyidae, *Chyromya flava* Linnaeus.
- luteola** Gmelin, 1790: 2879 (*Musca*). Europe. T A NMI. Syrphidae, *Myolepta dubia* Fabricius. Preocc. Scopoli 1763.
- luteola** Fabricius, 1805: 286 (*Musca*). Guinea. T A UZMC. Calliphoridae, *Auchmeromyia senegalensis* Macquart. Preocc. Scopoli 1763.
- luteus** Harris, 1780: 116, pl. 34, fig. 16 (*Musca*). England. T A Unknown. Psilidae.
- lychnidis** Fabricius, 1787: 353 (*Musca*). Germany, Kiel. T A UZMC F. Tephritidae, *Myoleja caesio* Harris.
- lyra** Harris, 1776: 42, pl. 10, fig. 5 (*Musca*). England. T A Unknown. Syrphidae, *Eristalis lyra* Harris.
- lycea** Walker, 1849: 873 (*Musca*). Uruguay, Montevideo. LT ♂ BMNH. Calliphoridae, *Compsomyiops fulvicrura* Robineau-Desvoidy. Lectotype designated by Dear 1985b: 148.
- macellaria** Fabricius, 1775: 776 (*Musca*). "America". HT ♀ UZMC F. Calliphoridae, *Cochliomyia macellaria* Fabricius. See Townsend 1931a: 70; Dear 1985b: 140.
- macellaria** Haliday, 1833: 165 (*Musca*). Ireland, Down, Holywood. T A NMI. Calliphoridae, *Pollenia* sp. Preocc. Fabricius 1775. N. Comb.
- macrocephala** Harris, 1780: 82 (*Musca*). Subsequent Usage of *Conops macrocephala* Linnaeus. T A Unknown. Syrphidae, *Rhingia campestris* Meigen. Misidentification.
- macrocephala** Rossi, 1794: 72 (*Musca*). Italy. T A ZMHU. Preocc. Harris 1780.
- macroceras** Gmelin, 1790: 2866 (*Musca*). N. n. *Rhagio longicornis* Fabricius. Keroplatidae, *Macrocerata macroceras* Gmelin. N. Comb.

- macrophthalma*** Preyssler, 1791: 68, pl. 2, fig. 6 (*Musca*). Czechoslovakia, Bohemia. T A DESTROY. Syrphidae, *Eristalinus aeneus* Scopoli. See Rozkosny, Chvála & Pont 1982: 352.
- mactans*** Fabricius, 1787: 344 (*Musca*). Cayenne. T A UZMC F.
- macularis*** Walker, 1858b: 104 (*Musca*). Moluccas, Aru Islands. LT ♂ BMNH. Calliphoridae, *Paramenia macularis* Walker. Lectotype designated by Crosskey 1965: 131 (HT designation Townsend 1931c: 375).
- maculata*** Scopoli, 1763: 326 (*Musca*). Carniola. T A DESTROY. Muscidae, *Graphomya maculata* Scopoli.
- maculata*** Harris, 1780: 140, pl. 41, fig. 37 (*Musca*). England. T ♀ Unknown. Muscidae, *Graphomya maculata* Scopoli. Preocc. Scopoli 1763.
- maculata*** de Villers, 1789a: 418 (*Musca*). Subsequent Usage of *Nemotelus maculatus* De Geer. Rhagionidae, *Rhagio maculatus* De Geer.
- maculata*** de Villers, 1789a: 433 (*Musca*). Subsequent Usage of *Volucella maculata* Geoffroy.
- maculata*** Meigen, 1804: 294 (*Atherix*). Germany. T ♂ MNHNP. Athericidae, *Atherix ibis* Fabricius.
- maculatus*** Fabricius, 1781: 420 (*Rhagio*). Germany. T A UZMC F.
- major larvarum*** De Geer, 1776: 24 (*Musca*). *Nomen nudum*. N. n. *Musca larvarum* Linnaeus. Tachinidae, *Exorista larvarum* Linnaeus. Not available; polynominal, not binomial.
- malaisei*** Emden, 1965: 91, fig. 2 (*Musca*). Burma, Kambaiti. HT ♂ NRS. Muscidae, *Musca malaisei* Emden.
- malansi*** Stein, 1857: 106 (*Musca*). *Nomen nudum*. Switzerland, Graubünden. Calliphoridae.
- mallochi*** Thompson & Pont, 1992: (*Musca*). N. n. *Musca squamata* Malloch. Muscidae, *Musca mallochi* Thompson & Pont. N. Name
- manicata*** Fabricius, 1794: 334 (*Musca*). France. HT ♂ MNHNP. Ephyrinae, *Ochthera manicata* Fabricius. See Claussen 1977: 511.
- mano*** Harris, 1780: 142, pl. 42, fig. 45 (*Musca*). England. T ♂ Unknown. Muscidae, *Eudasypheora cyanella* Meigen. Name improperly formed (verb).
- mantes*** Harris, 1780: 148, pl. 43, fig. 73 (*Musca*). England. T A Unknown. Muscidae, *Drymeia vicana* Harris.
- mantis*** De Geer, 1776: 143, pl. 8, fig. 17 (*Musca*). Sweden, Stockholm. NT ♂ NRS. Ephyrinae, *Ochthera mantis* De Geer. Neotype designation (Claussen 1977: 514).
- mantos*** Harris, 1780: 148 (*Musca*). Incosp. *mantes* Harris. Muscidae, *Drymeia vicana* Harris. Name status by present revision.
- marginalis*** Geoffroy, 1785: 497 (*Musca*). France, Paris. T A Author. Suppressed by I.C.Z.N. 1988: 236.
- marginalis*** Fallén, 1824: 66 (*Musca*). Sweden, Västergötland, Kinneskogen. LT ♂ NRS. Muscidae, *Spilogona marginifera* Hennig. Preocc. Geoffroy 1785. Suppressed by I.C.Z.N. 1988: 236. Lectotype designated by Pont 1984: 289.
- marginalis*** Wiedemann, 1830: 395 (*Musca*). South Africa, Nubia, Arabia. ST A UZMC NMW. Calliphoridae, *Chrysomya marginalis* Wiedemann. Conserved by I.C.Z.N. 1988: 236.
- marginata*** Fabricius, 1775: 761 (*Stratiomys*). England. T A BMNH B. Stratiomyidae, *Nemotelus pantherinus* Linnaeus.

- marginata** Fabricius, 1775: 784 (*Musca*). Germany, Saxony. T A UZMCF. Sciomyzidae, *Coremacera marginata* Fabricius.
- marginella** Gmelin, 1790: 2836 (*Musca*). N. n. *Stratiomys marginata* Fabricius. Stratiomyidae, *Nemotelus pantherinus* Linnaeus.
- marginella** Estlund, 1796: 131 (*Musca*). Sweden. T A UNKNOWN. Preocc. Gmelin 1790.
- marginifera** Walker, 1858b: 105 (*Musca*). Aru Islands. T ♀ BMNH. Calliphoridae, *Lucilia papuensis* Macquart.
- maritima** Scopoli, 1763: 344 (*Musca*). Yugoslavia, Tergestum. T A DESTROY. Syrphidae, *Xylota segnis* Linnaeus.
- marmorata** Geoffroy, 1785: 474 (*Musca*). France, Paris. T A Author.
- marmorata** Fabricius, 1805: 300 (*Musca*). Germany. T A MNHNP. Tachinidae, *Lespesia vertiginosa* Fallén. Preocc. Geoffroy 1785. N. Comb.
- maronea** Walker, 1849: 886 (*Musca*). Australia. HT ♀ BMNH. Muscidae, *Neomyia australis* Macquart. See Pont 1973: 203.
- marsya** Walker, 1849: 890 (*Musca*). Cape. HT ♂ BMNH. Muscidae, *Pyrellina marsya* Walker. See Pont 1973: 287.
- martia** Scopoli, 1763: 327 (*Musca*). Carniola. T A DESTROY. Sarcophagidae.
- maura** Linnaeus, 1758a: 590 (*Musca*). Europe. T A LSL. Bombyliidae, *Hemipenthes maurus* Linnaeus. 1 sp. in LSL (Dear MS).
- maura** Turton, 1801: 634 (*Musca*). N. n. *Rhagio morio* Fabricius. Sciaridae, *Bradysia forcipata* Fabricius. Preocc. Linnaeus 1758.
- maura** Fabricius, 1805: 302 (*Musca*). Barbaria [= Morocco: Tangiers]. LT ♂ UZMCF. Sarcophagidae, *Nyctia halterata* Panzer. Preocc. Linnaeus 1758. Lectotype designated by Pape 1986: 302 (also Townsend 1931c: 382).
- mazaca** Walker, 1849: 889 (*Musca*). South Africa, Cape of Good Hope. T ♀ BMNH. Muscidae, *Neomyia peronii* Robineau-Desvoidy.
- mediana** Wiedemann, 1830: 657 (*Musca*). China, Macao. LT ♂ NMW. Muscidae, *Musca sorbens* Wiedemann. Lectotype designated by Pont 1973: 288.
- meditabunda** Fabricius, 1781: 444 (*Musca*). Italy & Denmark, Copenhagen. LT ♀ UZMCF. Muscidae, *Myospila meditabunda* Fabricius. Lectotype designated by Michelsen 1979a: 186.
- meditalnenda** Gmelin, 1790: 2845 (*Musca*). N. n. *Musca meditabunda* Fabricius. Muscidae, *Myospila meditabunda* Fabricius.
- meditata** Fallén, 1825: 87 (*Musca*). Sweden. LT ♂ NRS. Muscidae, *Macrorchis meditata* Fallén. Lectotype designated by Pont 1984: 289.
- megacephala** Fabricius, 1794: 317 (*Musca*). Guinea. T A UZMCF. Calliphoridae, *Chrysomya megacephala* Fabricius.
- melancholica** Turton, 1801: 648 (*Musca*). Subsequent Usage of *Syrphus melanocholica* Fabricius. Syrphidae, *Merodon natans* Fabricius.
- melanius** Harris, 1776: 59, pl. 15, fig. 12 (*Musca*). England. T A Unknown. Syrphidae, *Eristalinus sepulchralis* Linnaeus.
- melanochrysa** Gmelin, 1790: 2862 (*Musca*). Europe. T A NMI.
- melanochrysa** Gmelin, 1790: 2879 (*Musca*). Europe. T A NMI. Syrphidae, *Xylota segnis* Linnaeus. Preocc. Gmelin 1790: 2862.
- melanopis** Gmelin, 1790: 2836 (*Musca*). Europe. T A NMI. Stratiomyidae.
- melanopis** Turton, 1801: 631 (*Musca*). Subsequent Usage of *Musca melanopis* Gmelin. Stratiomyidae.

- melanoptera** Gmelin, 1790: 2833 (*Musca*). Non-European. T A NMI. Bombyliidae.
- melanoptera** Fallén, 1817: 253 (*Musca*). Sweden, Östergötland & Västergötland. T A NRS. Calliphoridae, *Morinia doronici* Scopoli. Preocc. Gmelin 1790. N. Syn.
- As this name is preoccupied and must be replaced, we prefer to use the oldest probable name for the taxon (Schiner 1856: 412). Specialists should select a neotype for *Morinia doronici* (Scopoli) to fix this interpretation.
- melanopyrrha** Forster, 1770: 15 (*Musca*). *Nomen nudum*. Britain. Syrphidae, *Volucella bombylans* Linnaeus. Published without diagnosis or indication.
- melanopyrrha** Forster, 1771: 98 (*Musca*). England & Germany. T A LSL. Syrphidae, *Volucella bombylans* Linnaeus. See Day & Fitton 1977.
- melanura** Schrank, 1781: 457 (*Musca*). Austria, Linz. T A Unknown.
- melanura** Turton, 1801: 603 (*Musca*). Subsequent Usage of *Musca melanura* Schrank.
- melinus** Harris, 1780: 74 (*Musca*). Incosp. *mellinus* Harris. Dryomyzidae. Name status by present revision.
- mellina** Linnaeus, 1758a: 594 (*Musca*). Sweden. ST ♀ LSL. Syrphidae, *Melanostoma mellinum* Linnaeus. See Thompson *et al.* 1982: 157.
- mellina** Harris, 1780: 80, pl. 24, fig. 22 (*Musca*). England. T A Unknown. Syrphidae, *Scaeva pyrastri* Linnaeus. Preocc. Linnaeus 1758.
- mellinus** Harris, 1780: 74, pl. 21, fig. 4 (*Musca*). England. T A Unknown. Dryomyzidae. Preocc. Linnaeus 1758.
- menthastri** Linnaeus, 1758a: 594 (*Musca*). Sweden. ST ♀ LSL. Syrphidae, *Sphaerophoria menthastri* Linnaeus. See Thompson *et al.* 1982: 157.
- merdaria** Fabricius, 1794: 344 (*Musca*). Germany, Kiel. T A UZMC F. Scathophagidae, *Scathophaga stercoraria* Linnaeus.
- meridiana** Linnaeus, 1758a: 595 (*Musca*). Europe. HT ♂ LSL. Muscidae, *Mesembrina meridiana* Linnaeus. See Pont 1981: 170.
- meridiana** Bloch, 1776: 184, pl. 5, fig. 24 (*Musca*). Subsequent Usage of *Musca meridiana* Linnaeus. Muscidae, *Mesembrina meridiana* Linnaeus.
- meruensis** Zielke, 1973: 368 (*Musca*). Tanzania, Mt. Meru. HT ♂ CNC. Muscidae, *Musca meruensis* Zielke.
- merulina** Scopoli, 1763: 344 (*Musca*). Carniola. T A DESTROY. Syrphidae, *Cheilosia* sp.
- mesembrinoides** Walker, 1861b: 244 (*Musca*). New Guinea, Dorey. T ♀ BMNH. Calliphoridae, *Onesia obscurata* Walker.
- mesoleuca** Lichtenstein, 1796: 216 (*Musca*). Sinai. T A Auction.
- mesopotamiensis** Patton, 1920: 770, pl. 69 (*Musca*). Iraq, Euphrates & Tigris Rivers. ST A UNKNOWN. Muscidae, *Musca larvipara* Portschinsky. See Pont 1990: 167.
- metallina** Gmelin, 1790: 2874 (*Musca*). Subsequent Usage of *Syrphus metallinus* Fabricius. Syrphidae, *Lejogaster metallina* Fabricius.
- metallina** Gravenhorst, 1807: 378 (*Musca*). Unknown, probably Germany. T A Lost. Tachinidae.
- meteorica** Linnaeus, 1758a: 597 (*Musca*). Europe. T A Lost. Muscidae, *Hydrotaea meteorica* Linnaeus. See Pont 1981: 171.
- meteorica** Schrank, 1781: 464 (*Musca*). Subsequent usage of *Musca meteorica* Linnaeus. Muscidae, *Hydrotaea meteorica* Linnaeus.
- meticulosa** Scopoli, 1763: 344 (*Musca*). Carniola. T A DESTROY. Syrphidae, *Neoascia meticulosa* Scopoli. See Thompson 1981b: 475.

- metilia* Walker, 1849: 898 (*Musca*). Nepal. T ♂ BMNH. Calliphoridae, *Lucilia papuensis* Macquart.
- metus* Harris, 1780: 114, pl. 33, fig. 27 (*Musca*). England. T A Unknown. Tachinidae. *micans* Turton, 1801: 637 (*Musca*). Subsequent Usage of *Syrphus micans* Fabricius. Syrphidae, *Volucella inanis* Linnaeus.
- micans* Fabricius, 1805: 291 (*Musca*). India Orientali. HT A UZMC. Calliphoridae, *Catapicephala micans* Fabricius. See Townsend 1931c: 375.
- micans* Meigen, 1838: 303 (*Musca*). N. n. *Calliphora rufipalpis* Macquart. Calliphoridae, *Onesia floralis* Robineau-Desvoidy. Preocc. Fabricius 1805.
- microleon* Linnaeus, 1758a: 589 (*Musca*). Europe. T A LSL. Stratiomyidae, *Odonomyia microleon* Linnaeus. 2 sp. in LSL (Dear MS).
- miliaria* Schrank, 1781: 476 (*Musca*). Austria, Vienna, Prater. T A Unknown. Tephritidae, *Xyphosia miliaria* Schrank.
- minima* Zetterstedt, 1837: 42 (*Musca*). *Nomen nudum*. [Sweden, Marklin]. T A ZIL. Calliphoridae, *Melanomya nana* Meigen.
- minima* Zetterstedt, 1838: 654 (*Musca*). Sweden, Marklin. LT ♂ ZIL. Calliphoridae, *Melanomya nana* Meigen. Preocc. Harris 1780. Lectotype designated by Rognes 1991a: 199.
- minima* Rondani, 1868: 52 (*Musca*). Italy, Parma. T ♀ USB. Muscidae, *Musca tempestiva* Fallén. Preocc. Harris 1780.
- minimus* Harris, 1780: 87 (*Musca*, as *vomitor minimus*). England. T A Unknown. Calliphoridae, *Calliphora vomitoria* Linnaeus.
- minimus* Harris, 1780: index, pl. 35, fig. 43 (*Musca*, as *vibrans minimus*). England. T A UNKNOWN.
- minor* Harris, 1780: 153, pl. 45, fig. 83 (*Musca*, as *socio*). England. T A Unknown. Fanniidae, *Fannia canicularis* Linnaeus.
- minor* Turton, 1801: 655 (*Musca*). N. n. *Musca pusillus* Fabricius. Stratiomyidae, *Oplodontha minuta* Fabricius. Preocc. Harris 1780.
- minor* Macquart, 1851: 226 (*Musca*). Tasmania. LT ♂ MNHNP. Muscidae, *Musca domestica* Linnaeus. Preocc. Harris 1780. Lectotype designated by Pont 1967: 187.
- minor domestica* De Geer, 1776: 26 (*Musca*). *Nomen nudum*. [Sweden]. T A NRS. Fanniidae, *Fannia scalaris* Fabricius. Not available; polynominal, not binomial.
- minor larvarum* De Geer, 1776: 25 (*Musca*). *Nomen nudum*. [Sweden]. T A NRS. Tachinidae. Not available; polynominal, not binomial.
- minuta* Fabricius, 1787: 353 (*Musca*). Germany, Kiel. T A UZMC F. Chloropidae, *Meromyza saltatrix* Linnaeus. Preocc. Harris 1780.
- minuta* Fabricius, 1794: 268 (*Stratiomys*). India, Tranquebar. T A UZMC F. Stratiomyidae, *Oplodontha minuta* Fabricius. N. Comb.
- minuta* Awati, 1916b: 138 (*Musca*). India, Delhi. ST ♀ CRIK. Muscidae, *Musca confiscata* Speiser. Preocc. Harris 1780.
- minutior* Turton, 1801: 631 (*Musca*). N. n. *Stratiomyia minuta* Fabricius. Stratiomyidae, *Oplodontha minuta* Fabricius.
- minutissima* de Razoumowsky, 1789: 230 (*Musca*). Switzerland, Jorat. T A UNKNOWN.
- minutus* Harris, 1780: 74, pl. 21, fig. 7 (*Musca*). England. T A Unknown. Pallopteridae.
- mirabilis* Guérin-Méneville, 1838: 296 (*Musca*). Subsequent Usage of *Rutilia mirabilis* Guérin-Méneville. Tachinidae, *Formosia mirabilis* Guérin-Méneville.

- modesta* de Meijere, 1904: 106 (*Musca*). Java, Pasuruan. LT ♂ ITZA. Muscidae, *Musca crassirostris* Stein. Lectotype designated by Pont 1970a: 94.
- moerens* Fabricius, 1794: 349 (*Musca*). Guinea. T A UZMC F. Platystomatidae, *Engistoneura moerens* Fabricius.
- moesta* Gmelin, 1790: 2874 (*Musca*). N. n. *Syrphus lugubris* Fabricius. Syrphidae, *Pipiza lugubris* Fabricius.
- moesta* Gravenhorst, 1807: 379 (*Musca*). Unknown, probably Germany. T A UNKNOWN. Tachinidae. Preocc. Gmelin 1790.
- molio* Harris, 1780: 111, pl. 33, fig. 54 (*Musca*). England. T A Unknown. Syrphidae, *Neoascia podagraria* Fabricius. Name improperly formed (verb). See Thompson 1981b: 475.
- molitus* Harris, 1780: 110 (*Musca*). England. T A Unknown. Syrphidae, *Sphaerophoria scripta* Linnaeus.
- mollicula* Fallén, 1825: 90 (*Musca*). Sweden, Skåne. LT ♂ NRS. Muscidae, *Coenosia mollicula* Fallén. Lectotype designated by Pont 1984: 289.
- mollis* Walker, 1849: 892 (*Musca*). Ontario, St. Martin's Falls. T ♀ BMNH. Calliphoridae, *Phormia regina* Meigen.
- monachae* Ratzeburg, 1844: 173 (*Musca*). Germany. T A Unknown. Tachinidae, *Exorista larvarum* Linnaeus.
- moneta* Walker, 1849: 904 (*Musca*). France? Lapland? T ♀ BMNH. Sarcophagidae, *Ravinia moneta* Walker.
- monoculus* Swederus, 1787: 287 (*Musca*). North America. T A Unknown. Syrphidae.
- morio* Linnaeus, 1758a: 590 (*Musca*). Europe. T A LSL. Bombyliidae, *Hemipenthes morio* Linnaeus. 1 sp. in LSL (Dear MS).
- morio* Fabricius, 1794: 275 (*Rhago*). Germany. T A UZMC F. Sciaridae, *Bradysia forcipata* Fabricius.
- mortisequa* Kirby, 1837: 316 (*Musca*). Alaska ["North America, lat. 65°"]. T A Unknown. Calliphoridae, *Cynomya cadaverina* Robineau-Desvoidy.
- mortuorum* Linnaeus, 1761: 452 (*Musca*). Sweden. T A LSL. Calliphoridae, *Cynomya mortuorum* Linnaeus. 1 sp. in LSL (Dear MS).
- muliebris* Harris, 1780: 75, pl. 21, fig. 9 (*Musca*). England. T A Unknown. Palloppteridae, *Palloptera muliebris* Harris.
- mulinus* Harris, 1780: index (*Musca*). Incosp. *mellinus* Harris (1780: 74). Dryomyzidae. Name status by present revision.
- mulsus* Harris, 1780: 116, pl. 34, fig. 20 (*Musca*). England. T A Unknown. Lauxaniidae.
- multispina* Awati, 1916b: 138 (*Musca*). India. ST ♀ CRIK. Muscidae, *Musca domestica* Linnaeus.
- munroa* Wiedemann, 1830: 398 (*Musca*). Java. T ♀ UZMC. Calliphoridae, *Isomyia viredaurea* Wiedemann.
- munroi* Patton, 1936: 487, fig. 11 (*Musca*). South Africa, Natal, Prospect. ST ♂ UNKNOWN. Muscidae, *Musca munroi* Patton. See Pont 1990: 167.
- muralis* Walker, 1849: 888 (*Musca*). Ontario, St. Martin's Falls. HT ♀ BMNH. Calliphoridae, *Lucilia illustris* Meigen. [teste Pont].
- muraria* Scopoli, 1763: 338 (*Musca*). Carniola. T A DESTROY.
- murina* Schrank, 1776: 95 (*Musca*). Austria, Linz. T A Unknown. Syrphidae.
- murorum* Turton, 1801: 641 (*Musca*). Subsequent Usage of *Syrphus murorum* Fabricius. Syrphidae, *Eristalis murorum* Fabricius.

- muscaria* Turton, 1801: 631 (*Musca*). Subsequent Usage of *Stratiomys muscaria* Fabricius. Stratiomyidae, *Oxycera muscaria* Fabricius.
- mussitanus* Gmelin, 1790: 2867 (*Musca*). Subsequent Usage of *Syrrhus mussitanus* Fabricius. Syrphidae, *Arctophila superbii* Müller.
- mutabilis** Linnaeus, 1758a: 592 (*Musca*). Sweden. LT ♀ LSL. Syrphidae, *Microdon mutabilis* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 157.
- mutabilis** Fabricius, 1787: 331 (*Stratiomys*). Cayenne. T A UZMC F. Stratiomyidae, *Hoplitomyia mutabilis* Fabricius.
- mutata** Fallén, 1825: 92 (*Musca*). Sweden. LT ♂ NRS. Muscidae, *Lophosceles mutatus* Fallén. Lectotype designated by Pont 1984: 290.
- myennis** Coquebert, 1798: (*Musca*). France. T A MNHNP. Otitidae.
- myopina** Fallén, 1824: 65 (*Musca*). Sweden, Skåne, Lund. LT ♂ NRS. Anthomyiidae, *Myopina myopina* Fallén. Lectotype designated by Michelsen 1983: 114.
- mystacea** Linnaeus, 1758a: 591 (*Musca*). Europe. HT ♀ LSL. Muscidae, *Mesembrina mystacea* Linnaeus. See Pont 1981: 172.
- naevia** Gmelin, 1790: 2879 (*Musca*). Europe. T A NMI. Syrphidae, *Milesia semiluctifera* de Villers.
- naevosus** Harris, 1780: 75, pl. 21, fig. 10 (*Musca*). England. T A Unknown. Platystomatidae.
- nana** Stephens, 1829a: 60 (*Musca*). *Nomen nudum*. England. T A BMNH. Calliphoridae.
- nana** Meigen, 1830: 375 (*Musca*). Germany, Berlin District. HT ♀ Lost. Muscidae, *Musca tempestiva* Fallén. See Pont 1986b: 232.
- nana** Meigen, 1838: 306 (*Musca*). Subsequent Usage of *Nitellia nana* Robineau-Desvoidy. Calliphoridae, *Pollenia rufa* Fabricius.
- nancauriensis** Schiner, 1868: 306 (*Musca*). *Nomen nudum*. Nicobar Is. T ♂ NMW. Muscidae, *Musca domestica* Linnaeus. Published in synonymy, not subsequently validated by usage. Proposed in synonymy with *domestica* Linnaeus.
- napobrassicae** Bjerkanter, 1780: 196 (*Musca*). Sweden. T A UNKNOWN. Anthomyiidae, *Delia radicum* Linnaeus.
- nasatus** Harris, 1780: 105, pl. 32, fig. 35 (*Musca*). England. T A Unknown. Syrphidae, *Rhingia* sp.
- nasuta** Schrank, 1781: 475 (*Musca*). Austria, Vienna, Brigitteau. T A Unknown. Chloropidae, *Chlorops pumilionis* Bjerkanter.
- natalensis** Villeneuve, 1916: 512 (*Musca*). South Africa, Natal, Durban. ST ♂ SAMCT. Muscidae, *Musca gabonensis* Macquart.
- natans** Turton, 1801: 639 (*Musca*). Subsequent Usage of *Syrrhus natans* Fabricius. Syrphidae, *Merodon natans* Fabricius.
- navus** Harris, 1780: 109, pl. 33, fig. 50 (*Musca*). England. T A Unknown. Syrphidae, *Platycheirus* sp.
- nebulo** Fabricius, 1794: 321 (*Musca*). India Orientali. T A DESTROY. Muscidae, *Musca domestica* Linnaeus. See Michelsen 1979a: 187.
- nebulosa** Panzer, 1798b: 20 (*Musca*). Germany, Nuremberg. T ♂ ZMHU. Tachinidae, *Allophora obesa* Fabricius.
- nebulosa** Lamarck, 1816: 363 (*Musca*). Subsequent Usage of *Musca nebulosa* Panzer. Tachinidae, *Allophora obesa* Fabricius.
- nectarea** Linnaeus, 1767: 983 (*Musca*). Europe. T A LSL. Scatopsidae, *Scatopse notata* Linnaeus. 1 sp. in LSL (Dear MS).

- nectareus* Fabricius, 1787: 341 (*Syrphus*). Denmark. T A UZMC. Syrphidae, *Episyphus cannabina* Scopoli.
- nectarina* Gmelin, 1790: 2876 (*Musca*). N. n. *Musca nectareus* Fabricius. Syrphidae, *Episyphus cannabina* Scopoli.
- nectarivora* Turton, 1801: 656 (*Musca*). N. n. *Musca nectarea* Linnaeus. Scatopsidae, *Scatopse notata* Linnaeus.
- negriabdomina* Awati, 1916b: 138 (*Musca*). India. HT A CRIK. Muscidae, *Musca negriabdomina* Awati.
- nemorum* Linnaeus, 1758a: 591 (*Musca*). Sweden. LT ♀ LSL. Syrphidae, *Eristalis arbustorum* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 158.
- nemorum* Fallén, 1823: 55 (*Musca*). Sweden, Skåne. LT ♂ NRS. Muscidae, *Phaonia incana* Wiedemann. Preocc. Linnaeus 1758. Lectotype designated by Pont 1984: 290.
- neustriæ* Ratzeburg, 1844: 173 (*Musca*). Germany. T A Unknown. Tachinidae, *Tachina neustriæ* Ratzeburg.
- nevilli* Kleynhans, 1987: 115, fig. 1 (*Musca*). South Africa, N Transvaal, 15 km N of Pretoria, Farm ‘‘Kaalplaas’’. HT ♂ NCIP. Muscidae, *Musca nevilli* Kleynhans.
- nigra* Linnaeus, 1764a: 285 (*Musca*). *Nomen nudum*. [Sweden]. Chloropidae, *Oscinella frit* Linnaeus. Published in non-binominal work. N. Syn.
- nigra* Forskål, 1775: 24 (*Musca*). Arabia. T A Unknown. Syrphidae, *Ischiodon* sp.
- nigra* de Villers, 1789a: 432 (*Musca*). Subsequent Usage of *Volucella nigra* Geoffroy.
- nigra* de Villers, 1789a: 420 (*Musca*). Subsequent Usage of *Nemotelus niger* De Geer. Scenopinidae, *Scenopinus niger* De Geer.
- nigrana* Turton, 1801: 643 (*Musca*). N. n. *Syrphus nigrita* Fabricius. Syrphidae, *Sterphus jamaicensis* Gmelin.
- nigricans* de Villers, 1789a: 509 (*Musca*). Sweden? T A Unknown. Anisopodidae, *Sylvicola fuscatus* Fabricius. N. Syn.
- nigricans* Gmelin, 1790: 2836 (*Musca*). Europe. T A NMI. Stratiomyidae, *Odonomyia microleon* Linnaeus. Preocc. de Villers 1789.
- nigricans* Robineau-Desvoidy, 1830: 394 (*Sphora*). Australia. T A DESTROY. Muscidae, *Musca domestica* Linnaeus.
- nigricans* Macquart, 1835: 268 (*Musca*). Subsequent Usage of *Sphora nigricans* Robineau-Desvoidy. Muscidae, *Musca domestica* Linnaeus.
- nigriceps* Walker, 1849: 881 (*Musca*). Subsequent Usage of *Lucilia nigriceps* Macquart. Calliphoridae, *Lucilia nigriceps* Macquart.
- nigricolor* Fallén, 1825: 81 (*Musca*). Sweden, Skåne. HT ♂ ZIL. Muscidae, *Hebecnema nigricolor* Fallén. See Pont 1984: 290.
- nigricornis* Fabricius, 1794: 331 (*Musca*). India Orientali. T A UNKNOWN. Tachinidae.
- nigricruris* Rainbow, 1897: 98 (*Ebenia*). Ellice Is, Funafuti Atoll, Tuvalu. HT ♂ AMS. Muscidae, *Musca vetustissima* Walker. See Pont 1968: 90.
- nigrina* Fallén, 1817: 241 (*Musca*). Sweden. T A NRS. Tachinidae, *Thelaira nigrina* Fallén. N. Status
- nigrina* Meigen, 1838: 305 (*Musca*). Germany, Stolberg. HT ♂ MNHNP. Calliphoridae, *Pollenia amentaria* Scopoli. Preocc. Fallén 1817. See Rognes 1991a: 221.
- nigripennis* Fabricius, 1794: 346 (*Musca*). France. T A UNKNOWN. Phaeomyiidae, *Pelidnoptera nigripennis* Fabricius.
- nigripes* Gmelin, 1790: 2859 (*Musca*). Europe. T A NMI. Tachinidae?

- nigripes* Fabricius, 1794: 341 (*Musca*). France. T A MNHNP. Dolichopodidae,  
*Chrysotus laesus* Wiedemann. Preocc. Gmelin 1790. N. Syn.
- nigripes* Fabricius, 1794: 319 (*Musca*). Germany. T A UZMCF. Tachinidae, *Thelaira nigrina* Fallén. Preocc. Gmelin 1790. N. Comb.
- nigripilis* Gmelin, 1790: 2859 (*Musca*). Europe. T A NMI.
- nigrita* Scopoli, 1763: 328 (*Musca*). Idrija. T A DESTROY. Sarcophagidae, *Nyctia halterata?* Panzer.
- nigrita* Fabricius, 1781: 427 (*Syrphus*). Jamaica. T A Unknown. Syrphidae, *Sterphus jamaicensis* Gmelin. Preocc. Scopoli 1763.
- nigrita* Gmelin, 1790: 2832 (*Musca*). Subsequent Usage of *Bibio nigritus* Fabricius. Bombyliidae, *Thyridanthrax fenestratus* Fallén.
- nigrita* Fallén, 1823: 60 (*Musca*). Sweden, Gotland, Kjellunge. HT ♂ Lost. Muscidae, *Hebecnema vespertina* Fallén. Preocc. Scopoli 1763. See Pont 1984: 290.
- nigrithorax* Stein, 1909: 212 (*Musca*). Java, Semarang. LT ♂ ITZA. Muscidae, *Musca ventrosa* Wiedemann. Lectotype designated by Pont 1970a: 96.
- nigromarginata* Stephens, 1829a: 60 (*Musca*). *Nomen nudum*. England. T A BMNH. Calliphoridae.
- nitens* de Villers, 1789a: 549 (*Musca*). Southern France. T A Author. Syrphidae, *Ferdinandea cuprea* Scopoli.
- nitens* Zetterstedt, 1845: 1340 (*Musca*). N. n. *Musca vespollo sensu* Meigen. Sweden. LT ♂ ZIL. Calliphoridae, *Pollenia amentaria* Scopoli. Preocc. de Villers 1789. Lectotype designated by Rognes 1991a: 221.
- nitescens* Walker, 1861d: 302 (*Musca*). Aru Islands, Makian. T ♀ BMNH. Calliphoridae, *Lucilia papuensis* Macquart.
- nitida* Wiedemann, 1830: 410 (*Musca*). *Nomen nudum*. T A NMW. Muscidae, *Morellia nigricosta* Hough. Published in synonymy, not subsequently validated by usage. Proposed in synonymy with *violacea* Fabricius.
- nivalis* Bjerkander, 1777: 34 (*Musca*). Sweden. T A UNKNOWN.
- niveisquama* Thomson, 1868: 547 (*Musca*). China, Philippines, Singapore. T A NRS. Muscidae, *Musca sorbens* Wiedemann.
- nobilis* Gmelin, 1790: 2829 (*Musca*). N. n. *Bibio nobilitata* Fabricius. Therevidae, *Thereva nobilitata* Fabricius.
- nobilis* Meigen, 1826: 56 (*Musca*). Germany. ST ♂ ♀ NMW. Calliphoridae, *Lucilia sericata* Meigen. Preocc. Gmelin 1790. See Collin 1926: 259.
- nobilis* Gistel, 1856: 430 (*Musca*). *Nomen nudum*. Egypt. T A UNKNOWN. Published without diagnosis or indication.
- nobilitata* Linnaeus, 1767: 995 (*Musca*). Europe. T A LSL. Dolichopodidae, *Poecilobothrus nobilitatus* Linnaeus. 3 sp. in LSL (Dear MS).
- nobilitata* Fabricius, 1775: 757 (*Bibio*). Denmark. T A UZMCF. Therevidae, *Thereva nobilitata* Fabricius.
- noctiluca* Linnaeus, 1758a: 593 (*Musca*). Sweden. LT ♀ LSL. Syrphidae, *Pipiza noctiluca* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 158.
- nora* Walker, 1849: 910 (*Musca*). France & Lapland. LT ♂ BMNH. Muscidae, *Myospila meditabunda* Fabricius. Lectotype designated by Pont 1970b: 111.
- norwegica* Gmelin, 1790: 2865 (*Musca*). N. n. *Rhagio fasciatus* Fabricius. Rhagionidae.
- nosatus* Harris, 1780: 105 (*Musca*). Incosp. *nasatus* Harris. Syrphidae, *Rhingia* sp.

- notata** Fabricius, 1781: 448 (*Musca*). England. T A Yeats. Dolichopodidae, *Scellus notatus* Fabricius.
- notata** Estlund, 1796: 130 (*Musca*). Sweden, Elfskarby. T A UNKNOWN. Preocc. Fabricius 1781.
- notata** Fallén, 1823: 62 (*Musca*). Sweden, Skåne, Äsperöd. LT ♂ NRS. Muscidae, *Limnophora tigrina* Am Stein. Preocc. Fabricius 1781. Lectotype designated by Pont 1984:291.
- nova** Schrank, 1776: 93 (*Musca*). Austria. T A Unknown. Syrphidae, *Microdon* sp.
- novemmaculata** de Villers, 1789b: cxxxvii (*Musca*). N. n. *Musca tuberculata* de Villers [1789a: 428]. Bibionidae, *Dilophus novemmaculatus* de Villers. Published as *9-maculata*.
- novemmaculata** Gmelin, 1790: 2862 (*Musca*). Europe. T A NMI. Preocc. de Villers 1789.
- novempunctata** Gmelin, 1790: 2862 (*Musca*). Europe. T A NMI.
- nugator** Harris, 1780: 120, pl. 35, fig. 36 (*Musca*). Incosp. *nugator* Harris. Muscidae, *Phaonia rufiventris* Scopoli.
- nugator** Harris, 1780: index (*Musca*). England. T ♀ Unknown. Muscidae, *Phaonia rufiventris* Scopoli.
- obesa** Gmelin, 1790: 2868 (*Musca*). Subsequent Usage of *Syrphus obesus* Fabricius. Syrphidae, *Ornidia obesa* Fabricius.
- obfuscata** Gmelin, 1790: 2880 (*Musca*). N. n. *Conops fuscus* Scopoli. Syrphidae, *Eristalis interrupta* Poda.
- obscoena** Eschscholtz, 1822: 113 (*Musca*). Alaska, "Insel Unalaschka, an der Meeresküste". T A Unknown. Calliphoridae, *Calliphora vomitoria* Linnaeus.
- obscura** Fabricius, 1794: 315 (*Musca*). Germany. HT ♂ UZMC F. Calliphoridae, *Pollenia rufis* Fabricius. See Rognes 1987: 496.
- obscura** Meigen, 1830: 374 (*Musca*). Germany, Berlin. T A MNHNP. Calliphoridae. Preocc. Fabricius 1794.
- obscura** Walker, 1853b: 346 (*Musca*). Senegal. T ♂ BMNH. Muscidae, *Pyrellina distincta* Walker. Preocc. Fabricius 1794.
- obscura** Pont, 1968: 89 (*Degeeria*). Ellice Island, Funafuti Atoll, Tuvalu. ST ♀ AMS. Muscidae, *Musca domestica* Linnaeus. Published in synonymy, not subsequently validated by usage. Rainbow MS name cited in synonymy with *domestica* Linnaeus.
- obscurata** Walker, 1858b: 105 (*Musca*). Moluccas, Aru Islands. T ♀ Lost. Calliphoridae, *Onesia obscurata* Walker. Types not in BMNH (Pont).
- obsidianus** Harris, 1776: 37, pl. 9, fig. 8 (*Musca*). England. T ♀ Unknown. Tachinidae, *Zophomyia temula* Scopoli.
- obsoleta** Fallén, 1817: 238 (*Musca*). Sweden. LT ♀ NRS. Sarcophagidae, *Sarcophila latifrons* Fallén. Lectotype designated by Pape 1986: 305.
- obtrusa** Walker, 1858b: 105 (*Musca*). Aru Islands. LT ♀ BMNH. Muscidae, *Neomyia timorensis* Robineau-Desvoidy. Lectotype designated by Pont 1966: 94.
- obtusipennis** Fallén, 1823: 57 (*Musca*). Sweden, Skåne, Äsperöd. LT ♂ NRS. Muscidae, *Helina obtusipennis* Fallén. Lectotype designated by Pont 1984: 291.
- occidentis** Walker, 1853b: 347 (*Musca*). United States. T ♀ BMNH. Muscidae, *Eudasyphepha cyanicolor* Zetterstedt. Preocc. Walker 1853: 332.
- occidentis** Walker, 1853b: 332 (*Musca*). United States. HT ♂ BMNH. Muscidae, *Stomoxys calcitrans* Linnaeus. See Pont 1990: 167.

- occidua** Fabricius, 1794: 315 (*Musca*). West Indies. T A UZMC F. Sarcophagidae,  
*Tricharaea occidua* Fabricius. See Pape 1986: 302.
- oceanica** Le Guillou, 1842: 316 (*Musca*). Tasmania, Hobart. LT ♂ MNHNP. Muscidae, *Musca domestica* Linnaeus. Lectotype designated by Pont 1973: 170.
- ochracea** Ratzeburg, 1840: 125 (*Musca*). Germany. Tachinidae, *Zenillia fulva* Fallén.
- ochrapesus** Rathvon, 1855: 10 (*Musca*). Pennsylvania, Lancaster. T A Unknown. Chyromyidae, *Chyromya flava* Linnaeus.
- ochricornis** Wiedemann, 1830: 408 (*Musca*). Brazil. T A ZMHU NMW. Muscidae, *Morellia ochricornis* Wiedemann.
- ochroleuca** Gmelin, 1790: 2878 (*Musca*). Europe. T A NMI. Syrphidae.
- ochroptera** Gmelin, 1790: 2860 (*Musca*). Europe. T A NMI.
- ochropus** Turton, 1801: 635 (*Musca*). N. n. *Musca flavipes* Fabricius. Therevidae, *Dialineura anilis* Linnaeus.
- octopunctata** Coquebert, 1798: 145 (*Musca*). France, Reims. T A MNHNP. Otitidae, *Myennis octopunctata* Coquebert.
- ocymi** Turton, 1801: 651 (*Musca*). Subsequent Usage of *Syrphus ocymi* Fabricius. Syrphidae, *Platycheirus granditarsus* Forster.
- oenopota** Scopoli, 1763: 337 (*Musca*). Carniola. T A DESTROY. Drosophilidae, *Drosophila funebris?* Fabricius.
- oestracea** Linnaeus, 1758a: 592 (*Musca*). Sweden. LT ♀ LSL. Syrphidae, *Eristalis oestracea* Linnaeus. Lectotype designated by Thompson et al. 1982: 158.
- oleae** Gmelin, 1790: 2844 (*Musca*). France & Italy. T A Unknown. Tephritidae, *Dacus oleae* Rossi. Preocc. Rossi 1790.
- oleae** Rossi, 1790: 317 (*Musca*). Italy, Tuscany, Firenze & Pisa. T A ZMHU. Tephritidae, *Dacus oleae* Rossi.
- oleae** Petagna, 1792: 685 (*Musca*). Italy, Calabria. T A UNKNOWN. Tephritidae, *Dacus oleae* Rossi. Preocc. Rossi 1790.
- oleae** Fabricius, 1794: 349 (*Musca*). Subsequent Usage of *Musca oleae* Rossi. Tephritidae, *Dacus oleae* Rossi.
- olens** Fuesslin, 1775: 54 (*Musca*). Switzerland, Utliberg. T A Unknown. Xylophagidae, *Coenomyia ferruginea* Scopoli. N. Syn.
- olivacea** Meigen, 1838: 308 (*Musca*). Subsequent Usage of *Pollenia olivacea* Macquart. Calliphoridae, *Pollenia rudis* Fabricius.
- onopordinis** Fabricius, 1775: 787 (*Musca*). Denmark. T A UZMC F. Tephritidae, *Euleia heraclii* Linnaeus.
- oppidana** Scopoli, 1763: 349 (*Musca*). Carniola. T A DESTROY. Chyromyidae, *Chyromya oppidana* Scopoli.
- optata** Walker, 1859: 137 (*Musca*). Sulawesi, Makassar. T ♂ BMNH. Muscidae, *Neomyia diffidens* Walker.
- opulenta** Walker, 1858b: 104 (*Musca*). Moluccas, Aru Islands. HT ♀ BMNH. Calliphoridae, *Platytropesa auriceps* Macquart. See Crosskey 1965: 85.
- oratoria** Turton, 1801: 614 (*Musca*). Emend. *aratoria* Fabricius. Sciomyzidae, *Ilione albisetosa* Scopoli.
- orbiculata** Geoffroy, 1785: 484 (*Musca*). France, Paris. T A Author.
- orbiculus** Gmelin, 1790: 2875 (*Musca*). Subsequent Usage of *Syrphus orbiculus* Fabricius. Acroceridae, *Acrocera orbiculus* Fabricius.
- orientalis** Walker, 1849: 870 (*Musca*). Subsequent Usage of *Lucilia orientalis* Macquart. Calliphoridae, *Chrysomya rufifacies* Macquart.

- orior Harris, 1780: 143, pl. 42, fig. 48 (*Musca*). England. Calliphoridae, *Protophor-mia terraenovae* Robineau-Desvoidy. Name improperly formed (verb).
- ornate Harris, 1780: 146, pl. 43, fig. 84 (*Musca*). England. T A Unknown. Muscidae, *Azelia nebulosa* Robineau-Desvoidy. Name improperly formed (adverb).
- ornithovora Scopoli, 1763: 342 (*Musca*). Carniola. T A DESTROY. Piophilidae?
- osiris Wiedemann, 1830: 420 (*Musca*). Egypt. HT ♂ NMW. Muscidae, *Musca osiris* Wiedemann. See Pont 1986b: 251.
- ovipara Portschinsky, 1910: 25, fig. 21 (*Musca*). European Russia. ST ♂ ♀ ASL. Muscidae, *Musca autumnalis* De Geer.
- pabulorum Fallén, 1817: 252 (*Musca*). Sweden. LT ♀ ZIL. Muscidae, *Muscina prolapsa* Harris. Lectotype designated by Pont 1984: 291.
- pacifica Gmelin, 1790: 2872 (*Musca*). N. n. *Syrphus cingulatus* Fabricius. Syrphidae, *Helophilus cingulatus* Fabricius.
- pagana Fabricius, 1794: 326 (*Musca*). Denmark. LT ♂ UZMC. Muscidae, *Mydaea corni* Scopoli. Preocc. Harris 1780. Lectotype designated by Michelsen 1979a: 187.
- paganus Harris, 1780: 153, pl. 45, fig. 84 (*Musca*). England. T A Unknown. Scathophagidae.
- pallens Walker, 1849: 868 (*Musca*). Subsequent Usage of *Bengalia pallens* Robineau-Desvoidy. Calliphoridae, *Bengalia labiata* Robineau-Desvoidy.
- pallida Fabricius, 1787: 344 (*Musca*). Denmark, Copenhagen. LT ♂ UZMC F. Muscidae, *Phaonia pallida* Fabricius. Lectotype designated by Michelsen 1979a: 187.
- pallida de Villers, 1789a: 509 (*Musca*). Sweden, ‘Oelandica’. T A Unknown. Preocc. Fabricius 1787.
- pallipes Gmelin, 1790: 2834 (*Musca*). Subsequent Usage of *Stratiomys pallipes* Fabricius. Stratiomyidae, *Stratiomys pallipes* Fabricius.
- pallipes Fabricius, 1794: 340 (*Musca*). Germany. T A UZMC F. Dolichopodidae, *Syntormon pallipes* Fabricius.
- palmipes Rossi, 1794: 74 (*Musca*). Italy. T A ZMHU. Platypezidae, *Platypeza fasciata*? Meigen.
- palpator Harris, 1780: 119, pl. 35, fig. 37 (*Musca*). England. T A Unknown. Sepsidae, *Orygma* sp.
- palustris Scopoli, 1763: 345 (*Musca*). Carniola. T A DESTROY. Syrphidae.
- pampasiana Bigot, 1888: 607 (*Musca*). Argentina, Buenos Aires. HT ♂ BMNH. Muscidae, *Musca domestica* Linnaeus. See Pont 1990: 167.
- pandia Walker, 1849: 898 (*Musca*). France? HT ♀ BMNH. Calliphoridae, *Bellardia pandia* Walker. See Schumann 1974: 253.
- pantherina Linnaeus, 1758a: 590 (*Musca*). Sweden. T A LSL. Stratiomyidae, *Nemotelus pantherinus* Linnaeus. 1 sp. in LSL (Dear MS).
- papatasi Gmelin, 1790: 2866 (*Musca*). Subsequent Usage of *Bibio papatasi* Scopoli. Psychodidae, *Phlebotomus papatasi* Scopoli.
- paralleli Harris, 1782: 42 (*Musca*). Missp. *parralleli* Harris. Syrphidae, *Eristalis arbustorum* Linnaeus.
- parallelus Harris, 1776: 57, pl. 1, fig. 8 (*Musca*). England. T A Unknown. Syrphidae, *Helophilus parallelus* Harris.
- parasitica Hartig, 1838: 301 (*Musca*). Germany, Brunswick. T A UNKNOWN. Tachinidae.

- parcus** Harris, 1780: 144, pl. 42, fig. 56 (*Musca*). England. T A Unknown. Rhinophoridae, *Rhinophora lepida* Meigen.
- parietina** Linnaeus, 1758a: 599 (*Musca*). Europe. T A LSL. Tephritidae, *Oxyna parietina* Linnaeus. See White 1987: 104.
- parralleli** Harris, 1776: 43, pl. 10, fig. 7 (*Musca*). England. T A Unknown. Syrphidae, *Eristalis arbustorum* Linnaeus.
- parrallelus** Harris, 1780: index (*Musca*). Missp. *parallelus* Harris. Syrphidae, *Helophilus parallelus* Harris.
- paralleli** Harris, 1776: 43 (*Musca*). Incosp. *parralleli* Harris. Syrphidae, *Eristalis arbustorum* Linnaeus. Name revised by Goffe 1946: 74.
- parviceps** Zetterstedt, 1845: 1325 (*Musca*). *Nomen nudum*. Muscidae, *Pyrellia rapax* Harris. Published in synonymy, not subsequently validated by usage. Proposed in synonymy with *aenea* Zetterstedt.
- parvula** Fallén, 1825: 75 (*Musca*). Sweden, Skåne, Äsperöd. LT ♀ ZIL. Anthomyiidae, *Subhylemyia longula* Fallén. Preocc. Harris 1776. Lectotype designated by Michelsen 1983: 114.
- parvula** Meigen, 1826: 55 (*Musca*). Germany. ST ♂ NMW. Calliphoridae, *Lucilia illustris* Meigen. Preocc. Harris 1776. See Collin 1926: 259.
- parvulus** Harris, 1776: 48, pl. 11, fig. 11 (*Musca*). England. T ♀ Unknown. Stratiomyidae, *Microchrysa polita* Linnaeus.
- pascuorum** Meigen, 1826: 74 (*Musca*). Germany, probably Stolberg. LT ♀ MNHNP. Muscidae, *Muscina pascuorum* Meigen. Lectotype designated by Pont 1986b: 236.
- patersoni** Zielke, 1971c: 133 (*Musca*). Zaire, P. N. A. Mubiliba. HT ♂ SAIMR. Muscidae, *Musca patersoni* Zielke.
- patients** Walker, 1858: 106 (*Musca*). Moluccas, Aru Islands. T ♀ BMNH. Calliphoridae, *Polleniopsis patients* Walker.
- pattoni** Austen, 1910: 115 (*Musca*). India, Madras, Saidapet, Guindy. LT ♂ BMNH. Muscidae, *Musca pattoni* Austen. Lectotype designated by Pont 1990: 168.
- pavida** Illiger, 1807: 465 (*Musca*). Germany. T A ZMHU. Tachinidae.
- pedella** Fallén, 1825: 88 (*Musca*). Sweden, Skåne, Äsperöd. LT ♀ NRS. Muscidae, *Coenosia pedella* Fallén. Lectotype designated by Pont 1984: 291.
- pedissequus** Harris, 1776: 61, pl. 15, fig. 19 (*Musca*). England. T A Unknown. Syrphidae, *Xanthogramma pedissequum* Harris.
- pedo** Harris, 1780: 115, pl. 34, fig. 15 (*Musca*). England. T A Unknown. Micropezidae.
- pellita** Forster, 1770: 15 (*Musca*). Subsequent Usage of *Tabanus pellitus* Sulzer. Tabanidae, *Heptatoma pellucens* Fabricius.
- pellucens** Linnaeus, 1758a: 595 (*Musca*). Sweden. LT ♀ LSL. Syrphidae, *Volucella pellucens* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 159.
- pellucens** Meigen, 1835: 71 (*Musca*). Germany, Bavaria, Munich. HT ♀ DESTROY. Muscidae, *Musca domestica* Linnaeus. Preocc. Linnaeus 1758. See Pont 1986b: 237.
- pellucens** Meigen, 1838: 303 (*Musca*). Germany, Bavaria. T ♀ DESTROY. Muscidae, *Musca domestica* Linnaeus. Preocc. Linnaeus 1758.
- pendula** Linnaeus, 1758a: 591 (*Musca*). Sweden. LT ♀ LSL. Syrphidae, *Helophilus pendulus* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 159.
- pennipes** Forster, 1770: 15 (*Musca*). *Nomen nudum*. England. T A UNKNOWN.

- pennipes** Fabricius, 1781: 450 (*Musca*). North America. HT ♂ UZMCF. Tachinidae, *Trichopoda pennipes* Fabricius. See Townsend 1931a: 83.
- peragro** Harris, 1780: 119, pl. 35, fig. 29 (*Musca*). England. T A Unknown. Micropezidae. Name improperly formed (verb).
- percussus** Harris, 1780: 120, pl. 35, fig. 37 (*Musca*). England. T A Unknown. Psilidae.
- perelegand** Harris, 1780: 74, pl. 21, fig. 5 (*Musca*). England. T A Unknown. Tephritidae, *Trypetia zoe* Meigen.
- perelegans** Harris, 1782: index (*Musca*). Missp. *perelegand* Harris. Tephritidae, *Trypetia zoe* Meigen.
- perexilis** Harris, 1780: 81, pl. 24, fig. 24 (*Musca*). England. T A Unknown. Syrphidae, *Baccha perexilis* Harris.
- perexillis** Harris, 1780: index (*Musca*). Incosp. *perexilis* Harris. Syrphidae, *Baccha perexilis* Harris. Name status by present revision.
- perfixa** Walker, 1856a: 26 (*Musca*). Malaysia, Mt. Ophir. T ♀ BMNH. Muscidae, *Neomyia diffidens* Walker.
- perlata** Walker, 1861a: 314 (*Musca*). South Africa, Natal. T ♀ BMNH. Calliphoridae, *Pericallimyia perlata* Walker.
- permundus** Harris, 1782: index (*Musca*). Missp. *purmundus* Harris. Tephritidae, *Anomoia purmundus* Harris.
- pernix** Harris, 1780: 84, pl. 25, fig. 13 (*Musca*). England. T A Unknown. Sarcophagidae, *Ravinia pernix* Harris.
- personata** Geoffroy, 1785: 476 (*Musca*). France, Paris. T A Author. Preocc. Harris 1780.
- personatus** Harris, 1780: 79, pl. 24, fig. 20 (*Musca*). England, London, Stepney. T A Unknown. Syrphidae, *Pocota personata* Harris.
- pertinax** Linnaeus, 1764b: 331 (*Musca*). Sweden. T A LSL.
- pervenio** Harris, 1780: 139, pl. 41, fig. 23 (*Musca*). England. Calliphoridae, *Protocalliphora azurea* Fallén. Name improperly formed (verb).
- petiolata** Turton, 1801: 612 (*Musca*). N. n. *Musca compressa* Fabricius. Tachinidae, *Cordyligaster petiolata* Fabricius.
- petronella** Linnaeus, 1758a: 598 (*Musca*). Europe. T A LSL. Micropezidae, *Calobata petronella* Linnaeus. 3 sp. in LSL (Dear MS).
- phalaenarum** Scharfenberg, 1805: 1003 (*Musca*). Germany. T A Unknown. Tachinidae.
- phara** Walker, 1849: 889 (*Musca*). Unknown. T ♀ BMNH. Muscidae, *Neomyia peronii* Robineau-Desvoidy.
- phasiaeformis** Meigen, 1826: 72 (*Musca*). Austria or Southern France. LT ♂ MNHNP. Muscidae, *Musca tempestiva* Fallén. Lectotype designated by Pont 1986b: 238.
- phauda** Walker, 1849: 896 (*Musca*). Galapagos. T ♂ BMNH. Calliphoridae, *Cochliomyia macellaria* Fabricius.
- phellia** Walker, 1849: 884 (*Musca*). North Bengal. T ♀ BMNH. Calliphoridae, *Hemipyrellia pulchra* Wiedemann.
- pica** Fabricius, 1805: 293 (*Musca*). South America. HT ♂ UZMC. Tachinidae, *Rhamphinina pica* Fabricius. See Townsend 1931a: 100.
- picata** Gmelin, 1790: 2862 (*Musca*). Europe. T A NMI.
- picea** Gmelin, 1790: 2860 (*Musca*). Europe. T A NMI.

- piceae* Manuel, 1811: 39 (*Musca*). Emend. *picea* Gmelin.
- picta* Fabricius, 1781: 452 (*Musca*). North America. T A UNKNOWN. Otitidae, *Delphinia picta* Fabricius.
- pigra* Geoffroy, 1785: 499 (*Musca*). France, Paris. T A Author. Chloropidae, *Meromyza saltatrix* Linnaeus. N. Syn.
- pigra* Schrank, 1803: 114 (*Musca*). Germany, Bavaria, Gern & Ingolstadt. T A Unknown. Syrphidae, *Xylota ignava* Panzer. Preocc. Geoffroy 1785.
- pilifacies* Emden, 1965: 58 (*Musca*, as *interrupta* ssp.). Taiwan, Chia-i, U-fen, Mt. Lu-lin. HT ♂ BMNH. Muscidae, *Musca pilifacies* Emden. See Pont 1990: 168.
- pilosa* Drury, 1773: 109 (*Musca*). Jamaica. T A Author. Tachinidae, *Archytas pilosus* Drury.
- pilosa* Bloch, 1776: 176, pl. 4, fig. 16 & 184, pl. 5, fig. 16 (*Musca*). Nomina nuda. While the word *pilosa* is italicized so that *Musca pilosa* appears to be a binomen, Bloch was not describing new species. The word *pilosa* is only the second word of a diagnosis which was set in italics by mistake.
- pilosa* Awati, 1916b: 138 (*Musca*). India. ST ♀ CRIK. Muscidae, *Musca bezzii* Patton & Cragg. Preocc. Drury 1773.
- pinastri* De Geer, 1776: 113, pl. 7, fig. 7 (*Musca*). Sweden. T A NRS. Syrphidae, *Dasyphorus pinastri* De Geer. The syntypes of *Musca pinastri* are lost. They were not present when the De Geer Collection was accessioned in 1834 (Persson, *in litt.*). However, the species is easily recognized from the figures provided by De Geer. We select as lectotype of *pinastri* the specimen which was figured (De Geer 1776: pl. 7, fig. 1). The lectotype of *lunulatus* Meigen is a dark specimen of *Dasyphorus venustus* Meigen as was pointed out by Verrall (1901a: 353). *Musca pinastri* De Geer, however, is an available name for *lunulatus* of authors, as pointed out by Lundbeck (1917: 284).
- pinguis* Gmelin, 1790: 2868 (*Musca*). Subsequent Usage of *Syrphus pinguis* Fabricius. Syrphidae, *Meromacrus pinguis* Fabricius.
- pinguis* Fabricius, 1805: 302 (*Musca*). South America. HT ♂ UZMC. Tachinidae, *Winthemia pinguis* Fabricius. See Townsend 1931b: 174.
- piniperdae* Ratzeburg, 1844: 174 (*Musca*). Germany. T A Unknown. Tachinidae, *Gonia picea* Robineau-Desvoidy.
- pinivorae* Ratzeburg, 1844: 173 (*Musca*). Germany. T A Unknown. Tachinidae, *Blondelia nigripes* Fallén.
- pionia* Walker, 1849: 880 (*Musca*). Galapagos. T ♂ BMNH. Calliphoridae, *Lucilia pionia* Walker.
- pieniens* Linnaeus, 1758a: 594 (*Musca*). Sweden. LT ♀ LSL. Syrphidae, *Syritta pipiens* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 159.
- pipines* Geoffroy, 1785: 486 (*Musca*). Missp. *pieniens* Linnaeus. Syrphidae, *Syritta pipiens* Linnaeus.
- placida* Müller, 1764: 85 (*Musca*). Denmark, Sjaelland, Frederiksdal. T A UNKNOWN. Tephritidae, *Urophora stigma?* Loew.
- placida* Walker, 1853b: 343 (*Musca*). South Australia. T ♂ BMNH. Calliphoridae, *Calliphora dubia* Macquart. Preocc. Müller 1764.
- planiceps* Wiedemann, 1824: 48 (*Musca*). Java. HT ♂ UZMC. Muscidae, *Musca planiceps* Wiedemann. See Townsend 1931c: 369.

- planifrons** Fabricius, 1798: 565 (*Musca*). France, Paris. T A MNHNP. Chloropidae,  
*Platycephala planifrons* Fabricius.
- plebeja** Linnaeus, 1758a: 589 (*Musca*). Europe. T A LSL. Therevidae, *Thereva plebeja* Linnaeus.
- plebeja** Fabricius, 1781: 440 (*Musca*). Germany, Kiel. T A UZMC F. Tachinidae,  
*Dexia rustica* Fabricius. Preocc. Linnaeus 1758.
- plombea** de Villers, 1789a: 463 (*Musca*). Missp. *plumbea* Geoffroy.
- plumata** De Geer, 1776: 134, pl. 8, fig. 9 (*Musca*). Sweden. T A NRS. Syrphidae,  
*Volucella bombylans* Linnaeus.
- plumbea** Geoffroy, 1785: 492 (*Musca*). France, Paris. T A Author.
- plumipes** Scopoli, 1763: 334 (*Musca*). Carniola. T A DESTROY. Dolichopodidae,  
*Dolichopus plumipes* Scopoli.
- plumosa** Gmelin, 1790: 2868 (*Musca*). Europe. T A NMI. Syrphidae, *Volucella bombylans* Linnaeus.
- pluvialis** Linnaeus, 1758a: 597 (*Musca*). Europe. HT ♂ LSL. Anthomyiidae, *Anthomyia pluvialis* Linnaeus. See Pont 1981: 172.
- podagrlica** Gmelin, 1790: 2873 (*Musca*). Subsequent Usage of *Syrphus podagrica* Fabricius. Syrphidae, *Neoascia podagrlica* Fabricius.
- poeciloptera** Schrank, 1776: 96, pl. 3, fig. 22 (*Musca*). Austria, Linz. T A Unknown.  
 · Tephritidae, *Platyparea poeciloptera* Schrank.
- polisma** Walker, 1849: 878 (*Musca*). Unknown. T ♂ DESTROY. Calliphoridae.
- polita** Linnaeus, 1758a: 598 (*Musca*). Europe. T A LSL. Stratiomyidae, *Microchrysa polita* Linnaeus. 1 sp. in LSL (Dear MS).
- politula** Walker, 1853b: 338 (*Musca*). East Indies. T ♂ BMNH. Muscidae, *Neomyia lauta* Wiedemann. Preocc. Linnaeus 1758.
- politula** Malloch, 1929b: 112 (*Lissasterna*). Namibia, Farm Ojtjueza, 66km NE Windhoek. HT ♂ ZSZMH. Muscidae, *Musca lasiophthalma* Thomson. Preocc. Walker 1853. See Zielke 1971c: 109 & 1971d: 174.
- politus** Harris, 1780: 81, pl. 24, fig. 26 (*Musca*). England. T A Unknown. Syrphidae, *Lejogaster metallina* Fabricius. Preocc. Linnaeus 1758.
- pollinosa** Stein, 1909: 211 (*Musca*). Java, Batavia. LT ♂ ITZA. Muscidae, *Musca planiceps* Wiedemann. Lectotype designated by Pont 1970a: 99.
- polyzonias** Lepechin, 1775: 211, pl. 11, fig. 29 (*Musca*). *Nomen nudum*. [Russia]. Bombyliidae. Published in non-binominal work.
- polyzonias** Gmelin, 1790: 2833 (*Musca*). Siberia. T A Unknown. Bombyliidae.
- porca** Latreille, 1804: 196 (*Musca*). France. T A MNHNP. Otitidae, *Otites porca* Latreille. N. Status
- porcina** De Geer, 1776: 98 (*Musca*). N. n. *Musca tenax* Linnaeus. Syrphidae, *Eristalis tenax* Linnaeus.
- porphyrina** Walker, 1856a: 24 (*Musca*). Malaysia, Mt. Ophir. T ♀ BMNH. Calliphoridae, *Lucilia porphyrina* Walker.
- potens** Harris, 1780: 110, pl. 33, fig. 53 (*Musca*). England. T A Unknown. Syrphidae, *Myolepta potens* Harris.
- potentillae** de Villers, 1789a: 510 (*Musca*). Sweden? T A Unknown.
- praeceps** Scopoli, 1763: 333 (*Musca*). Carniola. T A DESTROY. Tachinidae, *Mintho praeceps* Scopoli.
- praecox** Walker, 1864: 236 (*Musca*). North Ceram. HT ♂ BMNH. Muscidae, *Musca conducens* Walker. See Pont 1990: 168.

- praedicens** Walker, 1859b: 139 (*Musca*). Sulawesi, Makasar. T ♀ BMNH. Muscidae, *Graphomya praedicens* Walker.
- praedo** Rossi, 1794: 75 (*Musca*). Italy. T A ZMHU.
- praepotens** Fallén, 1825: 83 (*Musca*). Subsequent Usage of *Anthomyia praepotens* Wiedemann. Anthomyiidae, *Eutrichota praepotens* Wiedemann.
- praeusta** Panzer, 1804: 209 (*Musca*). Germany, Regensburg. T A Unknown.
- prashadii** Patton, 1922b: 69 (*Musca*). India, Koharnag. LT ♂ BMNH. Muscidae, *Musca autumnalis* De Geer. Lectotype designated by Pont 1990: 168.
- prasina** Gmelin, 1790: 2879 (*Musca*). Europe. T A NMI. Syrphidae, *Cheilosia* sp.
- pratorum** Gmelin, 1790: 2870 (*Musca*). Subsequent Usage of *Syrphus pratorum* Fabricius. Syrphidae, *Meromacrus pratorum* Fabricius.
- pratorum** Meigen, 1826: 78 (*Musca*). Germany, probably Stolberg. LT ♂ MNHNP. Muscidae, *Dasyphora pratorum* Meigen. Lectotype designated by Pont 1986b: 240.
- primūtiva** Walker, 1849: 903 (*Musca*). China. HT ♀ BMNH. Muscidae, *Musca sorbens* Wiedemann. See Pont 1990: 168.
- princeps** Harris, 1780: 151, pl. 45, fig. 76 (*Musca*). England. T ♀ Unknown. Muscidae, *Mydaea corni* Scopoli.
- prisca** Walker, 1849: 903 (*Musca*). Australia. HT ♀ BMNH. Muscidae, *Musca vetustissima* Walker. See Pont 1973: 148 & 1990: 168.
- procedo** Harris, 1780: 141, pl. 41, fig. 42 (*Musca*). England. Tachinidae. Name improperly formed (verb).
- processioneae** Ratzeburg, 1840: 125 (*Musca (Tachina)*). Germany. T A UNKNOWN. Tachinidae, *Carcelia processioneae* Ratzeburg.
- processioneae** Ratzeburg, 1844: 170 (*Musca (Anthomyia)*). Germany. T A Unknown. Anthomyiidae. Preocc. Ratzeburg 1840.
- prodeo** Harris, 1780: 141, pl. 41, fig. 40 (*Musca*). England. T ♀ Unknown. Muscidae, *Muscina stabulans* Fallén. Name improperly formed (verb).
- prodromaria** Panzer, 1804: 160 (*Musca*). Germany, Regensburg. T A Unknown.
- proerna** Walker, 1849: 888 (*Musca*). Australia. HT ♀ BMNH. Muscidae, *Neomyia lauta* Wiedemann. See Pont 1973: 192.
- proferens** Walker, 1859b: 138 (*Musca*). Sulawesi, Makasar. T ♂ BMNH. Muscidae, *Pyrellia proferens* Walker.
- profuges** Harris, 1780: 81, pl. 24, fig. 25 (*Musca*). England, Essex, West Tilbury. T A Unknown. Syrphidae, *Doros profuges* Harris.
- profugus** Harris, 1782: index (*Musca*). Missp. *profuges* Harris. Syrphidae, *Doros profuges* Harris.
- prolabor** Harris, 1780: 139, pl. 41, fig. 31 (*Musca*). England. T A Unknown. Sarcophagidae, *Sarcophaga* sp. Name improperly formed (verb).
- prolapsa** Harris, 1780: 139, pl. 41, fig. 32 (*Musca*). England. T ♀ Unknown. Muscidae, *Muscina prolapsa* Harris.
- prolapsor** Harris, 1780: index (*Musca*). Incosp. *prolapsa* Harris. Muscidae, *Muscina prolapsa* Harris.
- promano** Harris, 1780: 143, pl. 42, fig. 47 (*Musca*). England. T A Unknown. Tachinidae, *Dufouria chalybeata* Meigen. Name improperly formed (verb).
- promans** Harris, 1780: 143 (*Musca*). Incosp. *promano* Harris. Tachinidae, *Dufouria chalybeata* Meigen. Name status by present revision.

- promisca* Awati, 1916b: 138 (*Musca*). India. ST ♂ ♀ CRIK. Muscidae, *Musca sorbens* Wiedemann.
- promittens* Walker, 1859b: 134 (*Musca*). Sulawesi, Makasar. T A BMNH. Calliphoridae, *Phumosia promittens* Walker.
- properans* Rossi, 1794: 74 (*Musca*). Italy. T A ZMHU. Platypezidae, *Platypeza fasciata?* Meigen.
- prospera* Walker, 1859b: 133 (*Musca*). Sulawesi, Makasar. LT ♀ BMNH. Calliphoridae, *Silbomyia albonotata* Bigot. Lectotype designated by Crosskey 1965: 56.
- prostrata* Scopoli, 1763: 328 (*Musca*). Carniola. T ♀ DESTROY. Muscidae, *Musca autumnalis* De Geer.
- provenio* Harris, 1780: 142, pl. 41, fig. 43 (*Musca*). England. T A Unknown. Tachinidae, *Dexia rustica* Fabricius. Name improperly formed (verb).
- proxima* Walker, 1853b: 341 (*Musca*). California. T ♀ BMNH. Calliphoridae, *Phormia regina* Meigen.
- pruinosa* Meigen, 1838: 306 (*Musca*). Subsequent Usage of *Pollenia pruinosa* Macquart. Calliphoridae, *Pollenia rufa* Fabricius.
- pseudocorvina* Emden, 1939: 80 (*Musca*, as *autumnalis* ssp.). Kenya, Aberdare Range, Mt Kinangop. HT ♂ BMNH. Muscidae, *Musca pseudocorvina* Emden. See Pont 1990: 168.
- psyllia* Schrank, 1803: 127 (*Musca*). Germany, Bavaria, Ingolstadt. T A Unknown.
- puber* Gmelin, 1790: 2850 (*Musca*). Missp. *pubera* Linnaeus. Scathophagidae, *Cordilura pubera* Linnaeus.
- pubera* Linnaeus, 1758a: 598 (*Musca*). Europe. T A LSL. Scathophagidae, *Cordilura pubera* Linnaeus. 4 sp. in LSL (Dear MS).
- puberula* Zetterstedt, 1837: 42 (*Musca*). *Nomen nudum*. [Sweden, Nordlandia, Tromsø]. T ♂ ZIL. Calliphoridae, *Bellardia pubicornis* Zetterstedt.
- puberula* Zetterstedt, 1838: 654 (*Musca*). Norway, Nordlandia, Tromsø. LT ♂ ZIL. Calliphoridae, *Bellardia pubicornis* Zetterstedt. Lectotype designated by Rognes 1991a: 51.
- pubiceps* Zetterstedt, 1845: 1333 (*Musca*). Sweden, Luleå. T A ZIL. Tachinidae, *Macquartia pubiceps* Zetterstedt.
- pudefactus* Harris, 1780: 121, pl. 35, fig. 42 (*Musca*). England. T A Unknown. Anthomyiidae.
- puella* Meigen, 1826: 57 (*Musca*). Germany, probably Kiel. HT ♂ NMW. Muscidae, *Neomyia cornicina* Fabricius. See Pont 1986b: 241.
- pulchella* Müller, 1776: 174 (*Musca*). Denmark. T A Author. Chloropidae, *Chlorops pulchella* Müller. N. Comb.
- pulchella* Rossi, 1790: 314, pl. 8, fig. 6 (*Musca*). Italy, Tuscany, Firenze & Pisa. T A ZMHU. Pallopteridae, *Palloptera muliebris* Harris. Preocc. Müller 1776.
- pulchra* Wiedemann, 1830: 406 (*Musca*). Unknown. T A NMW. Calliphoridae, *Hemipyrellia pulchra* Wiedemann.
- pulicaria* Geoffroy, 1785: 498 (*Musca*). France, Paris. T A Author.
- pulicaria* Schrank, 1803: 126 (*Musca*). Germany, Bavaria. T A Unknown. Preocc. Geoffroy 1785.
- pulla* Gmelin, 1790: 2859 (*Musca*). Europe. T A NMI. Syrphidae.
- pulla* Bezzi, 1911: 92 (*Musca*). South Africa, Pretoria. HT ♂ USNM. Muscidae, *Musca conducens* Walker. Preocc. Gmelin 1790.

- pullata** Müller, 1764: 86 (*Musca*). Denmark, Sjaelland, Frederiksdal. T A DESTROY.
- pumila** Fallén, 1825: 88 (*Musca*). Sweden. LT ♂ NRS. Muscidae, *Coenosia pumila* Fallén. Lectotype designated by Pont 1984: 291.
- pumila** Meigen, 1838: 307 (*Musca*). Subsequent Usage of *Pollenia pumila* Robineau-Desvoidy. Calliphoridae, *Pollenia rudis* Fabricius.
- pumila** Macquart, 1848: 218, pl. 6, fig. 12 (*Musca*). Australia. HT ♂ BMNH. Muscidae, *Musca vetustissima* Walker. Preocc. Fallén 1825. See Pont 1990: 168.
- pumilionis** Bjerkander, 1778: 201 (*Musca*). Sweden. T A UNKNOWN. Chloropidae, *Chlorops pumilionis* Bjerkander.
- pumilionis** Banks, 1791: 2 (*Musca*). England. T A BMNH B. Chloropidae, *Chlorops pumilionis* Bjerkander. Preocc. Bjerkander 1778. N. Syn.
- punctata** Poda, 1761: 116 (*Musca*). Austria, Graz. T A Unknown.
- punctata** Müller, 1764: 85 (*Musca*). Denmark. T A DESTROY. Syrphidae, *Eristalinus aeneus* Scopoli. Preocc. Poda 1761.
- punctata** Schrank, 1781: 474 (*Musca*). Austria, Vienna. T A Unknown. Tephritidae, *Orellia stictica* Gmelin. Preocc. Poda 1761. N. Comb.
- punctata** Fabricius, 1794: 347 (*Musca*). Germany, Kiel. T A UZMC F. Sciomyzidae, *Pherbellia schoenherri* Fallén. Preocc. Poda 1761.
- punctata** Walker, 1849: 868 (*Musca*). Subsequent Usage of *Ormia punctata* Robineau-Desvoidy. Tachinidae, *Ormia punctata* Robineau-Desvoidy.
- punctatus** Fabricius, 1787: 333 (*Rhagio*). Germany, Kiel. T A UZMC F. Anisopodidae, *Sylvicola punctatus* Fabricius.
- punctifera** Geoffroy, 1785: 473 (*Musca*). France, Paris. T A Author.
- punctulata** Scopoli, 1763: 338 (*Musca*). Idrija. T A DESTROY. Sciomyzidae, *Trypetoptera punctulata* Scopoli.
- punctulata** Wiedemann, 1819a: 21 (*Musca*). South Africa, Cape of Good Hope. HT ♀ UZMC. Calliphoridae, *Cosmina fuscipennis* Robineau-Desvoidy. Preocc. Scopoli 1763. See Townsend 1931c: 372.
- punctum** Fabricius, 1794: 351 (*Musca*). France. T A MNHNP. Sepsidae, *Sepsis punctum* Fabricius.
- pungens** De Geer, 1776: 78, pl. 4, fig. 12 (*Musca*). N. n. *Conops calcitrans* Linnaeus. Muscidae, *Stomoxyx calcitrans* Linnaeus.
- pungoana** Karsch, 1886: 259 (*Musca*). Angola, Pungo Andongo. T ♀ ZMHU. Muscidae, *Musca ventrosa* Wiedemann.
- puparum** Fabricius, 1794: 326 (*Musca*). Germany, Kiel. T A UZMC F. Tachinidae, *Ernestia puparum* Fabricius.
- purmundus** Harris, 1780: 74, pl. 21, fig. 6 (*Musca*). England, Kent, Dartford. T A Unknown. Tephritidae, *Anomoia purmundus* Harris.
- purpurascens** Walker, 1836: 355 (*Musca*). Brazil, Santa Catarina. T ♀ BMNH. Calliphoridae, *Lucilia purpurascens* Walker.
- purpurea** Walker, 1853b: 337 (*Musca*). South America. T ♂ BMNH. Calliphoridae, *Chloroprocta idioidea* Robineau-Desvoidy.
- pusilla** Gmelin, 1790: 2879 (*Musca*). Europe. T A NMI. Syrphidae, *Paragus* sp.
- pusilla** Meigen, 1826: 71 (*Musca*). Germany, Stolberg. LT ♂ MNHNP. Calliphoridae, *Bellardia viarum* Robineau-Desvoidy. Preocc. Gmelin 1790. Lectotype designated by Rognes 1991a: 40. N. Comb.

- pusilla* Macquart, 1848: 219 (*Musca*). Haiti. T ♀ BMNH. Sarcophagidae, *Anapuno-phyto lopesi* Thompson & Pont. Preocc. Gmelin 1790. See Lopes 1975: 541.
- pusilla* Macquart, 1851: 226 (*Musca*). Egypt. T ♂ UMO. Muscidae, *Musca sorbens* Wiedemann. Preocc. Gmelin 1790.
- pusilla* Rondani, 1868: 602 (*Musca*). *Nomen nudum*. Muscidae, *Musca tempestiva* Fallén.
- pusillus* Fabricius, 1794: 271 (*Nemotelus*). India, Tranquebar. T A UZMC F. Stratiomyidae, *Oplodontha minuta* Fabricius.
- putoria* Wiedemann, 1830: 403 (*Musca*). Sierra Leone. T A NMW. Calliphoridae, *Chrysomya chloropyga* Wiedemann.
- putrida* Fabricius, 1775: 775 (*Musca*). America. T A Auction. Calliphoridae. Type(s) originally in collection of Otto Thott.
- putrida* Fabricius, 1794: 316 (*Musca*). West Indies. T A UZMC F. Preocc. Fabricius 1775.
- putris* Linnaeus, 1758a: 597 (*Musca*). Europe. T A LSL. Sepsidae, *Themira putris* Linnaeus. 2 sp. in LSL (Dear MS).
- putris* Geoffroy, 1785: 490 (*Musca*). France, Paris & Sweden? T A Author. Preocc. Linnaeus 1758.
- pyrastri* Linnaeus, 1758a: 594 (*Musca*). Sweden. LT ♂ LSL. Syrphidae, *Scaeva pyrastri* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 159.
- pyrorum* Schrank, 1803: 114 (*Musca*). Germany, Bavaria. T A Unknown. Syrphidae, *Dasyphorus pinastri* De Geer.
- As Schrank included a reference to *Musca pinastri* De Geer in his definition of *pyrorum*, we select as lectotype of *pyrorum* the specimen illustrated by De Geer (De Geer 1776: pl. 7, fig. 1).
- quadrata* Geoffroy, 1785: 485 (*Musca*). France, Paris. T A Author.
- quadricolor* Gmelin, 1790: 2861 (*Musca*). Europe. T A NMI.
- quadrifasciata* Fabricius, 1781: 448 (*Musca*). Germany. T A UZMC F. Dolichopodidae, *Neurigona quadrifasciata* Fabricius.
- quadrifasciata* Geoffroy, 1785: 473 (*Musca*). France, Paris. T A Author. Preocc. Fabricius 1781.
- quadrilineata* Gmelin, 1790: 2835 (*Musca*). Subsequent Usage of *Stratiomys quadrilineata* Fabricius. Stratiomyidae, *Isomerocera quadrilineata* Fabricius.
- quadrilineata* Lichtenstein, 1796: 216 (*Musca*). India, Coromandel. T A Auction.
- quadrilineata* Fabricius, 1805: 286 (*Musca*). South America. HT A UZMC. Calliphoridae, *Eunesembrinellaflavipennis* Macquart. Preocc. Lichtenstein 1796. See Townsend 1931a: 69. N. Syn.
- quadrilineatus* Fabricius, 1787: 336 (*Syrphus*). India, Tranquebar. T A UZMC. Syrphidae, *Eristalinus arvorum* Fabricius.
- quadrimaculata* Fabricius, 1787: 347 (*Musca*). Germany, Kiel. T A UZMC F. Preocc. Swederus 1787.
- quadrimaculata* Swederus, 1787: 289 (*Musca*). New Zealand. HT ♀ BMNH B. Calliphoridae, *Calliphora quadrimaculata* Swederus. See Townsend 1931c: 375; Dear 1985a: 23.
- quadrimaculata* Fallén, 1823: 63 (*Musca*). Sweden, Skåne. LT ♂ NRS. Muscidae, *Helina celsa* Harris. Preocc. Swederus 1787. Lectotype designated by Pont 1984: 292.

- quadrimaculella** Hennig, 1958: 213 (*Helina*). N. n. *Musca quadrimaculata* Fallén. Muscidae, *Helina celsa* Harris.
- quadripunctata** Linnaeus, 1767: 997 (*Musca*). Sweden. NT ♂ LSL. Lauxaniidae, *Sapromyza quadripunctata* Linnaeus. Neotype designation (Yarom 1990: 290).
- quadripustulata** Fabricius, 1794: 324 (*Musca*). Germany. T A DESTROY. Tachinidae, *Winthemia quadripustulata* Fabricius. See Townsend 1932: 48.
- quadristrigata** Donovan, 1808: 72, pl. 467, fig. 3 (*Musca*). England. T A UNKNOWN. Chloropidae, *Chlorops quadristrigata* Donovan. N. Comb.
- quadrum** Fabricius, 1805: 297 (*Musca*). Austria. HT A UZMC F. Muscidae, *Helina quadrum* Fabricius. See Michelsen 1979a: 188.
- quinquefasciata** Geoffroy, 1785: 474 (*Musca*). France, Paris. T A Author.
- quinquelineata** Gmelin, 1790: 2870 (*Musca*). Subsequent Usage of *Syrphus quinquefasciatus* Fabricius. Syrphidae, *Eristalinus quinquelineatus* Fabricius.
- quinquepunctata** Schrank, 1781: 473 (*Musca*). Austria. T A Unknown.
- quinquestriata** Turton, 1801: 642 (*Musca*). Subsequent Usage of *Syrphus quinquestriatus* Fabricius. Syrphidae, *Eristalinus quinquestriatus* Fabricius.
- racilia** Walker, 1849: 884 (*Musca*). Sierra Leone. T ♂ BMNH. Muscidae, *Neomyia racilia* Walker.
- radiata** Fabricius, 1798: 565 (*Musca*). France, Paris. T A UZMC F. Tephritidae, *Trupanea radiata* Fabricius.
- radiata** Walckenaer, 1802: 399 (*Musca*). France, Paris. T A Unknown. Tephritidae, *Trupanea radiata* Fabricius. Preocc. Fabricius 1798. Probably the lack of attribution was an error.
- radicum** Linnaeus, 1758a: 596 (*Musca*). Sweden. HT ♂ LSL. Anthomyiidae, *Delia radicum* Linnaeus. See Pont 1981: 173.
- rapax** Harris, 1780: 144, pl. 42, fig. 55 (*Musca*). England. T ♀ Unknown. Muscidae, *Pyrellia rapax* Harris.
- rapax** Gmelin, 1790: 2848 (*Musca*). N. n. *Musca tigrina* Fabricius. Muscidae, *Coenosia tigrina* Fabricius. Preocc. Harris 1780.
- rara** Scopoli, 1763: 339 (*Musca*). Middle Carniola. T A DESTROY. Stratiomyidae, *Oxycera rara* Scopoli.
- rau** Robineau-Desvoidy, 1830: 396 (*Musca*). Missp. *tau* Schrank. Muscidae, *Musca autumnalis* De Geer.
- reccumbo** Harris, 1776: 35, pl. 9, fig. 4 (*Musca*). England. Tachinidae, *Tachina fera* Linnaeus. Name improperly formed (verb).
- recedans** Harris, 1780: 144, pl. 42, fig. 58 (*Musca*). England. T A Unknown. Tachinidae.
- recta** Meigen, 1838: 308 (*Musca*). Subsequent Usage of *Pollenia recta* Macquart. Calliphoridae, *Pollenia rudis* Fabricius.
- reccumbo** Harris, 1776: 35, pl. 9, fig. 4 (*Musca*). England. Tachinidae, *Tachina fera* Linnaeus. Name improperly formed (verb).
- redeambulo** Harris, 1780: 141, pl. 41, fig. 38 (*Musca*). England. Tachinidae, *Ernestia* sp. Name improperly formed (verb).
- redeans** Harris, 1782: index (*Musca*). Missp. *redians* Harris. Tachinidae.
- redeo** Harris, 1780: 140, pl. 41, fig. 34 (*Musca*). England. T ♀ Unknown. Muscidae, *Neomyia cornicina* Fabricius. Name improperly formed (verb).
- redians** Harris, 1780: 145, pl. 42, fig. 59 (*Musca*). England. T A Unknown. Tachinidae.

- reditus** Harris, 1780: 146, pl. 43, fig. 61 (*Musca*). England. T A Unknown. Anthomyiidae.
- refixa** Walker, 1856a: 26 (*Musca*). Singapore. T ♀ BMNH. Muscidae, *Neomyia diffidens* Walker.
- reflectens** Walker, 1856a: 24 (*Musca*). Malacca. T ♀ BMNH. Calliphoridae, *Isomyia viredaurea* Wiedemann.
- regalis** Meigen, 1826: 54 (*Musca*). Germany. ST ♂ NMW. Calliphoridae, *Lucilia regalis* Meigen. See Collin 1926: 258.
- regalis** Guérin-Méneville, 1831: , pl. 21, fig. 1 (*Rutilia*). Australia, A.C.T., Tharwa. NT ♂ ANIC. Tachinidae, *Rutilia regalis* Guérin-Méneville. Neotype designated by Crosskey 1973: 126.
- regalis** Guérin-Méneville, 1838: 295 (*Musca*). Subsequent Usage of *Rutilia regalis* Guérin-Méneville. Tachinidae, *Rutilia regalis* Guérin-Méneville.
- regina** Meigen, 1826: 58 (*Musca*). Germany, Stolberg. LT ♂ MNHN.P. Calliphoridae, *Phormia regina* Meigen. Lectotype designated by Rognes 1991a: 106. 1st revisor Seguy 1928: 169.
- relictus** Harris, 1780: 120, pl. 35, fig. 35 (*Musca*). England. T A Unknown. Coelopidae.
- remigro** Harris, 1780: 140, pl. 41, fig. 36 (*Musca*). England. T A Unknown. Calliphoridae, *Pollenia rufis* Fabricius. Name improperly formed (verb).
- remuria** Walker, 1849: 871 (*Musca*). China. T ♀ BMNH. Calliphoridae, *Chrysomya megacephala* Fabricius.
- repens** Harris, 1776: 37, pl. 9, fig. 10 (*Musca*). England. T A Unknown. Tachinidae.
- rerumbo** Harris, 1780: 74 (*Musca*). Missp. *reccumbo* Harris. Tachinidae, *Tachina fera* Linnaeus.
- resinellae** Schrank, 1781: 478 (*Musca*). Austria. T A Unknown. Tachinidae, *Actia nudibasis* Stein.
- resinosa** Presl, 1822: 207 (*Musca*). Ligurian Prussian Amber. T A Unknown.
- restituo** Harris, 1780: index (*Musca*). Missp. *restituo* Harris. Muscidae, *Musca autumnalis* De Geer.
- restituo** Harris, 1776: 36, pl. 9, fig. 5 (*Musca*). England. Muscidae, *Musca autumnalis* De Geer. Name improperly formed (verb).
- reticulata** Fabricius, 1781: 450 (*Musca*). Italy. T A UZMC F. Sciomyzidae, *Pherbina coryleti* Scopoli.
- reticulata** Geoffroy, 1785: 472 (*Musca*). France, Paris. T A Author. Preocc. Fabricius 1781.
- retusa** Fabricius, 1775: 775 (*Musca*). Australia. HT ♀ BMNH B. Tachinidae, *Rutilia retusa* Fabricius. See Crosskey 1973a: 137.
- reversio** Harris, 1780: 146, pl. 43, fig. 62 (*Musca*). England. T ♀ Unknown. Muscidae, *Helina reversio* Harris.
- reverto** Harris, 1780: 140, pl. 41, fig. 35 (*Musca*). England. T A Unknown. Sarcophagidae, *Sarcophaga* sp. Name improperly formed (verb).
- reviso** Harris, 1780: 141, pl. 41, fig. 39 (*Musca*). England. T A Unknown. Calliphoridae, *Protocalliphora azurea* Fallén. Name improperly formed (verb).
- revolo** Harris, 1780: 124, pl. 36, fig. 51 (*Musca*). England. T ♂ Unknown. Muscidae, *Helina impuncta* Fallén. Name improperly formed (verb).
- rex** Poda, 1761: 114 (*Musca*). Austria, Graz. T A Unknown. Stratiomyidae, *Beris* sp.

- ribesii** Linnaeus, 1758a: 593 (*Musca*). Sweden. LT ♀ LSL. Syrphidae, *Syrphus ribesii* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 159.
- riparia** Fallén, 1824: 67 (*Musca*). Sweden. LT ♂ NRS. Muscidae, *Limnophora riparia* Fallén. Lectotype designated by Pont 1984: 292.
- riparia** Robineau-Desvoidy, 1830: 398 (*Musca*). France. T ♀ DESTROY. Muscidae, *Musca autumnalis* De Geer. Preocc. Fallén 1824.
- rivalaris** Robineau-Desvoidy, 1863: 627 (*Musca*). France, Var. T ♀ DESTROY. Muscidae, *Musca domestica* Linnaeus.
- ralalis** Linnaeus, 1758a: 597 (*Musca*). Europe. HT ♀ LSL. Rhinophoridae, *Melanophora roralis* Linnaeus. See Crosskey 1977: 51.
- ralalis** Bloch, 1776: 170, pl. 3, fig. 9 (*Musca*). Subsequent Usage of *Musca roralis* Linnaeus. Rhinophoridae, *Melanophora roralis* Linnaeus.
- rosae** De Geer, 1776: 108, pl. 6, fig. 21 (*Musca*). N. n. *Musca pyrastri* Linnaeus. Syrphidae, *Scaeva pyrastri* Linnaeus.
- rosae** Fabricius, 1794: 356 (*Musca*). Germany, Kiel. T A UZMC F. Psilidae, *Chamaepsila hennigi* Thompson & Pont. Preocc. De Geer 1776.
- rosarum** Gmelin, 1790: 2877 (*Musca*). Subsequent Usage of *Syrphus rosarium* Fabricius. Syrphidae, *Platycheirus rosarium* Fabricius.
- rostrata** Fabricius, 1775: 783 (*Musca*). Denmark, Copenhagen. T A UZMC F. Dolichopodidae, *Medetera diadema* Linnaeus.
- rostrata** De Geer, 1776: 130 (*Musca*). Subsequent Usage of *Conops rostrata* Linnaeus. Syrphidae, *Rhingia rostrata* Linnaeus.
- rostrata** Preyssler, 1791: 113 (*Musca*). Subsequent Usage of *Musca rostrata* Fabricius. Dolichopodidae, *Medetera diadema* Linnaeus.
- rotundata** Linnaeus, 1758a: 596 (*Musca*). Europe. T A LSL. Tachinidae, *Gymnosoma rotundatum* Linnaeus. 1 sp. in LSL (Dear MS).
- rubiginea** Schrank, 1803: 116 (*Musca*). Germany, Bavaria, Ingolstadt. T A Unknown. Syrphidae.
- rubiginosa** Gmelin, 1790: 2878 (*Musca*). Non-European. T A NMI. Syrphidae.
- rubripes** Gmelin, 1790: 2862 (*Musca*). Europe. T A NMI.
- rudis** Fabricius, 1794: 314 (*Musca*). Germany, Schleswig-Holstein, Gromitz. NT ♂ UZMC F. Calliphoridae, *Pollenia rudis* Fabricius. Neotype designated by Rognes 1987b: 498.
- rufa** Scopoli, 1763: 335 (*Musca*). Carniola. T A DESTROY. Psilidae, *Psila fimetaria* Linnaeus.
- rufa** De Geer, 1776: 96, pl. 5, fig. 17 (*Musca*). Sweden. T A NRS. Psilidae? Preocc. Scopoli 1763.
- rufa** de Villers, 1789a: 418 (*Musca*). Subsequent Usage of *Nemotelus rufus* De Geer.
- rufa** Panzer, 1798a: 17 (*Musca*). Austria. T A ZMHU. Sciomyzidae, *Elgiva solicita* Harris. Preocc. Scopoli 1763.
- ruficapilla** Gmelin, 1790: 2860 (*Musca*). Europe. T A NMI.
- ruficauda** Fabricius, 1794: 353 (*Musca*). France. T A MNHNP. Tephritidae, *Terellia ruficauda* Fabricius.
- ruficaudis** De Geer, 1776: 127 (*Musca*). N. n. *Musca fallax* Linnaeus. Syrphidae, *Blera fallax* Linnaeus.
- ruficeps** Fabricius, 1805: 299 (*Musca*). Southern France. HT ♀ UZMC. Calliphoridae, *Rhyncomya ruficeps* Fabricius. See Townsend 1931c: 372.

- ruficeps* Meigen, 1826: 55 (*Musca*). Germany, Stolberg. T A MNHNP. Calliphoridae, *Lucilia caesar* Linnaeus. Preocc. Fabricius 1805.
- ruficeps* Meigen, 1838: 306 (*Musca*). N. n. *Pollenia violacea* Macquart. Calliphoridae, *Pollenia* sp. Preocc. Fabricius 1805.
- ruficornis* Gmelin, 1790: 2874 (*Musca*). Subsequent Usage of *Syrphus ruficornis* Fabricius. Syrphidae, *Ferdinandea ruficornis* Fabricius.
- ruficornis* Fabricius, 1794: 314 (*Musca*). India Orientali. LT ♀ UZMC. Sarcophagidae, *Parasarcophaga ruficornis* Fabricius. Lectotype designated by Pape 1986: 302.
- ruficornis* Meigen, 1838: 306 (*Musca*). Subsequent Usage of *Pollenia ruficornis* Macquart. Calliphoridae, *Pollenia vespillo* Fabricius.
- ruficornis* Walker, 1858a: 215 (*Musca*). Tasmania. T ♀ BMNH. Calliphoridae, *Onesia pubescens* Macquart. Preocc. Fabricius 1794.
- ruficerus* Ratzeburg, 1844: 174 (*Musca (Tachina)!*). Germany. T A UNKNOWN. Tachinidae.
- rufifrons* Fabricius, 1781: 449 (*Musca*). Italy. T A UZMC F. Sciomyzidae, *Euthycera fumigata* Scopoli.
- rufifrons* Fabricius, 1805: 295 (*Musca*). Guinea. HT ♂ UZMC. Sarcophagidae, *Craticulina seriata* Speiser. Preocc. Fabricius 1781. See Townsend 1931c: 379; Pape 1986: 302.
- rufifrons* Macquart, 1843: 309 (*Musca*). Java. T ♀ MNHNP. Muscidae, *Musca domestica* Linnaeus. Preocc. Fabricius 1781.
- rufina* Fallén, 1825: 92 (*Musca*). Sweden, Skåne. LT ♂ NRS. Anthomyiidae, *Pegomya rufina* Fallén. Lectotype designated by Michelsen 1983: 114.
- rufipalpis* Meigen, 1838: 307 (*Musca*). Subsequent Usage of *Pollenia rufipalpis* Macquart. Calliphoridae, *Pollenia rufidus* Fabricius.
- rufipes* Panzer, 1798c: 23 (*Musca*). Germany, Nuremberg. T A ZMHU. Sciomyzidae, *Sepedon sphaeus* Fabricius.
- rufipes* Fallén, 1825: 85 (*Musca*). Sweden, Skåne. LT ♂ NRS. Anthomyiidae, *Pegomya zonata* Zetterstedt. Preocc. Panzer 1798. Lectotype designated by Michelsen 1983: 114.
- rufipes* Descourtilz, 1826: 157, pl. 1, fig. 2 (*Musca*). France, "les bois de Meudon". T ♂ UNKNOWN. Preocc. Panzer 1798.
- rufipes* Eversmann, 1834: 428 (*Musca*). *Nomen nudum*. USSR, Volga to Urals. T A UNKNOWN. Calliphoridae. Published without diagnosis or indication.
- rufiventris* Scopoli, 1763: 329 (*Musca*). Carniola. T A DESTROY. Muscidae, *Phaonia rufiventris* Scopoli.
- rufiventris* Gmelin, 1790: 2859 (*Musca*). Europe. T A NMI. Preocc. Scopoli 1763.
- rufiventris* Fallén, 1817: 239 (*Musca*). Sweden, Kalmar. T A NRS. Tachinidae, *Mintho rufiventris* Fallén. Preocc. Scopoli 1763. N. Comb.
- rufiventris* Macquart, 1843: 312, pl. 20, fig. 9 (*Musca*). America (=Mauritius). T ♂ MNHNP. Muscidae, *Musca xanthomelaena* Wiedemann. Preocc. Scopoli 1763.
- rufiventris* Macquart, 1846: 328, pl. 17, fig. 8 (*Musca*). Brazil. LT ♂ BMNH. Muscidae, *Musca domestica* Linnaeus. Preocc. Scopoli 1763. Lectotype designated by Pont 1990: 168.
- rufomaculata* De Geer, 1776: 28, pl. 1, fig. 10 (*Musca*). Sweden. T A NRS. Tachinidae, *Eriothrix rufomaculata* De Geer.

- rugosa* Fabricius, 1794: 330 (*Musca*). Germany, Kiel. T A UZMC F. Scenopinidae, *Scenopinus niger* De Geer.
- runicis* Sommerfelt, 1827: 99 (*Musca*). Norway, S. Norrland, Saltdalen. T A UNKNOWN. Anthomyiidae, *Pegomya solennis* Meigen. See Michelsen 1985: 52.
- ruralis* Gravenhorst, 1807: 379 (*Musca*). Unknown, probably Germany. T A Lost. Muscidae.
- ruralis* Fallén, 1817: 237 (*Musca*). Sweden, Skåne. LT ♂ NRS. Sarcophagidae, *Angiometopa falleni* Pape. Preocc. Gravenhorst 1807. Lectotype designated by Pape 1986: 306.
- rustica* Fabricius, 1775: 777 (*Musca*). Denmark, Copenhagen. T A UZMC F. Tachinidae, *Dexia rustica* Fabricius.
- rustica* Schrank, 1803: 120 (*Musca*). Germany, Pottmes. T A Unknown. Tachinidae? Preocc. Fabricius 1775.
- rustica* Fallén, 1825: 79 (*Musca*, as *angelicae* var. β). Sweden. LT ♂ NRS. Muscidae, *Mydaea urbana* Meigen. Preocc. Fabricius 1775. Lectotype designated by Pont 1984: 292.
- rustica* Robineau-Desvoidy, 1863: 622 (*Musca*). France, Paris. T A DESTROY. Muscidae, *Musca autumnalis* De Geer. Preocc. Fabricius 1775.
- rutilans* Fabricius, 1781: 436 (*Musca*). West Indies. T A UZMC F. Tachinidae, *Billaea rutilans* Fabricius. Types not found by Townsend 1931a: 102.
- rutilus* Harris, 1780: 86, pl. 25, fig. 20 (*Musca*). England. T A Unknown. Tachinidae, *Mintho praeceps* Scopoli.
- rustilo* Harris, 1780: 80, pl. 24, fig. 21 (*Musca*). England. T A Unknown. Syrphidae, *Ferdinandea cuprea* Scopoli. Name improperly formed (verb).
- sabaeus* Gmelin, 1790: 2832 (*Musca*). Subsequent Usage of *Bibio sabaeus* Fabricius. Bombyliidae, *Lomatia sabaeus* Fabricius.
- sacalis* Bjerkander, 1783: 102 (*Musca*). Sweden. T A UNKNOWN.
- sacra* Fabricius, 1805: 291 (*Musca*). [South Africa, Cape of Good Hope] Error for New Zealand. HT ♀ UZMC. Calliphoridae, *Calliphora quadrimaculata* Swederus. See Townsend 1931c: 375; Dear 1985a: 23.
- saltatrix* Linnaeus, 1761: 555 (*Musca*). Sweden. T A LSL. Chloropidae, *Meromyza saltatrix* Linnaeus. 1 sp. in LSL (Dear MS).
- saltatrix* Geoffroy, 1785: 480 (*Musca*). France, Paris. T A Author. Chloropidae, *Chlorops pumilionis* Bjerkander. Preocc. Linnaeus 1758. N. Syn.
- saltitans* Scopoli, 1763: 350 (*Musca*). Carniola. T A DESTROY. Scenopinidae, *Scenopinus fenestralis* Linnaeus.
- saltuum* Linnaeus, 1758a: 599 (*Musca*). Europe. T A LSL. Pallopteridae, *Palloptera saltuum* Linnaeus. 1 sp. in LSL (Dear MS).
- salviae* Gmelin, 1790: 2877 (*Musca*). Subsequent Usage of *Syrphus salviae* Fabricius. Syrphidae, *Asarkina salviae* Fabricius.
- sanctae-helenae* Macquart, 1848: 218, pl. 6, fig. 14 (*Musca*). Saint Helena. LT ♂ IRSNB. Muscidae, *Musca domestica* Linnaeus. Lectotype designated by Pont 1977a: 127.
- sandaracha* Harris, 1780: 119, pl. 35, fig. 30 (*Musca*). England. T A Unknown. Heleomyzidae.
- sanio* Harris, 1780: index (*Musca*). Incosp. *sannio* Harris. Dolichopodidae. Name improperly formed (verb). Name status by present revision.
- saniosa* Westring, 1814: 56, pl. 4b (*Musca*). Sweden. T A NRS. Sphaeroceridae.

- sannio Harris, 1780: 158, pl. 47, fig. 5 (*Musca*). England. T A Unknown. Dolichopodidae. Name improperly formed (verb).
- santoshi** Joseph & Parui, 1972: 180 (*Musca*). India, NEFA, Kambang. HT ♂ ZSI. Muscidae, *Musca santoshi* Joseph & Parui.
- sarcophagoides** Walker, 1861b: 244 (*Musca*). New Guinea, Dorey. T ♂ BMNH. Calliphoridae, *Polleniopsis sarcophagoides* Walker.
- sarsina** Walker, 1849: 882 (*Musca*). South Africa. T ♀ BMNH. Muscidae, *Neomyia peronii* Robineau-Desvoidy.
- satyrus** Gmelin, 1790: 2831 (*Musca*). Subsequent Usage of *Bibio satyrus* Fabricius. Bombyliidae, *Ligyra satyrus* Fabricius.
- scabiosae** Fabricius, 1794: 361 (*Musca*). France. T A MNHNP. Tephritidae.
- scalaris** Fabricius, 1794: 332 (*Musca*). Denmark, Copenhagen. LT ♂ UZMC. Fanniidae, *Fannia scalaris* Fabricius. Lectotype designated by Michelsen 1979a: 188.
- scapularis** Rondani, 1875: 428 (*Musca*). Borneo, Sarawak. T ♂ MCSNG. Muscidae, *Musca conducens* Walker.
- scatophaga** Malloch, 1925: 373 (*Musca*). South Africa, Cape Province, Prospect. HT ♂ BMNH. Muscidae, *Musca aethiops* Stein. See Pont 1990: 169.
- scitula** Harris, 1780: 111, pl. 33, fig. 55 (*Musca*). England. T A Unknown. Syrphidae, *Episyphus cannabina* Scopoli. Name improperly formed (adverb).
- scitulus** Harris, 1780: 105, pl. 32, fig. 33 (*Musca*). England. T A Unknown. Syrphidae, *Episyphus cannabina* Scopoli.
- scitus** Harris, 1780: 107, pl. 32, fig. 41 (*Musca*). England. T A Unknown. Syrphidae, *Tropidia scita* Harris.
- scolopacea** Linnaeus, 1758a: 590 (*Musca*). Europe. T A LSL. Rhagionidae, *Rhagio scolopaceus* Linnaeus. 3 sp. in LSL (Dear MS).
- scolopacina** de Villers, 1789a: 429 (*Musca*). France, Bressia [=Bresse]. T A Author.
- scripta** Linnaeus, 1758a: 594 (*Musca*). Sweden. LT ♂ LSL. Syrphidae, *Sphaerophoria scripta* Linnaeus. Lectotype designated by Vockeroth 1971: 1634.
- scutata** Rossi, 1790: 314 (*Musca*). Italy, Tuscany, Firenze & Pisa. T A ZMHU. Chloropidae, *Capnoptera scutata* Rossi.
- scutellaris** Fabricius, 1805: 293 (*Musca*). South America. LT ♂ UZMC. Muscidae, *Scutellomusca scutellaris* Fabricius. Lectotype designated by Michelsen 1979a: 188 (HT designation Townsend 1931a: 68).
- scutellaris** Fallén, 1825: 90 (*Musca*). Sweden, Skåne, Åsperöd. LT ♂ NRS. Muscidae, *Phaonia rufiventris* Scopoli. Preocc. Fabricius 1805. Lectotype designated by Pont 1984: 293.
- scutellata** Meigen, 1838: 304 (*Musca*). Subsequent Usage of *Calliphora scutellata* Macquart. Calliphoridae, *Calliphora vicina* Robineau-Desvoidy.
- scybalaria** Linnaeus, 1758a: 599 (*Musca*). Europe. T A LSL. Scathophagidae, *Scathophaga scybalaria* Linnaeus. 4 sp. in LSL (Dear MS).
- secalis** Bjerkaander, 1777: 30 (*Musca*). Sweden. T A UNKNOWN.
- seelandica** Gmelin, 1790: 2870 (*Musca*). N. n. *Syrphus trilineatus* Fabricius. Syrphidae, *Helophilus seelandicus* Gmelin.
- segetum** Turton, 1801: 641 (*Musca*). Subsequent Usage of *Syrphus segetum* Fabricius. Syrphidae, *Merodon segetum* Fabricius.
- segmentaria** Panzer, 1804: 140 (*Musca*). Germany, Regensburg. T A Unknown.

- segmentaria* Fabricius, 1805: 292 (*Musca*). South America. HT ♀ UZMC. Calliphoridae, *Hemilucilia nubipennis* Rondani. Preocc. Panzer 1804. See Townsend 1931a: 71; Dear 1985b: 127. N. Syn.
- segnis* Linnaeus, 1758a: 595 (*Musca*). Sweden. LT ♂ LSL. Syrphidae, *Xylota segnis* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 160.
- selecta* Walker, 1859b: 135 (*Musca*). Sulawesi, Makasar. T ♀ BMNH. Calliphoridae, *Isomyia selecta* Walker. N. Comb.
- sellata* Fuesslin, 1775: 54 (*Musca*). Switzerland, Zürich & Bundten; Germany, Regensburg. ST A UNKNOWN. Stratiomyidae, *Clitellaria ephippium* Fabricius. N. Syn.
- semiargentata* Donovan, 1795: 92, pl. 142, fig. 3 (*Musca*). England, Epping Forest. T A UNKNOWN. Dolichopodidae, *Argyra leucocephala* Meigen.
- semiargentea* Turton, 1801: 606 (*Musca*). Emend. *semiargentata* Donovan. Dolichopodidae, *Argyra leucocephala* Meigen.
- semiatra* Wiedemann, 1830: 421 (*Musca*). Brazil. T A UZMC. Cuterebridae, *Pseudogametes semiatra* Wiedemann. HT designation for male in NMW by Townsend 1931a: 80.
- semicaerulea* Stephens, 1829a: 60 (*Musca*). *Nomen nudum*. England. T A BMNH. Calliphoridae.
- semiflava* Panzer, 1798b: 9 (*Musca*). Austria. T ♀ ZMHU.
- semiluctifera* de Villers, 1789a: 463, pl. 9, fig. 23 (*Musca*). Southern France. T A Author. Syrphidae, *Milesia semiluctifera* de Villers.
- seminationis* Fabricius, 1775: 786 (*Musca*). Germany, Leipzig. T A UZMC F. Platystomatidae, *Platystoma seminationis* Fabricius.
- semirufa* Turton, 1801: 648 (*Musca*). Subsequent Usage of *Syrphus semirufus* Fabricius. Syrphidae, *Blera fallax* Linnaeus.
- semulator* Harris, 1780: 112, pl. 33, fig. 58 (*Musca*). England. T A Unknown. Syrphidae, *Xylota* sp.
- semulator* Harris, 1780: index (*Musca*). Incosp. *semulator* Harris. Syrphidae, *Xylota* sp. Name revised by Goffe 1946: 84.
- senegalensis* Macquart, 1843: 308 (*Musca*). Senegal. T ♂ MNHNP. Muscidae, *Musca domestica* Linnaeus.
- senilis* Fabricius, 1794: 331 (*Musca*). Germany. T A UZMC F. Scenopinidae, *Scenopinus fenestralis* Linnaeus.
- senilis* Eversmann, 1834: 428 (*Musca*). *Nomen nudum*. USSR, Volga to Urals. Muscidae. Published without diagnosis or indication.
- seniorwhitei* Patton, 1922b: 73 (*Musca*). India, Krishna River & Kollar Lake. ST A BMNH. Muscidae, *Musca seniorwhitei* Patton. See Pont 1990: 169.
- sensifera* Walker, 1861a: 314 (*Musca*). Mexico. T ♂ BMNH. Tachinidae, *Billaea sensifera* Walker.
- seplasia* Walker, 1849: 891 (*Musca*). Unknown. T ♂ BMNH. Muscidae, *Eudasyphora cyanicolor* Zetterstedt.
- sepulchralis* Linnaeus, 1758a: 596 (*Musca*). Sweden. LT ♀ LSL. Syrphidae, *Eristalinus sepulchralis* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 160.
- sepulcralis* Meigen, 1826: 71 (*Musca*). Germany. ST A MNHNP. Calliphoridae, *Onesia floralis* Robineau-Desvoidy. Preocc. Linnaeus 1758.
- serena* Fallén, 1825: 76 (*Musca*). Sweden, Skåne. LT ♂ NRS. Fanniidae, *Fannia serena* Fallén. Lectotype designated by Pont 1984: 293.

- serena* Meigen, 1826: 59 (*Musca*). Germany, probably Stolberg. LT ♀ MNHNP.  
*Muscidae*, *Pyrellia rapax* Harris. Preocc. Fallén 1825. Lectotype designated by  
Pont 1986b: 243.
- serenissima* Walker, 1853b: 340 (*Musca*). East Indies. T ♀ BMNH. Calliphoridae,  
*Lucilia cuprina* Wiedemann.
- sericata* Meigen, 1826: 53 (*Musca*). Austria. T ♀ NMW. Calliphoridae, *Lucilia*  
*sericata* Meigen. Type lost (Rognes 1991a: 180).
- sericea* Geoffroy, 1785: 489 (*Musca*). France, Paris. T A Author.
- sericea* Donovan, 1808: 26, pl. 445, fig. 2 (*Musca*). England, Kent, Faversham. T A  
UNKNOWN. Tachinidae. Preocc. Geoffroy 1785.
- serpo* Harris, 1780: 144, pl. 42, fig. 54 (*Musca*). England. Sarcophagidae, *Sarcophaga*  
sp. Name improperly formed (verb).
- serrata* Linnaeus, 1758a: 597 (*Musca*). Europe. T A LSL. Heleomyzidae, *Heleomyza*  
*serrata* Linnaeus. 1 sp. in LSL (Dear MS).
- serratula* Manuel, 1811: 35 (*Musca*). Emend. *serratulae* Linnaeus. Tephritidae,  
*Terellia serratulae* Linnaeus.
- serratulae* Linnaeus, 1758a: 600 (*Musca*). Europe. T A LSL. Tephritidae, *Terellia*  
*serratulae* Linnaeus. See White 1987: 104.
- serva* Walker, 1853b: 349 (*Musca*). United States. T ♂ BMNH. Calliphoridae,  
*Opsodexia serva* Walker. N. Comb.
- setaria* Rossi, 1788: 130 (*Musca*). [Italy, Pisa]. T A UNKNOWN.
- seticornis* Gmelin, 1790: 2861 (*Musca*). Europe. T A NMI.
- setigera* Awati, 1916b: 138 (*Musca*). India. ST ♀ CRIK. Muscidae, *Musca xanthomelaena* Wiedemann.
- setosa* Gmelin, 1790: 2859 (*Musca*). Europe. T A NMI.
- setosa* Panzer, 1804: 72 (*Musca*). Germany, Regensburg. T A Unknown. Preocc.  
Gmelin 1790.
- setosa* Presl, 1822: 205 (*Musca*). Ligurian Prussian Amber. T A Unknown. Preocc.  
Gmelin 1790.
- setulosa* Zielke, 1970: 504 (*Musca*). *Nomen nudum*. Uganda, Kichwamba, Ankole.  
ST ♀ USNM. Muscidae, *Musca setulosa* Zielke. Published after 1930 without a  
description.
- setulosa* Zielke, 1971a: 104 (*Musca*). Uganda, 15mi S. Kichwamba, Kalinzu forest,  
Ankole Distr. HT ♀ USNM. Muscidae, *Musca setulosa* Zielke.
- sexdentata* Gmelin, 1790: 2836 (*Musca*). Subsequent Usage of *Stratiomys sexdentata*  
Fabricius. Stratiomyidae, *Beris chalybata* Forster.
- sexmaculata* Preyssler, 1791: 88, pl. 3, fig. 10 (*Musca*). Czechoslovakia, Prague,  
Botanical Garden. T ♀ DESTROY. Muscidae, *Helina sexmaculata* Preyssler. See  
Rozkosny, Chvála & Pont 1982: 352.
- sexpunctata* Fabricius, 1805: 300 (*Musca*). Denmark. HT ♂ UZMC. Sarcophagidae,  
*Pierretia sexpunctata* Fabricius. See Pape 1986: 303.
- shanghaiensis* Ôuchi, 1938a: 5 (*Musca*). China, Shanghai & Tienmushan. ST ♀ SIE.  
Muscidae, *Musca formosana* Malloch.
- sigmoidea* Geoffroy, 1785: 471 (*Musca*). France, Paris. T A Author. Acalyptrate.
- signata* Preyssler, 1791: 95, pl. 3, fig. 12 (*Musca*). Czechoslovakia, Prague, Botanical  
Garden. T ♀ DESTROY. Muscidae, *Helina celsa* Harris. See Rozkosny, Chvála  
& Pont 1982: 352.

- silentis** Harris, 1776: 59, pl. 15, fig. 14 (*Musca*). England. T ♂ Unknown. Syrphidae, *Sericomyia silentis* Harris.
- silvarum** Meigen, 1826: 53 (*Musca*). Austria. T ♀ NMW. Calliphoridae, *Lucilia silvarium* Meigen. Preocc. Linnaeus 1758 (*sylvarum*). Type lost (Rognes 1991a: 183).
- silvestris** Fallén, 1824: 70 (*Musca*). Sweden, Västergötland. LT ♀ NRS. Anthomyiidae, *Alliopsis silvestris* Fallén. Lectotype designated by Michelsen 1983: 115.
- similis** Forster, 1770: 15 (*Musca*). *Nomen nudum*. England. Stratiomyidae, *Beris similis* Forster. Published without diagnosis or indication.
- similis** Forster, 1771: 97 (*Musca*). England. T A LSL. Stratiomyidae, *Beris similis* Forster.
- simulator** Harris, 1780: 123, pl. 36, fig. 49 (*Musca*). England. T A Unknown. Acalyptrate.
- simulator** Harris, 1780: index (*Musca*). Incosp. *semulator* Harris. Syrphidae, *Xylota* sp. Name revised by Goffe 1946: 85.
- simulator** Harris, 1780: 152, pl. 45, fig. 77 (*Musca*). England. T A Unknown. Coelopidae.
- simulator** Harris, 1780: index (*Musca*). Incosp. *simulator* Harris. Acalyptrate. Name status by present revision.
- sinensis** Gmelin, 1790: 2872 (*Musca*). N. n. *Syrphus zonatus* Fabricius. Syrphidae, *Phytomia zonata* Fabricius.
- syngenesiae** Manuel, 1811: 37 (*Musca*). Emend. *syngenesiae* Fabricius. Platystomatidae, *Rivellia syngenesiae* Fabricius.
- singularius** Harris, 1776: 45, pl. 11, fig. 2 (*Musca*). England. T A Unknown. Stratiomyidae, *Stratiomys singularior* Harris.
- sinuata** Donovan, 1805: 88, pl. 41 (*Musca*). Australia. T A Author. Tachinidae, *Microtropesa sinuata* Donovan. See Townsend 1932: 40 (HT F in BMNH!), Crosskey 1973b: 135.
- sinuata** Wiedemann, 1830: 384 (*Musca*). Subsequent Usage of *Musca sinuata* Donovan. Tachinidae, *Microtropesa sinuata* Donovan.
- smaragdina** Geoffroy, 1785: 492 (*Musca*). France, Paris. T A Author.
- socia** Fallén, 1825: 82 (*Musca*). Sweden, Skåne. LT ♀ NRS. Anthomyiidae, *Emesomyia socia* Fallén. Lectotype designated by Michelsen 1983: 115.
- socio** Harris, 1780: 147, pl. 43, fig. 69 (*Musca*). England. T A Unknown. Fanniidae, *Fannia canicularis* Linnaeus. Name improperly formed (verb).
- solaia** Walker, 1849: 887 (*Musca*). Australia. T ♀ BMNH. Calliphoridae, *Hemipyrellia ligurriens* Wiedemann.
- solicitus** Harris, 1780: 116, pl. 34, fig. 18 (*Musca*). England. T A Unknown. Sciomyzidae, *Elgiva sollicita* Harris.
- solivagus** Harris, 1780: 85, pl. 25, fig. 15 (*Musca*). England. T A Unknown. Tachinidae, *Thelaira solivaga* Harris.
- solor** Harris, 1780: 146, pl. 43, fig. 65 (*Musca*). England. T A Unknown. Anthomyiidae. Name improperly formed (verb).
- solstitialis** Linnaeus, 1758a: 601 (*Musca*). Europe. T A LSL. Tephritidae, *Urophora solstitialis* Linnaeus. See White 1987: 105.
- solstitialis**, Sulzer, 1776: 216 (*Musca*). Switzerland. T A UNKNOWN. Preocc. Linnaeus 1758. *Subsequent usage*

- somalorum** Bezzı, 1892: 190 (*Musca*, as *corvina* var.). Somalia, Obbia, Uebi. T ♀ MCSNM. Muscidae, *Musca somalorum* Bezzı.
- sonchi** Linnaeus, 1767: 998 (*Musca*). Europe. T A LSL. Tephritidae, *Ensina sonchi* Linnaeus. See White 1987: 105.
- sorbens** Wiedemann, 1830: 418 (*Musca*). Sierra Leone. HT ♀ NMW. Muscidae, *Musca sorbens* Wiedemann. See Pont 1973: 289.
- sordida** Gmelin, 1790: 2861 (*Musca*). Europe. T A NMI.
- sordida** Zetterstedt, 1837: 42 (*Musca*). *Nomen nudum*. [Sweden, Lycksele]. T ♀ ZIL. Calliphoridae, *Protocalliphora rognesi* Thompson & Pont.
- sordida** Zetterstedt, 1838: 657 (*Musca*). Sweden, Lycksele. LT ♀ ZIL. Calliphoridae, *Protocalliphora rognesi* Thompson & Pont. Preocc. Gmelin 1790. Lectotype designated by Sabrosky 1956: 179. N. Syn.
- sordidissima** Walker, 1864: 216 (*Musca*). Mysol. HT ♂ BMNH. Muscidae, *Musca domestica* Linnaeus. See Pont 1990: 169.
- soror** Robineau-Desvoidy, 1830: 399 (*Musca*). India, Coromandel. T ♀ DESTROY. Muscidae, *Musca domestica* Linnaeus.
- spangleri** Zielke, 1970: 504 (*Musca*). *Nomen nudum*. Uganda, Kichwamba, Ankole. HT ♂ USNM. Muscidae, *Musca spangleri* Zielke. Published after 1930 without a description.
- spangleri** Zielke, 1971a: 103, fig. 1 (*Musca*). Uganda, Kichwamba, Ankole Distr. HT ♂ USNM. Muscidae, *Musca spangleri* Zielke.
- spatula** Scopoli, 1763: 341 (*Musca*). Idrıja. T A DESTROY. Stratiomyidae, *Stratiomys chamaeleon* Linnaeus.
- spectanda** Wiedemann, 1830: 419 (*Musca*). Sierra Leone. HT ♂ NMW. Muscidae, *Musca sorbens* Wiedemann. See Pont 1973: 290.
- speculifera** de Villers, 1789a: 464, pl. 9, fig. 25 (*Musca*). Europe. T A Author. Syrphidae, *Chrysogaster* sp.
- speculifera** Bezzı, 1911: 96 (*Musca*). Tunisia, Djerba Is. HT ♀ MCSNM. Muscidae, *Musca albina* Wiedemann. Preocc. de Villers 1789.
- sperata** Walker, 1859b: 136 (*Musca*). Sulawesi, Makassar. T ♂ BMNH. Muscidae, *Neomyia sperata* Walker.
- sphegea** Gmelin, 1790: 2873 (*Musca*). Subsequent Usage of *Syrphus sphegeus* Fabricius. Sciomyzidae, *Sepedon sphegeus* Fabricius.
- sphinx** Gmelin, 1790: 2832 (*Musca*). Subsequent Usage of *Bibio sphinx* Fabricius. Bombyliidae, *Ligyra sphinx* Fabricius.
- spilogaster** Wiedemann, 1824: 50 (*Musca*). South Africa, Cape of Good Hope. T ♂ UZMC. Sarcophagidae, *Poecilometopa spilogaster* Wiedemann.
- spinifera** Geoffroy, 1785: 494 (*Musca*). France, Paris. T A Author.
- spinipes** Scopoli, 1763: 342 (*Musca*). Carniola. T A DESTROY. Sciomyzidae, *Sepedon spinipes* Scopoli.
- spinipes** Fallén, 1823: 61 (*Musca*). Sweden. LT ♂ NRS. Muscidae, *Hydrotaea diabolus* Harris. Preocc. Scopoli 1763. Lectotype designated by Pont 1984: 293.
- spinohumera** Awati, 1916b: 138 (*Musca*). India. ST ♀ CRIK. Muscidae, *Musca lusoria* Wiedemann.
- spinosa** Awati, 1916b: 138 (*Musca*). India. ST ♀ CRIK. Muscidae, *Musca pattoni* Austen.
- spitzbergensis** Walker, 1849: 893 (*Musca*). Subsequent Usage of *Calliphora spitzbergenensis* Robineau-Desvoidy. Calliphoridae, *Calliphora vicina* Robineau-Desvoidy.

- splendens* Gmelin, 1790: 2833 (*Musca*). Subsequent Usage of *Stratiomys splendens* Fabricius. Stratiomyidae, *Pycnomalla splendens* Fabricius.
- splendens* Pont, 1980: 727 (*Musca*). N. n. *Musca splendida* Paterson. Muscidae, *Musca splendens* Pont.
- splendida* Donovan, 1805: 88, pl. 41 (*Musca*). Australia, New South Wales. NT ♂ BMNH. Tachinidae, *Rutilia splendida* Donovan. Crosskey 1973a: 123.
- splendida* Meigen, 1826: 56 (*Musca*). Germany, Stolberg. LT ♂ MHNTP. Calliphoridae, *Lucilia caesar* Linnaeus. Preocc. Donovan 1805. Lectotype designated by Rognes 1991a: 164.
- splendida* Paterson, 1957: 111, fig. 5 (*Musca*). South Africa, Cape, Knysna. HT ♂ SAIMR. Muscidae, *Musca splendens* Pont. Preocc. Donovan 1805.
- spoliata* Scopoli, 1763: 350 (*Musca*). Carniola. T A DESTROY. Scenopinidae, *Scenopinus fenestralis* Linnaeus.
- spondylii* Schrank, 1837: 18 (*Musca*). Austria? T A Unknown. Sepsidae, *Saltella sphondylii* Schrank. N. Syn.
- squalida* Gmelin, 1790: 2861 (*Musca*). Europe. T A NMI.
- squamata* Müller, 1764: 85 (*Musca*). Denmark, Sjaelland, Frederiksdal. T A Author. Dolichopodidae, *Dolichopus ungulata* Linnaeus. N. Syn.
- squamata* Malloch, 1934: 10 (*Viviparomusca*). Sulawesi, Menado. HT ♀ IRSNB. Muscidae, *Musca mallochi* Thompson & Pont. Preocc. Müller 1764.
- squamifera* Müller, 1776: 176 (*Musca*). N. n. *Musca squamata* Müller. Dolichopodidae, *Dolichopus ungulatus* Linnaeus.
- stabulans* Fallén, 1817: 252 (*Musca*). Sweden. LT ♂ NRS. Muscidae, *Muscina stabulans* Fallén. Lectotype designated by Pont 1984: 293.
- stellata* Fuesslin, 1775: 56 (*Musca*). Switzerland, Zürich. T A Unknown. Tephritidae, *Trupanea stellata* Fuesslin.
- stellata* Geoffroy, 1785: 470 (*Musca*). France, Paris. T A Author. Preocc. Fuesslin 1775.
- stellata* Roemer, 1789: 84 (*Musca*). Switzerland. T A Unknown. Tephritidae, *Trupanea stellata* Fuesslin. Preocc. Fuesslin 1775.
- stercoraria* Linnaeus, 1758a: 599 (*Musca*). Europe. T A LSL. Scathophagidae, *Scathophaga stercoraria* Linnaeus. 2 sp. in LSL (Dear MS).
- stictica* Gmelin, 1790: 2863 (*Musca*). N. n. *Musca punctata* Schrank. Tephritidae, *Orellia stictica* Gmelin. N. Comb.
- stigma* Fabricius, 1798: 563 (*Musca*). West Indies. T A UZMC. Otitidae, *Noto-gramma stigma* Fabricius.
- stigma* Panzer, 1798c: 21 (*Musca*). Austria. T A ZMHU. Sepsidae, *Sepsis punctum* Fabricius.
- stimulans* Robineau-Desvoidy, 1830: 393 (*Byomyia*). France, Saint-Sauveur. T A DESTROY. Muscidae, *Musca tempestiva* Fallén.
- stimulans* Macquart, 1835: 268 (*Musca*). Subsequent Usage of *Byomyia stimulans* Robineau-Desvoidy. Muscidae, *Musca tempestiva* Fallén.
- stipata* Walker, 1853b: 348 (*Musca*). Guyana, Demerara. T ♀ BMNH. Muscidae, *Graphomya stipata* Walker.
- stomoxidea* Robineau-Desvoidy, 1830: 396 (*Musca*). Brazil. T ♀ DESTROY. Muscidae, *Musca domestica* Linnaeus.
- striata* Gmelin, 1790: 2865 (*Musca*). N. n. *Tipula lineata* Fabricius. Keroplatidae, *Cerotelion striatum* Gmelin. N. Status

- striata* Fabricius, 1794: 315 (*Musca*). Denmark, Copenhagen. LT ♀ UZMC. Sarco-phagidae, *Ravinia pernix* Harris. Preocc. Gmelin 1790. Lectotype designated by Pape 1986: 303 (also Townsend 1931c: 377).
- striata* Townsend, 1931c: 378 (*Musca*). Subsequent Usage of *Musca striata* of Enderlein, not Fabricius. Sarcophagidae, *Thrysocnema incisilobata* Pandellé. Renamed as *Thrysocnema transita* based on HT M in ZMHU.
- striacta** Awati, 1916b: 138 (*Musca*). India. HT A CRIK. Muscidae, *Musca striacta* Awati.
- strigata* Gmelin, 1790: 2834 (*Musca*). Subsequent Usage of *Stratiomys strigata* Fabricius. Stratiomyidae, *Stratiomys longicornis* Scopoli.
- strigosa* Gmelin, 1790: 2843 (*Musca*). Europe. T A NMI.
- strigosa* Fabricius, 1794: 322 (*Musca*). Germany, Kiel. HT A UZMC F. Anthomyiidae, *Hylemya vagans* Panzer. Preocc. Gmelin 1790. See Michelsen 1979a: 189.
- strigula* Fabricius, 1794: 334 (*Musca*). France. T A MNHNP. Chloropidae, *Chlorops strigula* Fabricius.
- striolata* Fallén, 1824: 71 (*Musca*). Sweden, Skåne. LT ♂ NRS. Anthomyiidae, *Botanophila striolata* Fallén. Lectotype designated by Michelsen 1983: 118.
- structus* Harris, 1780: 147, pl. 43, fig. 68 (*Musca*). England. T A Unknown. Anthomyiidae.
- strumosa* Scopoli, 1763: 343 (*Musca*). Carniola. T A DESTROY. Micropezidae, *Micropeza corrigiolata* Linnaeus.
- stuans* Manuel, 1811: 35 (*Musca*). Missp. *aestuans* Linnaeus.
- stuckenbergi* Zielke, 1971c: 100 (*Musca*). South Africa, Natal, Pietermaritzburg. HT ♀ NMP. Muscidae, *Musca sorbens* Wiedemann.
- stygia* Fabricius, 1781: 438 (*Musca*). (Newfoundland) Error for New Zealand. HT ♀ BMNH B. Calliphoridae, *Calliphora stygia* Fabricius. See Osten Sacken 1882: 20; Townsend 1931c: 374; Dear 1985a: 24.
- stylata* Fabricius, 1775: 785 (*Musca*). England. T A UZMC F. Tephritidae, *Urophora stylata* Fabricius.
- subcoleoptrata* Gmelin, 1790: 2869 (*Musca*). Subsequent Usage of *Conops subcoleoptrata* Linnaeus. Tachinidae, *Allophora subcoleoptrata* Linnaeus.
- subcutanea* Linnaeus, 1764a: 297 (*Musca*). *Nomen nudum*. [France]. Tephritidae. Published in non-binominal work.
- subcutanea* Linnaeus, 1764a: 300 (*Musca*). *Nomen nudum*. [France]. Tephritidae, *Ensina sonchi* Linnaeus. Published in non-binominal work.
- subcutanea* Linnaeus, 1764a: 287 (*Musca*). *Nomen nudum*. [France]. Tephritidae. Published in non-binominal work.
- subcutanea* Bjerkander, 1793: 57 (*Musca*). Sweden. T A UNKNOWN.
- subcutanea* Turton, 1801: 623 (*Musca*). N. n. *Musca heraclii* Linnaeus. Tephritidae, *Euleia heraclii* Linnaeus. Preocc. Bjerkander 1793.
- sublatus* Harris, 1780: 125, pl. 36, fig. 56 (*Musca*). England. T ♂ Unknown. Muscidae, *Phaonia* sp.
- subpuncta* Fallén, 1825: 80 (*Musca*). Sweden. LT ♂ NRS. Muscidae, *Helina quadrum* Fabricius. Lectotype designated by Pont 1984: 294.
- subsultans** Linnaeus, 1767: 993 (*Musca*). Sweden. T A LSL. Phoridae, *Borophaga subsultans* Linnaeus. See Disney 1982: 115.
- subtomentosa* Gmelin, 1790: 2843 (*Musca*). Europe. T A NMI. Syrphidae, *Sericomyia silentis* Harris.

- subulata** Gmelin, 1790: 2860 (*Musca*). Europe. T A NMI.
- subvariegata** Jördens, 1801: 301 (*Musca*). *Nomen nudum*. Name published in synonymy of *Musca carnaria* and attributed to "Scopoli Ent. carn. 869" [= *variegata*].
- subventus** Harris, 1780: 145, pl. 43, fig. 60 (*Musca*). England. T ♂ Unknown. Muscidae, *Phaonia subventa* Harris.
- succincta** Gmelin, 1790: 2866 (*Musca*). N. n. *Rhagio cinctus* Fabricius. Anisopodidae, *Sylvicola cinctus* Fabricius.
- suecica** de Villers, 1789a: 510 (*Musca*). Sweden, Ostrogothia. T A Unknown.
- suffusa** Walker, 1853b: 336 (*Musca*). Brazil. T A DESTROY. Calliphoridae.
- sugillatrix** Robineau-Desvoidy, 1830: 392 (*Plaxemya*). France, Saint-Sauveur. T ♂ DESTROY. Muscidae, *Musca osiris* Wiedemann.
- suilla** Fabricius, 1794: 343 (*Musca*). Germany. T A UZMC F. Scathophagidae, *Scathophaga suilla* Fabricius.
- surinamensis** De Geer, 1776: 145, pl. 29, fig. 1 (*Musca*). Surinam. T A NRS. Syrphidae, *Palpada vinetorum* Fabricius. See Thompson 1973: 354.
- sylvanus** Gmelin, 1790: 2832 (*Musca*). Subsequent Usage of *Bibio sylvana* Fabricius. Bombyliidae, *Comptosia sylvana* Fabricius.
- sylvarum** Linnaeus, 1758a: 592 (*Musca*). Sweden. LT ♀ LSL. Syrphidae, *Xylota sylvarum* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 160.
- syngenesiae** Fabricius, 1781: 454 (*Musca*). Germany. T A UZMCF. Platystomatidae, *Rivellia syngenesiae* Fabricius.
- syrphus** Meuschen, 1787: 212 (*Musca*). Unknown. T A Auction.
- tabaniformis** Fabricius, 1805: 302 (*Musca*). Mogador [=Morocco, Essaouira]. HT ♀ UZMC. Sarcophagidae, *Craticulina tabaniformis* Fabricius. See Townsend 1931c: 379; Pape 1986: 303.
- tachinoides** Fallén, 1817: 244 (*Musca*). Sweden, Östergötland & Västergötland. ST A NRS. Tachinidae, *Stomina tachinoides* Fallén. See Townsend 1932: 43 (invalid HT designation).
- taeniata** Panzer, 1798c: 18 (*Musca*). Germany, Nuremberg. T ♂ ZMHU. Tachinidae, *Phasia crassipennis* Fabricius.
- taenionota** Wiedemann, 1819a: 22 (*Musca*). Java. T ♀ UZMC. Sarcophagidae.
- tahitiensis** Lichtenstein, 1796: 215 (*Musca*). Tahiti. T A Auction. Muscidae.
- taitensis** Macquart, 1843: 310, pl. 20, fig. 8 (*Musca*). Tahiti. T ♂ MNHNP. Muscidae, *Musca domestica* Linnaeus.
- taniaria** Wiedemann, 1830: 406 (*Musca*). Incosp. *Ianiaria* Wiedemann. Calliphoridae, *Cochliomyia macellaria* Fabricius. Name revised by Wiedemann 1830: 683. See Dear 1985b: 141.
- tarda** Linnaeus, 1761: 441 (*Musca*). Sweden. T A LSL. Scenopinidae, *Scenopinus fenestralis* Linnaeus. 4 sp. in LSL (Dear MS).
- tardigradus** Harris, 1776: 46, pl. 11, fig. 6 (*Musca*). England. T A Unknown. Stratiomyidae, *Oxycrera rara* Scopoli.
- tarditas** Harris, 1780: 106, pl. 32, fig. 40 (*Musca*). England. T A Unknown. Syrphidae, *Cheilosia* sp.
- tarditus** Harris, 1780: index (*Musca*). Incosp. *tarditas* Harris. Syrphidae, *Cheilosia* sp. Name revised by Goffe 1946: 85.
- tau** Schrank, 1781: 458 (*Musca*). Austria, Linz. T A Unknown. Muscidae, *Musca autumnalis* De Geer.

- tegularia* Wiedemann, 1830: 655 (*Musca*). Egypt. T A SMF. Calliphoridae, *Lucilia sericata* Meigen.
- temperata* Walker, 1853b: 340 (*Musca*). East Indies. T ♀ BMNH. Calliphoridae, *Lucilia cuprina* Wiedemann.
- tempestatum* Bezzi, 1908c: 101 (*Biomyia*). Ethiopia (Eritrea), Asmara. ST ♂ ♀ MCSNM. Muscidae, *Musca tempestatum* Bezzi. STs also in BMNH (Pont 1990: 169) & USNM (*teste* Pont).
- tempestiva* Fallén, 1817: 254 (*Musca*). Sweden, Skåne, near Kivik. LT ♂ NRS. Muscidae, *Musca tempestiva* Fallén. Lectotype designated by Pont 1984: 294.
- temula* Scopoli, 1763: 330 (*Musca*). Carniola. T A DESTROY. Tachinidae, *Zophomyia temula* Scopoli.
- tenax* Linnaeus, 1758a: 591 (*Musca*). Sweden. LT ♀ LSL. Syrphidae, *Eristalis tenax* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 160.
- tenebricus* Harris, 1776: 45, pl. 11, fig. 3 (*Musca*). England. T A Unknown. Stratiomyidae, *Stratiomys longicornis* Scopoli.
- tener* Harris, 1780: 116, pl. 34, fig. 19 (*Musca*). England. T A Unknown. Sciomyzidae.
- tentaculata* De Geer, 1776: 86 (*Musca*). Sweden. ST A Lost. Muscidae, *Lispe tentaculata* De Geer. See Persson, Pont & Michelsen 1984: 93.
- tenur* Harris, 1780: 112, pl. 33, fig. 59 (*Musca*). England. T A Unknown. Syrphidae, *Neoascia tenur* Harris. See Thompson 1981b: 476.
- tenur* Harris, 1780: index (*Musca*). Incosp. *tener* Harris. Sciomyzidae. Name status by present revision.
- teres* Gmelin, 1790: 2861 (*Musca*). Europe. T A UNKNOWN.
- terminata* Wiedemann, 1830: 414 (*Musca*). Sierra Leone. T ♂ NMW. Calliphoridae, *Isomyia terminata* Wiedemann.
- terraereginae* Johnston & Bancroft, 1920b: 20 (*Musca*). *Nomen nudum*. Australia, Queensland. Muscidae, *Musca terraereginae* Johnston & Bancroft. [also 1920c: 34, 1920d: 73].
- terraereginae* Johnston & Bancroft, 1920e: 31 (*Musca*). Australia, Queensland, Eidsvold. LT ♂ QMB. Muscidae, *Musca terraereginae* Johnston & Bancroft. Lectotype designated by Pont 1973: 180.
- terrestris* Linnaeus, 1767: 996 (*Musca*). Sweden. T A LSL.
- tessella* Turton, 1801: 604 (*Musca*). Emend. *Musca tessellata* Fabricius. Tachinidae, *Tachina tessella* Turton. N. Status
- tessellata* Geoffroy, 1785: 476 (*Musca*). France, Paris. T A Author.
- tessellata* Fabricius, 1794: 324 (*Musca*). Germany. T A UZMC F. Tachinidae, *Tachina tessella* Turton. Preocc. Geoffroy 1785. N. Syn.
- tessellata* Fabricius, 1805: 285 (*Musca*). South America. HT ♀ UZMC. Sarcophagidae, *Euboettcheria australis* Townsend. Preocc. Geoffroy 1785. See Townsend 1931a: 74; Pape 1986: 303.
- testacea* De Geer, 1776: 87 (*Musca*). Sweden. LT ♂ NRS. Anthomyiidae, *Pegomya testacea* De Geer. Lectotype designated by Persson, Pont & Michelsen 1984: 94.
- testacea* Fabricius, 1781: 440 (*Musca*). Germany. T A DESTROY. Muscidae, *Phaonia rufiventris* Scopoli. Preocc. De Geer 1776. See Michelsen 1979a: 190.
- testacea* Fabricius, 1787: 330 (*Stratiomys*). South America. T A UZMCF. Xylophagidae, *Coenomyia ferruginea* Scopoli.

- testaceus** Fabricius, 1805: 257 (*Sargus*). N. n. *Musca elongata* Fabricius. Stratiomyidae, *Ptecticus testaceus* Fabricius.
- teste** Zielke, 1971c: 126 (*Musca*). *Nomen nudum*. Muscidae, *Musca lusoria* Wiedemann. Published in synonymy, not subsequently validated by usage. Mistake for *Musca lusoria* Wiedemann *teste* Bezz!
- tetragona** Geoffroy, 1785: 484 (*Musca*). France, Paris. T A Author. Syrphidae.
- thalassina** Meigen, 1826: 54 (*Musca*). Germany, Stolberg. HT ♀ MNHNP. Calliphoridae, *Phormia regina* Meigen. See Rognes 1991a: 106.
- thuscia** Walker, 1849: 897 (*Musca*). Azores, Fayal. T ♀ BMNH. Calliphoridae, *Calliphora vicina* Robineau-Desvoidy.
- thymastri** Gmelin, 1790: 2876 (*Musca*). Subsequent Usage of *Syrphus thymastri* Fabricius. Syrphidae, *Paragus* sp.
- tiberina** Saccà, 1947: 128 (*Musca*, as *domestica* var.). *Nomen nudum*. T A UNKNOWN. Muscidae, *Musca domestica* Linnaeus.
- tibetana** Fan, 1978: 329, fig. 1 (*Musca*). Tibet, Nyingchi (*teste* Fan). T ♂ SIE. Muscidae, *Musca tibetana* Fan.
- tibialis** Gmelin, 1790: 2859 (*Musca*). Europe. T A NMI.
- tibialis** Walker, 1836: 355 (*Musca*). Brazil, Santa Catarina. T A BMNH. Muscidae, *Muscina stabulans* Fallén. Preocc. Gmelin 1790.
- tibiseta** Malloch, 1928: 334 (*Byomyia*). Sumatra, Fort de Kock. HT ♂ Lost. Muscidae, *Musca seniorwhitei* Patton. See Pont 1970a: 105.
- tifata** Walker, 1849: 871 (*Musca*). China. T ♂ BMNH. Calliphoridae, *Lucilia papuensis* Macquart.
- tigrina** Fabricius, 1775: 779 (*Musca*). England. T A DESTROY. Muscidae, *Coenosia tigrina* Fabricius. See Michelsen 1979a: 190.
- timeo** Harris, 1780: 107, pl. 32, fig. 44 (*Musca*). England. T A Unknown. Syrphidae, *Platycheirus peltatus* Meigen. Name improperly formed (verb).
- timidus** Harris, 1780: 116, pl. 34, fig. 17 (*Musca*). England. T A Unknown. Acalyptrate.
- tipulaea** Geoffroy, 1785: 497 (*Musca*). France, Paris. T A Author.
- tipularia** Fabricius, 1794: 337 (*Musca*). Barbaria. T A MHN. Scathophagidae.
- tipuliformis** Gmelin, 1790: 2865 (*Musca*). Subsequent Usage of *Rhagio tipuliformis* Fabricius.
- tomentosa** Turton, 1801: 634 (*Musca*). Subsequent Usage of *Rhagio tomentosus* Fabricius. Rhagionidae, *Chrysopilus auratus* Fabricius.
- topiaria** Schrank, 1803: 126 (*Musca*). Germany, Bavaria, Ingolstadt. T A UNKNOWN.
- torosa** Wiedemann, 1819a: 21 (*Musca*). Bengal. T ♀ UZMC. Calliphoridae, *Bengalia torosa* Wiedemann.
- torvis** Harris, 1780: index (*Musca*). Incosp. *torvus* Harris. Tachinidae, *Ernestia rufis?* Fallén. Name status by present revision.
- torvus** Harris, 1780: 113, pl. 33, fig. 26 (*Musca*). England. T A Unknown. Tachinidae, *Ernestia rufis?* Fallén.
- totliana** Gmelin, 1790: 2850 (*Musca*). N. n. *Musca fasciata* Fabricius. Micropezidae, *Hoplocheiloma totliana* Gmelin. N. Comb.
- tranquebarica** Gmelin, 1790: 2870 (*Musca*). N. n. *Syrphus quadrilineatus* Fabricius. Syrphidae, *Eristalinus arvorum* Fabricius.

- transfuga** Linnaeus, 1758a: 594 (*Musca*). Sweden. LT ♀ LSL. Syrphidae, *Lejops transfugus* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 160.
- transvaalensis** Zielke, 1971c: 120 (*Musca*). South Africa, Transvaal, Bloemhof. HT ♀ NMP. Muscidae, *Musca transvaalensis* Zielke.
- transversa** Fallén, 1825: 92 (*Musca*). Sweden, Skåne. LT ♂ NRS. Anthomyiidae, *Pegomya transversa* Fallén. Lectotype designated by Michelsen 1983: 118.
- trelineata** Harris, 1782: 58 (*Musca*). Missp. *trilenva* Harris. Syrphidae, *Helophilus pendulus* Linnaeus.
- tremula** Linnaeus, 1767: 991 (*Musca*). Emend. *temula* Scopoli. Tachinidae, *Zophomyia temula* Scopoli.
- triangula** Fallén, 1825: 74 (*Musca*). Sweden, Skåne, Äsperöd. LT ♂ NRS. Muscidae, *Limnophora triangula* Fallén. Lectotype designated by Pont 1984: 294.
- tricincta** Fabricius, 1805: 301 (*Musca*). South America. HT ♀ UZMC. Tachinidae, *Comatacta tricincta* Fabricius. See Townsend 1931a: 91.
- tricolor** Gmelin, 1790: 2861 (*Musca*). Europe. T A NMI.
- tridens** Bosc, 1792: 54 (*Musca*). France, Paris. T A MNHNP.
- trifascia** Walker, 1861b: 245 (*Musca*). New Guinea, Dorey. T ♀ BMNH. Calliphoridae, *Dexopollenia trifascia* Walker.
- trifasciata** Schrank, 1781: 453 (*Musca*). Subsequent Usage of *Conops trifasciata* Scopoli. Syrphidae, *Volucella inanis* Linnaeus.
- trigrammica** Schrank, 1803: 125 (*Musca*). Germany, Bavaria, Ingolstadt. T A Unknown.
- trileniata** Harris, 1780: index (*Musca*). Missp. *trilenva* Harris. Syrphidae, *Helophilus pendulus* Linnaeus.
- trilrena** Harris, 1776b: 58, pl. 15, fig. 9 (*Musca*). England. T A Unknown. Syrphidae, *Helophilus pendulus* Linnaeus.
- trilineata** Linnaeus, 1767: 980 (*Musca*). Sweden. T A LSL. Stratiomyidae, *Oxycera trilineata* Linnaeus. 1 sp. in LSL (Dear MS).
- trilineata** Harris, 1776b: 58 (*Musca*). Incosp. *trilenva* Harris. Syrphidae, *Helophilus pendulus* Linnaeus. Name status by present revision.
- trilineatus** Fabricius, 1775: 766 (*Syrphus*). New Zealand. T A BMNH B. Syrphidae, *Helophilus seelandicus* Gmelin.
- trimaculata** Geoffroy, 1785: 472 (*Musca*). France, Paris. T A Author.
- trinervis** Geoffroy, 1785: 489 (*Musca*). France, Paris. T A Author.
- tringaria** Linnaeus, 1758a: 590 (*Musca*). Sweden. T A LSL. Rhagionidae, *Rhagio tringarius* Linnaeus. 1 sp. in LSL (Dear MS).
- tripunctata** Scopoli, 1763: 329 (*Musca*). Idrija. T A DESTROY.
- tripunctata** Bjerkander, 1777: 33 (*Musca*). Sweden. T A UNKNOWN. Preocc. Scopoli 1763.
- tripunctata** Gmelin, 1790: 2861 (*Musca*). Europe. T A NMI. Preocc. Scopoli 1763.
- triquetra** Fallén, 1825: 73 (*Musca*). Subsequent Usage of *Anthomyia triquetra* Wiedemann. Muscidae, *Azelia triquetra* Wiedemann.
- tristis** Sulzer, 1776: 216 (*Musca*). Switzerland, Zürich. T A Unknown.
- tristis** Harris, 1780: 115, pl. 34, fig. 13 (*Musca*). England. T A Unknown. Sciomyzidae, *Coremacera marginata* Fabricius. Preocc. Sulzer 1776.
- tristis** Herbst, 1787: 110 (*Musca*). T A UNKNOWN. Tachinidae, *Allophora hemiptera* Fabricius. Preocc. Sulzer 1776.

- tristis* Schrank, 1803: 110 (*Musca*). Germany. Gern. T A Unknown. Syrphidae. Preocc. Sulzer 1776.
- tristor* Harris, 1780: 107, pl. 32, fig. 42 (*Musca*). England. T A Unknown. Syrphidae, *Pipiza noctiluca* Linnaeus. Name improperly formed (verb).
- trita* Walker, 1856a: 24 (*Musca*). Malacca. T ♀ BMNH. Muscidae, *Neomyia coeruleifrons* Macquart.
- troglodyta* Gmelin, 1790: 2832 (*Musca*). Subsequent Usage of *Bibio troglodyta* Fabricius. Bombyliidae, *Villa troglodyta* Fabricius.
- truncata* Bjerkander, 1777: 34 (*Musca*). Sweden. T A UNKNOWN.
- tuberculata* Müller, 1776: 177 (*Musca*). Denmark. T A Author.
- tuberculata* de Villers, 1789a: 428, pl. 9, fig. 12 (*Musca*). Southern France, St. Jean de Gardonенque. T A Author. Bibionidae, *Dilophus novemmaculatus* de Villers. Preocc. Müller 1776.
- tuberculata* de Villers, 1789a: 550 (*Musca*). France, Cevennes Mts. T A Author. Preocc. Müller 1776.
- tuberis* Vallot, 1802: 131 (*Musca*). France. T A UNKNOWN. Heleomyzidae, *Suillia* sp. N. Comb.
- tuguriorum* Scopoli, 1763: 328 (*Musca*). Carniola. T A DESTROY. Muscidae, *Phaonia tuguriorum* Scopoli.
- turbida* Wiedemann, 1830: 396 (*Musca*). South Africa, Cape of Good Hope. T ♀ ZMHU. Calliphoridae.
- turbida* Walker, 1853b: 336 (*Musca*). Jamaica. T A DESTROY. Calliphoridae, *Cochliomyia macellaria* Fabricius. Preocc. Wiedemann 1830. [teste Pont].
- turcica* Geoffroy, 1785: 487 (*Musca*). France, Paris. T A Author.
- tussilaginis* Fabricius, 1775: 787 (*Musca*). Denmark. T A UZMC F. Tephritidae, *Terellia tussilaginis* Fabricius.
- ugandae* Emden, 1939: 84 (*Musca*, as *autumnalis* ssp.). Uganda, Burunga. LT ♂ BMNH. Muscidae, *Musca ugandae* Emden. Lectotype designated by Pont 1990: 169.
- uliginosa* Linnaeus, 1767: 983 (*Musca*). Europe. T A LSL. Stratiomyidae, *Nemotelus uliginosus* Linnaeus. 1 sp. in LSL (Dear MS).
- uliginosa* Fallén, 1825: 81 (*Musca*). Sweden. LT ♂ NRS. Muscidae, *Helina sexmaculata* Preissler. Preocc. Linnaeus 1758. Lectotype designated by Pont 1984: 295.
- umbellatarum* Fabricius, 1775: 785 (*Musca*). England. T A UZMC F. Pallopteridae, *Palloptera umbellatarum* Fabricius.
- umbelliferarum* Scopoli, 1763: 349 (*Musca*). Carniola. T A DESTROY. Chloropidae, *Chlorops pumilionis* Bjerkander.
- umbraculata* Fabricius, 1794: 348 (*Musca*). France. T A MNHNP. Chloropidae, *Platycephala umbraculata* Fabricius.
- umbraculata* Fabricius, 1805: 294 (*Musca*). Denmark, Copenhagen. HT ♂ UZMC. Muscidae, *Musca domestica* Linnaeus. Preocc. Fabricius 1794. See Michelsen 1979a: 190.
- umbrarum* Linnaeus, 1758a: 599 (*Musca*). Europe. T A LSL. Sciomyzidae, *Dictya umbrarum* Linnaeus. 3 sp. in LSL (Dear MS).
- umbrifera* Walker, 1861c: 267 (*Musca*). Sulawesi, Tondano. HT ♂ BMNH. Calliphoridae, *Pollenia umbrifera* Walker. [teste Pont]. N. Comb.
- unguicornis* Scopoli, 1763: 335 (*Musca*). Carniola. T A DESTROY. Sciomyzidae, *Limnia unguicornis* Scopoli.

- ungulata** Linnaeus, 1758a: 598 (*Musca*). Europe. T A LSL. Dolichopodidae, *Dolichopus ungulatus* Linnaeus. 4 sp. in LSL (Dear MS).
- unicolor** Fabricius, 1805: 307 (*Musca*). Germany, Kiel. T A UZMC F. Pallopteridae, *Palloptera saltuum* Linnaeus.
- unisetula** Fan, 1965: 122 (*Musca*, as *tempestiva* var.). *Nomen nudum*. Korea, Kaicheng. HT ♂ SIE. Muscidae, *Musca tempestiva* Fallén. Varietal name proposed after 1960.
- univalvis** Turton, 1801: 632 (*Musca*). N. n. *Musca clavicornis* Fabricius. Syrphidae, *Ceriana clavicornis* Fabricius.
- unxia** Walker, 1849: 899 (*Musca*). Italy. LT ♂ BMNH. Calliphoridae, *Bellardia pandia* Walker. Lectotype designated by Schumann 1974: 253.
- urbanus** Harris, 1780: 153, pl. 45, fig. 85 (*Musca*). England. T A Unknown. Tachinidae, *Siphona geniculata* De Geer.
- urticæ** Linnaeus, 1758a: 600 (*Musca*). Europe. T A LSL. Otitidae, *Ceroxyx urticæ* Linnaeus. 2 sp. in LSL (Dear MS).
- urticae**, Fabricius, 1794: 357 (*Musca*). Misid. *urticæ* Linnaeus. Otitidae, *Seioptera vibrans* Linnaeus.
- vaccarum** De Geer, 1776: 85, pl. 5, fig. 1 (*Musca*). N. n. *Musca meteorica* Linnaeus. Muscidae, *Hydrotaea meteorica* Linnaeus.
- vaccina** Robineau-Desvoidy, 1863: 625 (*Musca*). France, Paris. T A DESTROY. Muscidae, *Musca domestica* Linnaeus.
- vacua** Scopoli, 1763: 346 (*Musca*). Carniola. T A DESTROY. Syrphidae, *Syrphus ribesii* Linnaeus.
- vacua** Fallén, 1817: 240 (*Musca*). Sweden. T A NRS. Tachinidae, *Dexia cincta* Robineau-Desvoidy. Preocc. Scopoli 1763. N. Comb.
- vagabunda** Meigen, 1826: 72 (*Musca*). Germany, Stolberg. LT ♀ MHNTP. Calliphoridae, *Pollenia vagabunda* Meigen. Lectotype designated by Rognes 1991aa: 238 & 1992: 109.
- vagans** Panzer, 1798b: 18 (*Musca*). Austria. T ♀ ZMHU. Anthomyiidae, *Hylemya vagans* Panzer.
- vagans** Fallén, 1825: 78 (*Musca*). Sweden. LT ♂ NRS. Muscidae, *Phaonia falleni* Michelsen. Preocc. Panzer 1798. Lectotype designated by Pont 1984: 295.
- vagatoria** Robineau-Desvoidy, 1830: 399 (*Musca*). France. T ♀ DESTROY. Muscidae, *Musca domestica* Linnaeus.
- vagus** Harris, 1780: 153, pl. 45, fig. 82 (*Musca*). England. T A Unknown. Psilidae, *Loxocera* sp.
- valens** Harris, 1780: 113, pl. 33, fig. 25 (*Musca*). England. T A Unknown. Sarcophagidae, *Sarcophaga* sp.
- valentina** Müller, 1766: 198 (*Musca*). Italy, Piedmont, Torino. T A DESTROY. Syrphidae, *Volucella zonaria* Poda.
- valga** Gmelin, 1790: 2879 (*Musca*). Europe. T A NMI. Syrphidae, *Chalcosyrphus valgus* Gmelin.
- valida** Wiedemann, 1830: 387 (*Musca*). Unknown (=South Carolina?). HT ♀ NMW. Tachinidae, *Theresa rutilans* Fabricius. Preocc. Harris 1780. See Townsend 1931a: 102.
- validus** Harris, 1780: 124, pl. 36, fig. 50 (*Musca*). England. T ♀ Unknown. Muscidae, *Phaonia valida* Harris.

- vallata Forster, 1770: 15 (*Musca*). *Nomen nudum*. England. T A LSL. Stratiomyidae, *Beris vallata* Forster.
- vallata Forster, 1771: 96 (*Musca*). England. T A LSL. Stratiomyidae, *Beris vallata* Forster.
- vanella Turton, 1801: 632 (*Musca*). Subsequent Usage of *Rhagio vanellus* Fabricius. Rhagionidae, *Rhagio tringaria* Linnaeus.
- varensis Robineau-Desvoidy, 1863: 627 (*Musca*). France, Var. T ♀ DESTROY. Calliphoridae, *Pollenia* sp.
- varia Gmelin, 1790: 2843 (*Musca*). Europe. T A NMI.
- varia Fabricius, 1794: 327 (*Musca*). India Orientali. HT A UZMC. Tachinidae, *Cuphocera iavana* Wiedemann. Preocc. Gmelin 1790. See Townsend 1932: 42; Crosskey 1973b: 134. N. Comb.
- varia Meigen, 1826: 66 (*Musca*). Germany, Stolberg. LT ♂ MNHNP. Calliphoridae, *Pollenia rufis* Fabricius. Preocc. Gmelin 1790. Lectotype designated by Rognes 1991a: 238. N. Comb.
- varia Walker, 1853b: 342 (*Musca*). South Australia. HT ♀ BMNH. Calliphoridae, *Amenia chrysane* Walker. Preocc. Gmelin 1790. See Crosskey 1965: 120.
- variabilis Gmelin, 1790: 2834 (*Musca*). N. n. *Stratiomys mutabilis* Fabricius. Stratiomyidae, *Hoplitimya mutabilis* Fabricius.
- variabilis Fallén, 1823: 58 (*Musca*). Sweden. LT ♂ NRS. Muscidae, *Thricops nigrifrons* Robineau-Desvoidy. Preocc. Gmelin 1790. Lectotype designated by Pont 1984: 295.
- varians Wiedemann, 1830: 655 (*Musca*). Egypt. T ♀ SMF. Calliphoridae, *Lucilia cuprina* Wiedemann.
- varians Meigen, 1975: 418 (*Musca*). *Nomén nudum*. [Germany?]. T A MNHNP. Calliphoridae, *Pollenia vagabunda* Meigen. See Rognes 1992: 109.
- variata Fallén, 1823: 59 (*Musca*). Sweden, Skåne. LT ♂ NRS. Anthomyiidae, *Hylemya variata* Fallén. Lectotype designated by Hackett 1968: 346.
- varicolor Fabricius, 1805: 296 (*Musca*). India, Tranquebar. HT ♂ UZMC. Calliphoridae, *Bengalia varicolor* Fabricius. See Townsend 1931c: 371.
- varicus Harris, 1780: 114, pl. 34, fig. 12 (*Musca*). England. T A Unknown. Sciomyzidae.
- varicus Harris, 1780: 115, pl. 34, fig. 14 (*Musca*). England, Kent. T A Unknown. Sciomyzidae, *Limnia unguicornis* Scopoli. Preocc. Harris 1780.
- variegata Scopoli, 1763: 326 (*Musca*). Idrija. T A DESTROY. Sarcophagidae, *Sarcophaga variegata* Scopoli.
- variegata De Geer, 1776: 96, pl. 6, fig. 1 (*Musca*). [Sweden]. T A NRS. Sciomyzidae, *Euthycera chaerophylli* Fabricius. Preocc. Scopoli 1763.
- varieus Harris, 1780: index (*Musca*). Incosp. varicus Harris. Sciomyzidae, *Limnia unguicornis* Scopoli. Name status by present revision.
- variolosa Schrank, 1837: 18 (*Musca*). Germany, Bavaria. Platystomatidae, *Platystoma seminationis* Fabricius.
- vecors Harris, 1780: 154, pl. 45, fig. 89 (*Musca*). England. T A Unknown. Tephritidae.
- velox Bjerckander, 1777: 33 (*Musca*). Sweden. T A UNKNOWN.
- venatoria Fabricius, 1805: 285 (*Musca*). South America. HT ♂ UZMC. Tachinidae, *Scotiaptera venatoria* Fabricius. See Townsend 1931a: 100.
- venosa Presl, 1822: 203 (*Musca*). Ligurian Prussian Amber. T A Unknown.

- ventito** Harris, 1780: 139, pl. 41, fig. 30 (*Musca*). England. T A Unknown. Calliphoridae. Name improperly formed (verb).
- ventrosa** Wiedemann, 1830: 656 (*Musca*). China. LT ♂ NMW. Muscidae, *Musca ventrosa* Wiedemann. Lectotype designated by Pont 1973: 164.
- verena** Walker, 1849: 874 (*Musca*). Venezuela. T ♀ BMNH. Calliphoridae, *Compsomyiops verena* Walker. N. Comb.
- vermilio** Linnaeus, 1758a: 590 (*Musca*). France. T A LSL. Vermileonidae, *Vermileo vermilio* Linnaeus.
- verna** Fabricius, 1794: 330 (*Musca*). France. T A Lost. Muscidae, *Lispocephala verna* Fabricius. See Michelsen 1979a: 190; type figured by Coquebert 1804: 106.
- vernicina** Scopoli, 1763: 333 (*Musca*). Upper Carniola. T A DESTROY. Lonchaeidae, *Lonchaea* sp.
- versicolor** Gmelin, 1790: 2859 (*Musca*). Europe. T A NMI.
- versicolor** Fabricius, 1794: 283 (*Syrphus*). Germany. T A UZMC F. Syrphidae, *Parhelophilus versicolor* Fabricius.
- versicolor** Panzer, 1804: 140 (*Musca*). Germany, Regensburg. T A Unknown. Preocc. Gmelin 1790.
- versicolor** Meigen, 1826: 77 (*Musca*). Austria. ST A Lost. Muscidae, *Dasyphora penicillata* Egger. Preocc. Gmelin 1790. See Pont 1986b: 250.
- versicolorata** Turton, 1801: 639 (*Musca*). N. n. *Syrphus versicolor* Fabricius. Syrphidae, *Parhelophilus versicolor* Fabricius.
- versipellis** Gmelin, 1790: 2859 (*Musca*). Europe. T A NMI.
- vesiculosa** Turton, 1801: 646 (*Musca*). Subsequent Usage of *Syrphus vesiculosus* Fabricius. Syrphidae, *Copestylum vesiculosum* Fabricius.
- vespaesimilis** Barbut, 1781: 300, pl. 16, fig. 2 (*Musca*). N. n. *Musca vespiformis* Linnaeus. Syrphidae, *Tennostoma vespiformis* Linnaeus. *Xanthogramma festiva* L. was the species actually figured!
- vespertina** Fallén, 1823: 58 (*Musca*). Sweden. LT ♂ NRS. Muscidae, *Hebecnema vespertina* Fallén. Lectotype designated by Pont 1984: 295.
- vespiformis** Linnaeus, 1758a: 593 (*Musca*). Sweden. LT ♂ LSL. Syrphidae, *Tennostoma vespiformis* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 161.
- vespiformis** Geoffroy, 1785: 478 (*Musca*). France, Paris. T A Author. Syrphidae, *Xanthogramma festiva* Linnaeus. Preocc. Linnaeus 1758.
- vespillo** Fabricius, 1794: 318 (*Musca*). Germany, Kiel. LT ♂ UZMC. Calliphoridae, *Bellardia vespillo* Fabricius. "HT not in Kiel, nor label" Townsend 1931a: 70. Lectotype designated by Rognes 1991a: 59.
- vetustissima** Walker, 1849: 902 (*Musca*). Australia. HT ♀ BMNH. Muscidae, *Musca vetustissima* Walker. See Pont 1973: 148 & 1990: 169.
- vibrans** Linnaeus, 1758a: 599 (*Musca*). Europe. T A LSL. Otitidae, *Seioptera vibrans* Linnaeus. 5 sp. in LSL (Dear MS).
- vibrans** Harris, 1780: 122, pl. 35, fig. 43 (*Musca*). England. T A Unknown. Sepsidae, *Sepsis cynipsea* Linnaeus. Preocc. Linnaeus 1758.
- vicanus** Harris, 1780: 152, pl. 45, fig. 78 (*Musca*). England. T A Unknown. Muscidae, *Drymeia vicana* Harris.
- vicaria** Walker, 1853b: 348 (*Musca*). New Zealand. HT ♀ BMNH. Muscidae, *Musca domestica* Linnaeus. See Pont 1973: 169 & 1990: 170.

- vicina* Walker, 1849: 893 (*Musca*). Subsequent Usage of *Calliphora vicina* Robineau-Desvoidy. Calliphoridae, *Calliphora vicina* Robineau-Desvoidy.
- vicina* Macquart, 1851: 226 (*Musca*). America [=Martinique]. T ♂ MNHNP. Muscidae, *Musca domestica* Linnaeus.
- viduata* Linnaeus, 1758a: 598 (*Musca*). Sweden. LT ♂ LSL. Syrphidae, *Pipizella viduata* Linnaeus. Lectotype designated by Thompson *et al.* 1982: 161.
- villis* Gravenhorst, 1807: 380 (*Musca*). Unknown, probably Germany. T A Lost. Muscidae.
- villeneuvii* Patton, 1922b: 74 (*Musca*). India, Kerala, Nilambur. ST A BMNH. Muscidae, *Musca villeneuvii* Patton. See Pont 1990: 170.
- villica* Gravenhorst, 1807: 378 (*Musca*). Unknown, probably Germany. T A Lost. Tachinidae.
- villosa* Turton, 1801: 655 (*Musca*). Subsequent Usage of *Stratiomys villosa* Fabricius. Stratiomyidae, *Lasiopa villosa* Fabricius.
- vinculata* Scudder, 1877: 758 (*Musca*). Colorado, White River [Oligocene]. T ♂ USNM.
- vinulus* Harris, 1780: 117, pl. 34, fig. 21 (*Musca*). England. T A Unknown. Tephritidae.
- violacea* Scopoli, 1763: 340 (*Musca*). Carniola. T A DESTROY. Stratiomyidae, *Sargus cuprarius* Linnaeus.
- violacea* Panzer, 1804: 209 (*Musca*). Germany, Regensburg. T A Unknown. Preocc. Scopoli 1763.
- violacea* Fabricius, 1805: 288 (*Musca*). South America. HT ♀ UZMC. Calliphoridae, *Choloroprocta idioidea* Robineau-Desvoidy. Preocc. Scopoli 1763. See Michel-sen 1979a: 191 & Dear 1985b: 114.
- violacea* Robineau-Desvoidy, 1830: 393 (*Byomyia*). France, Saint-Sauveur. T ♀ DESTROY. Muscidae, *Musca tempestiva* Fallén.
- violacea* Meigen, 1838: 301 (*Musca*). Germany, Stolberg. HT ♂ MNHNP. Calliphoridae, *Protocalliphora azurea* Fallén. Preocc. Scopoli 1763. See Rognes 1991a: 116.
- violacea* Walker, 1853b: 335 (*Musca*). New Zealand. T ♀ BMNH. Calliphoridae, *Calliphora quadrimaculata* Swederus. Preocc. Scopoli 1763.
- viredaurea* Wiedemann, 1819a: 22 (*Musca*). Java. T ♂ UZMC. Calliphoridae, *Isomyia viredaurea* Wiedemann.
- virens* Scopoli, 1763: 342 (*Musca*). Carniola. T A DESTROY. Dolichopodidae, *Liancalus virens* Scopoli.
- virens* Thunberg, 1789: 91 (*Musca*). T A UNKNOWN. Preocc. Scopoli 1763.
- virens* Wiedemann, 1830: 409 (*Musca*). Java. T ♀ RNH. Calliphoridae, *Cosmina limbipennis* Macquart. Preocc. Scopoli 1763. N. Syn.
- virescens* Stephens, 1829: 307 (*Musca*). England. T A UNKNOWN. Muscidae, *Helina impuncta* Fallén. Published in synonymy, not subsequently validated by usage. Published in synonymy with *Anthomyia impuncta* Fallén.
- virescens* Meigen, 1838: 305 (*Musca*). Subsequent Usage of *Pollenia virescens* Macquart. Calliphoridae, *Pollenia vespillo* Fabricius.
- virginiensis* Drury, 1773: 993 (*Musca*). Virginia. T A Author. Syrphidae, *Milesia virginiensis* Drury.
- viridana* Panzer, 1804: 154 (*Musca*). Germany, Regensburg. T A Unknown.

- viridaurea* Wiedemann, 1830: 397 (*Musca*). Missp. *viredaurea* Wiedemann. Calliphoridae, *Isomyia viredaurea* Wiedemann.
- viridescens* de Villers, 1789a: 548 (*Musca*). France, Massiliae [=Marseille]. T A Author. Tephritidae. Preocc. de Villers 1789: 463.
- viridescens* de Villers, 1789a: 463, pl. 9, fig. 24 (*Musca*). Southern France, Marseille. T A Author. Syrphidae, *Microdon* sp.
- viridescens* Walker, 1849: 895 (*Musca*). Subsequent Usage of *Calliphora viridescens* Robineau-Desvoidy. Calliphoridae, *Calliphora viridescens* Robineau-Desvoidy.
- viridi-caerulea* De Geer, 1776: 62 (*Musca*). N. n. *Musca cadaverina* Linnaeus. Calliphoridae, *Lucilia caesar* Linnaeus.
- viridula* Gmelin, 1790: 2835 (*Musca*). Subsequent Usage of *Stratiomys viridula* Fabricius. Stratiomyidae, *Oplodontha viridula* Fabricius.
- vitrata* Geoffroy, 1785: 475 (*Musca*). France, Paris. T A Author.
- vitreus* Harris, 1776: 48, pl. 11, fig. 10 (*Musca*). England. T A Unknown. Stratiomyidae, *Microchrysa polita* Linnaeus.
- vitripennis* Meigen, 1826: 73, pl. 43, fig. 34 (*Musca*). France, Fontainebleau. HT ♂ MNHNP. Muscidae, *Musca vitripennis* Meigen. See Pont 1986b: 250.
- vittata* Fabricius, 1794: 355 (*Musca*). Guinea. T A UZMC F. Tephritidae, *Carpophthoromyia vittata* Fabricius.
- vivax* Fabricius, Otto, 1780: 206 (*Musca*). Greenland, Millutarsuk. T A UZMC. Syrphidae, *Platycheirus cyaneus* Müller.
- vivipara* Geoffroy, 1785: 471 (*Musca*). France, Paris. T A Author.
- vivipara* Bezzi, 1911: 86 (*Musca*). Missp. *larvipara* Portschinsky. Muscidae, *Musca larvipara* Portschinsky.
- vivipara major* De Geer, 1776: 63, pl. 3, fig. 18 (*Musca*). *Nomen nudum*. N. n. *Musca carnaria* Linnaeus. Sarcophagidae, *Sarcophaga carnaria* Linnaeus. Not available; polynominal, not binomial.
- vivipara minor* De Geer, 1776: 70 (*Musca*). *Nomen nudum*. [Sweden]. T A NRS. Calliphoridae. Not available; polynominal, not binomial.
- volets* Harris, 1780: 84, pl. 25, fig. 12 (*Musca*). England. T A Unknown. Tachinidae, *Dexiosoma caninum* Fabricius.
- volito* Harris, 1780: 122, pl. 36, fig. 46 (*Musca*). England. T A Unknown. Anthomyiidae, *Hylemya vagans* Panzer. Name improperly formed (verb).
- volvulus* Fabricius, 1794: 328 (*Musca*). Italy. T A UZMC F. Tachinidae, *Phyllomya volvulus* Fabricius.
- vomitoria* Linnaeus, 1758a: 595 (*Musca*). Europe & America. T A LSL. Calliphoridae, *Calliphora vomitoria* Linnaeus. 2 sp. in LSL (Dear MS).
- vorax* Geoffroy, 1785: 486 (*Musca*). France, Paris. T A Author. Syrphidae.
- vulpina* Fabricius, 1775: 776 (*Musca*). Europe. LT ♂ UZMC F. Muscidae, *Graphomya maculata* Scopoli. Lectotype designated by Michelsen 1979a: 191.
- vulpina* de Villers, 1789: 554 (*Musca*). Subsequent Usage of *Musca vulpina* Fabricius. Muscidae, *Graphomya maculata* Scopoli.
- wittweri* Zielke, 1974: 12 (*Musca*). Kenya, 26 Miles SW Nairobi. HT ♂ CAS. Muscidae, *Musca wittweri* Zielke.
- xanthomela* Walker, 1859b: 139 (*Musca*). Sulawesi, Makassar. HT ♀ BMNH. Muscidae, *Musca ventrosa* Wiedemann. See Pont 1990: 170.
- xanthomelas* Wiedemann, 1824: 49 (*Musca*). Java. T ♀ UZMC. Muscidae, *Musca xanthomelaena* Wiedemann.

- yerburyi** Patton, 1923b: 329 (*Musca*). N. n. *Musca incerta* Patton. Muscidae, *Musca pattoni* Austen.
- zonalis** Turton, 1801: 644 (*Musca*). Subsequent Usage of *Syrphus zonalis* Fabricius. Syrphidae, *Phytomia zonata* Fabricius.
- zonaria** Schrank, 1781: 454 (*Musca*). Subsequent Usage of *Conops zonaria* Poda. Syrphidae, *Volucella zonaria* Poda.
- zonatus** Fabricius, 1787: 337 (*Syrphus*). China. T A UZMC. Syrphidae, *Phytomia zonata* Fabricius.

## Index to names by biogeographical realm

**Afrotropical:** *afra*, *africa*, *africana*, *alba*, *albiceps*, *albomaculata*, *alpesa*, *beckeri*, *bibula*, *bigemmis*, *biseta*, *braesia*, *calisia*, *calleva*, *capensis*, *chloropyga*, *chrysotelus*, *congolensis*, *curviforceps*, *cuthbertsoni*, *cyanea*, *dasyops*, *dichotoma*, *distincta*, *dorsomaculata*, *elara*, *elatior*, *elongata*, *ethiopica*, *eustolia*, *exalbida*, *fasciata*, *freedmani*, *fucina*, *gabonensis*, *guineensis*, *hauriens*, *hebes*, *heidiae*, *himella*, *hottentota*, *hugonis*, *inconstans*, *interrupta*, *interruptella*, *irwinrossi*, *kihuris*, *kweilinensis*, *lasiopa*, *lasiophthalma*, *leucocephala*, *leucomelas*, *liberia*, *lindneri*, *longipes*, *lothari*, *lugens*, *lusoria*, *luteola*, *marginalis*, *marsya*, *mazaca*, *megacephala*, *meruensis*, *moerens*, *munroi*, *natalensis*, *nevilli*, *obscura*, *patersoni*, *perlata*, *phara*, *polit*, *pseudocorvina*, *pulla*, *punctulata*, *pungoana*, *putoria*, *quadrilineata*, *quinquelineata*, *racilia*, *rufifrons*, *rufiventris*, *sanctae-helenae*, *sarsina*, *scatophaga*, *senegalensis*, *setulosa*, *somalorum*, *sorbens*, *spangleri*, *spectanda*, *spilogaster*, *splendens*, *splendida*, *stuckenbergi*, *tempestatum*, *terminata*, *teste*, *transvaalensis*, *turbida*, *ugandae*, *varians*, *vittata*, *wittweri*.

**Australian:** *angustifrons*, *antiquissima*, *arcioides*, *augur*, *australia*, *australis*, *benedicta*, *bilineata*, *bimaculata*, *calliphoroïdes*, *cassara*, *cingulatus*, *costalis*, *cuprea*, *dawsoni*, *diffusa*, *ditissima*, *donysa*, *dorsalis*, *dubia*, *dux*, *eristaloïdes*, *extrema*, *fergusoni*, *fieldi*, *flavifacies*, *flavinervis*, *formosa*, *fulvimaculata*, *germana*, *gloriosa*, *hilli*, *hortona*, *icela*, *inconclusa*, *laemica*, *leonina*, *liris*, *lupina*, *macularis*, *marginifera*, *maronea*, *mesembrinoides*, *minor*, *mirabilis*, *nigricans*, *nigricurvis*, *nitescens*, *obscura*, *obscurata*, *obtrusa*, *oceanica*, *opulenta*, *pacifica*, *patiens*, *placida*, *praecox*, *prisca*, *pumila*, *quadrimaculata*, *regalis*, *retusa*, *ruficornis*, *sacra*, *sarcophagoïdes*, *satyrus*, *seelandica*, *sinuata*, *solaia*, *sordidissima*, *splendida*, *sylvanus*, *tahitiensis*, *taitensis*, *terraereginae*, *trifascia*, *trilineatus*, *varia*, *vetustissima*, *vicaria*, *violacea*.

**Nearctic:** *americana*, *amoena*, *ascarides*, *aurulans*, *bibosa*, *cadaverum*, *cerasina*, *certima*, *clavata*, *cloacaris*, *compressa*, *contigua*, *familiaris*, *filata*, *flavipennis*, *groenlandica*, *harpyia*, *heraea*, *hirticollis*, *hydropica*, *idessa*, *ilerda*, *lilaea*, *mollis*, *monoculus*, *mortisequa*, *muralis*, *occidentis*, *ochrapesus*, *pennipes*, *picta*, *proxima*, *serva*, *stygia*, *valida*, *vicina*, *vinculata*, *virginiensis*, *viridescens*, *vivax*.

**Neotropical:** *acromion*, *aequinoctialis*, *agrorum*, *aldrichi*, *americana*, *analis*, *anceps*, *annonae*, *annulata*, *atrifrons*, *auriflua*, *aurulans*, *australis*, *basalis*, *basilaris*, *bicolor*, *bifasciata*, *bipuncta*, *brunnea*, *caruca*, *chilensis*, *cincta*, *cingulata*, *cluvia*, *compressa*, *concolor*, *consanguinea*, *contigua*, *costalis*, *crassa*, *crassicornis*, *crucis*, *cylindracea*, *cylindricus*, *dentipes*, *diaphana*, *dilecta*, *dispar*, *elongata*, *equestris*, *erythrocephala*, *esuriens*, *eximia*, *exinanita*, *fabricii*, *fasciata*, *fucata*, *fulvescens*, *gamelia*, *gibba*, *grisea*, *gulo*, *hirta*, *hortorum*, *hystrix*, *illucens*, *inanis*, *incerta*, *insularis*, *jamaicensis*, *laniaria*, *lasciua*, *lateralis*, *leprae*, *leucopa*, *longicornis*, *longipes*, *lucifer*, *lyrcea*, *macellaria*, *mactans*, *mutabilis*, *nigrana*, *nigriceps*, *nigrita*, *nitida*, *obesa*, *occidua*, *ochricornis*, *pallipes*, *pampasiana*, *petiolata*, *phauda*, *pica*, *pinguis*, *pionia*, *pratorum*, *punctata*, *purpurascens*, *purpurea*, *pusilla*, *putrida*,

*quadrilineata, rufiventris, rutilans, scutellaris, segmentaria, semiatra, sensifera, stigma, stipata, stomoxidea, striata, suffusa, taniaria, tessellata, testacea, testaceus, tibialis, totiana, tricincta, turbida, variabilis, venatoria, verena, vesiculosa, violacea.*

**Oriental:** *abdominalis, adumbrata, aenea, affixa, albina, aruorum, arvorum, asiatica, aucta, azurea, bakeri, bata, bezzii, bivittata, chrysoides, chui, cinerea, cingalaisina, coerulata, coerulea, collecta, combrea, conducens, confiscata, confixa, conflagrans, convexifrons, craggi, crassa, crax, crux, cuprina, defixa, delectans, determinata, diffidens, dioclea, discolor, divaricata, divisa, dotata, electa, ericetorum, errans, eutaeniata, exempta, favillacea, ferruginea, flavicans, flaviceps, fletcheri, formosana, fortunata, fumipennis, fuscipennis, gavisa, gibsoni, greeni, gurnei, gurneyi, hervei, histrionica, hoi, hortensia, hortulana, humilis, ieuna, illingworthi, incerta, indica, inducta, inferior, infixa, ingens, inscribens, insignis, intrahens, jacobsoni, jeiuna, jejuna, kasauliensis, lar, larvipara, latifrons, latiparafrons, lauta, ligurriens, lineata, lucens, malaisei, mallochi, mediana, metilia, micans, minor, minuta, minutior, modesta, multispina, munda, nancauriensis, nebulo, negriabdomina, nigricornis, nigrithorax, niveisquama, optata, orientalis, pallens, pattoni, perfixa, phellia, pilifacies, pilosa, planiceps, polita, pollinosa, porphyrina, praedicens, prashadii, primitiva, proerna, proferens, promisca, promittens, prospera, pusillus, quadrilineata, quadrilineatus, quinquestriata, refixa, reflectens, remuria, ruficornis, rufifrons, salviae, santoshi, scapularis, selecta, seniorwhitei, serenissima, setigera, shanghaiensis, soror, sperata, sphinx, spinohumera, spinosa, squamata, striacta, taenionota, temperata, tibiseta, tifata, torosa, tranquebarica, trita, troglodyta, umbrifera, unisetula, varia, varicolor, ventrosa, villeneuvii, viredaurea, virens, viridaurea, xanthomela, xanthomelas, yerburyi, zonalis, zonatus.*

**Palaearctic:** *abbreviata, abdominalis, ablectus, abrupta, accincta, acuminata, adiaphora, adspersus, adunata, aenea, aerea, aestiuia, aestivationis, aestuans, aethiops, affinis, agilis, agilitas, agromyzina, alata, albicincta, albifrons, albimana, albipennis, albipes, albisetia, albolineata, albomicans, albula, alceae, alevo, allevo, allii, alni, alterabilis, alternata, altratus, aluta, ambigua, ambulans, amentaria, amica, amicula, amita, angelicae, angusticornis, anilis, annexa, annulata, annulatus, anteaimbulo, anteambulo, anthracina, anthrax, aphidioides, aphidivora, aparia, apiformis, apivora, aquatica, aratoria, arbustorum, arctii, arcuata, ardea, argentata, argentea, argyrostoma, aristata, armipes, arnicae, arrogans, artemisiae, aselliformis, asiliformis, assimilis, ater, aterrima, atherix, atra, atramentaria, atrata, atratula, atratus, atricolor, atropos, auctus, audaculus, aurata, aurifacies, aurifrons, ausus, autumnalis, avarus, avenae, azurea, balteata, bardus, bicincta, bicinctata, bicolor, bicolora, bidentata, bifasciata, bilineata, bipunctata, blandus, boganidae, bohemica, bombinatrix, bombylans, bombyliformis, bombyolina, bombylius, bombyloides, borealis, bovina, brachyptera, brassicaria, brevis, brunnea, buccata, buphthalmi, cadaverina, caemeteriorum, caerulea, caeruleophthalmica, caerulescens, caesar, caesarina, caesarion, caesia, caesio, calamitosa, calcarata, calcitrans, callosus, campestris, campicola, canicularis, canina, cannabina, canus, capitata, capito, capraria, capucina, caracteristica, carbonaria, cardui, carinifrons, carnaria, carnarida, carnifex, carnivora, casei, caudata, caudex, cautus, cellaris, celsus, cemiteriorum, centaureae, cephalotes, cerasi, cerinus, chaerophylli, chalybaea,*

chalybata, chalybea, chalybeata, chamaeleon, chloris, chorea, chrysocephala, chrysoptera, chrysopus, chrysorrhaea, chrysorrhoea, chrysostoma, chrysozonias, cibaria, cicur, ciliata, cimecoides, cimicoides, cincta, cinctus, cineraria, cinerascens, cinerea, cinerella, cinereus, cinerius, cingulata, citreofasciata, citrofasciata, clandestina, clauata, clauda, clauipes, clavicornis, clavicus, claviformis, clavipes, claviventris, clemens, coalescens, coarctata, coccinea, coeo, coerulea, cognata, coleoptrata, colossus, combinata, comio, comis, comito, commoror, comoror, compressa, compuncitor, compunctus, comtus, concinna, conferta, confinis, conflu, conformis, confusus, conica, connexa, conopsea, conopsoides, conosoma, consimilis, consisto, consobrinus, conspersus, conssecis, constane, constans, continua, contracta, convenio, convolo, corni, cornicena, cornicina, cornuta, corollae, corriginata, coruina, corvina, corvinoides, coryden, corydon, coryleti, coryletti, coryli, cossenico, costata, cothurnata, crabroniformis, crassipennis, crassipes, crassiostris, crassiseta, cibrata, cristata, crocus, cruciata, cruentata, cryptarum, cucularia, cupraria, cuprea, cursitans, cursoria, curvinervis, curvipes, cyanea, cyanella, cyanophthalma, cylindrica, cylindricornis, cynipsea, cynophila, dauci, deceptoria, decolor, decor, deduco, degener, dejectus, delicate, demandata, demano, denigrata, dentipes, depressa, depuncta, despecta, devia, dexter, diabolus, diadema, diaphana, dimano, dimidiata, diophthalma, dirae, discoidea, discolor, dispar, divaricata, divisa, dizonias, dolens, doliata, domestica, domestica major, domitor, doronici, dorsalis, dubia, dubitata, duplicita, efflatouni, elata, elater, elegans, elephantopus, eligans, elongata, emoda, ephippium, equestris, erinacea, erinaceus, erratica, erro, erucarum, erythrocephala, erythrophthalma, evectus, evonymellae, exactus, excors, exilis, extenuata, extranea, facultas, falcata, fallax, familiaris, fasciata, fasciatus, fasciolata, felina, felix, femorata, fenestratalis, fera, ferina, ferruginea, fervida, festinans, festiva, fector, filiformis, fimetaria, fimet, fimetorum, flauescens, flauicans, flava, flaveola, flavescentis, flavicans, flaviceps, flavidornis, flavidrons, flavidalpis, flavidipennis, flavidipes, flavidissima, floralis, florea, florescens, florescentiae, florid, formosa, formosus, frigida, frit, frondescentiae, frontalis, frutetorum, fucatus, fuciformis, fulges, fulva, fulvicornis, fulvipalpis, fulvipes, fulvithorax, fulviventrism, fulvus, fumigata, funebres, funebris, funesta, fungorum, funibres, furcata, furt, fusca, fuscata, fuscipennis, fuscipes, fuscula, fuscus, fusiformis, galeata, galpinus, gangraenosa, gastrodes, genarum, geniculata, gentilis, geographica, germanica, germinationis, gibba, gibbosa, glabrata, glabricula, glaucia, glaucoptera, globosa, globulipes, graeca, graminum, grandis, granditarsa, grisea, grisella, grisescens, grossa, grossificationis, grossipes, guttata, gymnosomea, haemorrhhoa, haemorrhoidalis, halterata, hamata, hartigii, hebes, hederacea, helenii, helianthi, helluo, hemiptera, hemipterus, hemisphaerica, heraclei, heraclii, hieracii, hilaris, hirsuta, hirsutissima, hirta, hirticeps, hordei, hortensis, horticola, hortorum, hortulana, hottentotta, hyalinata, hydroleon, hydrophila, hyoscyami, hypoleon, iacobaeae, ichneumonea, ichneumonoides, ignavus, iliaca, illustratus, illustris, imbellis, impavida, importuna, impuncta, inaequalis, inanis, incurvata, inda, indicus, inflata, ingredior, ingrue, ingrus, intermedia, interpuncta, interpunctionis, interpunctus, interrupta, interruptus, intersecta, interventum, interventus, intricaria, intricata, investigator, invisito, ionia, iridata, irritans, jacobaeae, japonica, jejuna, joco, labiata, lacustris, laeta, lagyra, lamed, lancifer, lanio, lappae, lappona, lardaria, larvarum, larvata, larvincola, lateralis, lateritia, laternaria, laticornis, latifrons, latipes, latrinarum, latro, leneatus,

lenio, lenis, leontodontis, lepida, lepidus, leporinus, leptogaster, leucocephala, leuconyx, leucophaeus, leucopis, leucoptera, leucozona, levidus, libatrix, limatus, limbata, lincensis, lineata, lineatus, lineola, lineolae, lineolata, lingens, linzensis, lithophila, liturea, litoriae, liturata, litus, livens, longerro, longervo, longicornis, longipennis, longipes, longisco, longula, lucida, lucidula, lucophaeus, lucopheus, lucorum, lucozona, luctuosa, ludeus, ludicrous, ludicrus, ludifacies, ludifica, lugubris, lunata, lunula, lupulina, lurida, lustrator, lutaria, lutea, luteipennis, luteola, luteus, lychnidis, lyra, macellaria, macrocephala, macroceras, macrophthalmia, maculata, maculatus, major larvarum, malansi, manicata, mano, mantes, mantis, mantos, marginalis, marginata, marginella, maritima, marmorata, martia, maura, meditabunda, meditalnenda, meditata, melancholica, melanius, melanochrysa, melanopsis, melanoptera, melanopyrrha, melanura, melinus, mellina, mellinus, menthastris, merdaria, meridiana, merulina, mesoleuca, mesopotamiensis, metallina, meteorica, meticulosa, metus, micans, microleon, miliaria, minima, minimus, minor, minor domestica, minor larvarum, minuta, minutissima, minutus, moesta, molio, molitus, mollicula, monachae, moneta, morio, mortuorum, muliebris, mulinus, mulsus, muraria, murina, murorum, muscaria, mussitans, mutabilis, mutata, myennis, myopina, mystacea, naevia, naevosus, nana, napobrassicae, nasatus, nasuta, natans, navus, nebulosa, nectarea, nectarous, nectarina, nectarivora, nemorum, neustriae, nigra, nigricans, nigricolor, nigrina, nigripennis, nigripes, nigripilis, nigrita, nigromarginata, nitens, nivalis, nobilis, nobilitata, noctiluca, nora, norwegica, nosatus, notata, nova, novemmaculata, novempunctata, nugater, nugator, obfuscata, obscoena, obscura, obsidianus, obsoleta, obtusipennis, ochracea, ochroleuca, ochroptera, ochropus, octopunctata, ocymi, oenopota, oestracea, oleae, olens, olivacea, onopordinis, oppidana, oratoria, orbiculata, orbiculus, orior, ornate, ornithovora, osiris, ovipara, pabulorum, pagana, paganus, pallida, pallipes, palmipes, palpator, palustris, pandia, pantherina, papatas, parallelis, parallelus, parasitica, parcus, parietina, parralleli, parrallelus, parrelleli, parviceps, parvula, parvulus, pascuorum, pavida, pedella, pedissequus, pedo, pellita, pellucens, pendula, pennipes, peragro, percussus, perelegand, perelegans, perexilis, perexillis, permundus, pernix, personata, personatus, pertinax, pervenio, petronella, phalaenarum, phasiaeformis, picata, picea, piceae, pigra, pilosa, pinastri, piniperdae, pinivorae, pipiens, pipines, placida, planifrons, plebeja, plombea, plumata, plumbea, plumipes, plumosa, pluvialis, podagraria, poeciloptera, polisma, polita, politus, polyzonias, porca, porcina, potens, potentillae, praeceps, praedo, praepotens, praeusta, prasina, pratorum, princeps, procedo, processionae, prodeo, prodromaria, profuges, profugus, prolabor, prolapsa, prolapsor, promano, promans, properans, prostrata, provenio, pruinosa, psyllia, puber, pubera, puberula, pubiceps, pudefactus, puella, pulchella, pulchra, pulicaria, pulla, pullata, pumila, pumilionis, punctata, punctatus, punctifera, punctulata, punctum, pungens, puparum, purmundus, pusilla, putris, pyrastri, pyrorum, quadrata, quadricolor, quadrifasciata, quadrimaculata, quadrimaculella, quadripunctata, quadripustulata, quadristrigata, quadrum, quinquefasciata, quinquepunctata, radiata, radicum, rapax, rara, rau, reccumbo, recedans, recta, recurvo, redambulo, redeans, redeo, redians, redditus, regalis, regina, relictus, remigro, repens, rerumbo, resinellae, resinosa, restituo, restituo, reticulata, reversio, revertio, reviso, revolo, rex, ribesii, riparia, rivularis, roralis, rosae, rosarum, rostrata, rotundata, rubiginea, rubripes, rudis, rufa, ruficapilla, ruficauda, ruficaudis, ruficeps, ruficornis, rufifcrus, rufifrons, rufina,

rufipalpis, rufipes, rufiventris, rufomaculata, rugosa, ruminis, ruralis, rustica, rutillus, rutilo, sabaeus, sacalis, saltatrix, saltitans, saluum, sandaracha, sanio, saniosa, sannio, scabiosae, scalaris, scitule, scitulus, scitus, scolopacea, scolopacina, scripta, scutata, scutellaris, scutellata, scybalaria, secalis, segetum, segmentaria, segnis, sellata, semiargentata, semiargentea, semicaerulea, semiflava, semiluctifera, seminationis, semirufa, semulater, semulator, senilis, seplasia, sepulchralis, sepulcralis, serena, sericata, sericea, serpo, serrata, serratula, serratulae, setaria, seticornis, setosa, sexdentata, sexmaculata, sexpunctata, sigmoidea, signata, silentis, silvarum, silvestris, similis, simulator, sinensis, singenesiae, singularius, smaragdina, socia, socio, sollicitus, solivagus, solor, solstitialis, sonchi, sordida, spatula, speculifera, sphegea, spinifera, spinipes, spitzbergensis, splendens, splendida, spoliata, spondylii, squalida, squamata, squamifera, stabulans, stellata, stercoraria, stictica, stigma, stimulans, striata, strigata, strigosa, strigula, striolata, structus, strumosa, stuans, stylata, subcoleoptrata, subcutanea, sublatus, subpuncta, subsultans, subtomentosa, subulata, subvariegata subventus, succincta, suecica, sugillatrix, suilla, surinamensis, sylvarum, syngenesiae, tabaniformis, tachinoides, taeniata, tarda, tardigradus, tarditas, tartidus, tau, tegularia, tempestiva, temula, tenax, tenebris, tener, tentaculata, tenur, teres, terrestris, tessella, tessellata, testacea, tetragona, thalassina, thuscia, thymastri, tiberina, tibetana, tibialis, tigrina, timeo, timidus, tipulaea, tipularia, tipuliformis, tomentosa, topiaria, torvis, torvus, transfuga, transversa, trelineata, tremula, triangula, tricolor, tridens, trifasciata, trigrammica, trileniata, trilenna, trilineata, trimaculata, trinervis, tringaria, tripunctata, triquetra, tristis, tristor, truncata, tuberculata, tuberis, tuguriorum, turcica, tussilaginis, uliginosa, umbellatarum, umbelliferarum, umbraculata, umbrarum, unguicornis, unguulata, unicolor, univalvis, unxia, urbanus, urticae, vaccarum, vaccina, vacua, vagabunda, vagans, vagatoria, vagus, valens, valentina, valga, validus, vallata, vanella, varensis, varia, variabilis, varians, variata, varicus, variegata, varieus, variolosa, vecors, velox, venosa, ventito, vermileo, verna, vernicina, versicolor, versicolorata, versipellis, vespaesimilis, vespertina, vespidiformis, vespillo, vibrans, vicanus, viduata, villosa, vinulus, violacea, vires, virescens, viridana, viridescens, viridi-caerulea, viridula, vitrata, vitreus, vitripennis, vivipara, vivipara major, vivipara minor, volets, volito, volvulus, vomitoria, vorax, vulpina, zonaria.

**Unknown:** atra, bicolor, capensis, carinata, castanea, colombaschensis, costata, doclea, elevata, flavipes, fusca, gigantea, helvolo, leucosticta, melanoptera, rubiginosa, ruralis, solstitialis, syrphus, vilis, villica.

## Index to names by current taxonomic position

- Acalyprate*: commoror, comoror,  
 hirsuta; leporinus, sigmaidea,  
 simulater, simulator, timidus.  
*Acanthiophilus*: helianthi.  
*Achanthiptera*: inanis.  
*Acinia*: fucata, furcata.  
*Aciura*: coryli.  
*Acrocera*: orbiculus.  
*Acroceridae*: gibbosa, orbiculus.  
 acromion: longipes.  
*Actia*: resinellae.  
*Adia*: cinerella.  
*aeneus*: leucocephala, macrourhala,  
 punctata.  
*aethiops*: scatophaga.  
*Agria*: affinis.  
*alba*: cinerea.  
*albiceps*: bibula, elara, emoda, felix,  
 himella.  
*albicinctus*: albicincta.  
*albifrontalis*: australis.  
*albina*: beckeri, speculifera.  
*albiseta*: aratoria, crocus, oratoria.  
*albonotata*: prospera.  
*alceae*: demandata.  
*aldrichi*: esuriens.  
*Alliopsis*: silvestris.  
*Allophora*: affinis, hemiptera, nebulosa,  
 subcoleoptrata, tristis.  
*alpaea*: congolensis.  
*Amenia*: leonina, varia.  
*amentaria*: nigrina, nitens.  
*amita*: amica.  
*Anabarynchus*: bilineata.  
*Anapunophyo*: pusilla.  
*Angiometopa*: ruralis.  
*angulicornis*: erinacea.  
*anilis*: flavipes, ochropus.  
*Anisopodidae*: bilineata, cinctus, fuscata,  
 nigricans, punctatus, succincta.  
*annexa*: bifasciata, dizonias.  
*annulatus*: annulata.  
*Anomoia*: permundus, purmundus.  
*Anthomyia*: litus, pluvialis.  
*Anthomyiidae*: albicincta, albula,  
 ambigua, auctus, audaculus, bicolor,  
 buccata, cinerea, coarctata,  
 comio, comis, conformis, conica,  
 decore, dubia, dubitata, exactus,  
 excors, exilis, flavipennis, flavipes,  
 floralis, fungorum, grisea, hilaris,  
 lancifer, laticornis, lenio, lenis,  
 liturariae, litus, longula, myopina,  
 napobrassicae, parvula, pluvialis,  
 paeoptens, processionae,  
 pudefactus, radicum, redditus, rufina,  
 rufipes, rumicis, silvestris, socia,  
 solor, strigosa, striolata, structus,  
 testacea, transversa, vagans, variata,  
 volito.  
*Anthomyiidae?*: cristata.  
*Anthrax*: aethiops, anthrax.  
*Anthrax*: hottentotta.  
*antiqua*: liturariae.  
*Aphyssura*: dubia.  
*aquatica*: graminum.  
*arbustorum*: horticola, nemorum,  
 parallelis, parralleli, parrelleli.  
*Archytas*: diaphana, hystrix, pilosa.  
*Arctophila*: fulvus, mussitans.  
*arcuatum*: arcuata, bipunctata, festiva,  
 imbellis.  
*Argyra*: diaphana, ludeus, semiargentata,  
 semiargentea.  
*argyrocephala*: frontalis, leucocephala.  
*arrognis*: cimocoides, cimicoides.  
*arvorum*: aruorum, quadrilineatus,  
 tranquebarica.  
*Asarkina*: ericetorum, salviae.  
*aterrima*: brevis.  
*Athericidae*: atherix, chrysopus, flavipes,  
 maculatus.  
*Atherix*: atherix, chrysopus, flavipes,  
 maculatus.  
*atra*: aterrima.  
*atriceps*: boganiidae.  
*Auchmeromyia*: luteola.  
*audacula*: audaculus.  
*augur*: dorsalis.  
*auratus*: tomentosa.  
*auriceps*: opulenta.  
*aurifinis*: auriflua.  
*aurulenta*: evonymellae.  
*australis*: donysa, maronea, tessellata.

- autissiodorensis: flavigalpus.  
 autumnalis: bovina, campestris,  
     continua, coruina, corvina, floralis,  
     grisella, ludificies, ludifica, ovipara,  
     prashadii, prostrata, rau, restitituo,  
     restituo, riparia, rustica, tau.
- Azelia*: ornata, triquetra.  
*azurea*: pervenio, reviso, violacea.  
*Baccha*: elongata, perekilis, perekillis.  
*Bactrocera*: ferruginea.  
*Bactromyia*: evonymellae.  
*barda*: americana.  
*Baumhaueria*: angusticornis.  
*Bellardia*: agilis, dolens, pandia,  
     puberula, pusilla, unxia, vespillo.  
*Belvosia*: aldrichi, bifasciata, esuriens.  
*Bengalia*: favillacea, iejuna, iejuna,  
     jejuna, pallens, torosa, varicolor.  
*Beris*: chalybata, chalybea, chalybeata,  
     clavipes, rex, sexdentata, similis,  
     vallata.  
*bezzi*: pilosa.  
*Bibio*: bidentata, hortulana.  
*Bibionidae*: bidentata, ciliata, hortulana,  
     novemmaculata, tuberculata.  
*bicinctum*: bicincta, bicinctata, callosus.  
*bilineatus*: bilineata.  
*Billaea*: rutilans, sensifera.  
*bipunctatus*: bipunctata.  
*Blaesoxiphia*: lineata.  
*Blattariac*: lithophila.  
*Blera*: fallax, ruficaudis, semirufa.  
*Blondelia*: pinivora.  
*Bogosiella*: fasciata.  
*bombylans*: hirsutissima, melanopyrrha,  
     plumata, plumosa.  
*Bombyliidae*: aethiops, anthrax, bicolora,  
     capucina, denigrata, fusca, hirsuta,  
     hottentotta, lar, lucifer, maura,  
     melanoptera, morio, nigrata,  
     polyzonias, sabaeus, satyrus, sphinx,  
     sylvanus, troglodyta.  
*Bombyliidae?*: capensis.  
*Borophaga*: subsultans.  
*Botanophila*: cinerea, striolata.  
*Bothrophora*: lupina.  
*Brachyopa*: arcuata, conica.  
*Bradysia*: maura, morio.  
  
*brassicaria*: cylindrica.  
*brunneum*: brunnea, esuriens, exinanita,  
     inanis.  
*cadaverina*: aurulans, compressa, ilerda,  
     lilaea, mortisequa.  
*caesar*: azurea, cadaverina, fulges,  
     ruficeps, splendida, viridi-caerulea.  
*caesio*: lychnidis.  
*calcitrans*: occidentis, pungens.  
*calleva*: cuthbertsoni.  
*Calliopomyia*: glabrata.  
*Calliphora*: affinis, aucta, augur,  
     australis, carnaria, carnivora,  
     coerulea, dorsalis, erythrocephala,  
     flavigalpus, fulvimaculata, genarum,  
     laemica, laticornis, minimus,  
     obscoena, placida, quadrimaculata,  
     sacra, scutellata, spitzbergensis,  
     stygia, thuscia, vicina, violacea,  
     viridescens, vomitoria.  
*Calliphoridae*: abdominalis, accincta,  
     affinis, agilis, albiceps, alni,  
     amentaria, anthracina, atramentaria,  
     aucta, augur, aurulans, australis,  
     azurea, bata, bibula, bicolor,  
     bigemmis, bivittata, boganiidae,  
     cadaverina, cadaverum, caerulea,  
     caerulescens, caesar, caesarina,  
     caesarion, calliphoroides, carnaria,  
     carnarida, carnivora, caruca, certima,  
     chalybaea, chilensis, chloropyga,  
     chrysocephala, chrysoides,  
     chrysorrhoea, cinerea, cluvia,  
     coerulea, cognata, collecta, combrea,  
     compressa, concinna, confixa,  
     conflagrans, consobrinus, costalis,  
     cuprea, cuprina, defixa, degener,  
     delectans, depressa, diffusa, dioclea,  
     discolor, ditissima, divisa, dolens,  
     doronici, dorsalis, dotata, dubia, dux,  
     elara, emoda, equestris, eristaloides,  
     erythrocephala, exempta, eximia,  
     familiaris, fasciata, favillacea, felina,  
     felix, flavicans, flaviceps, flavigalpus,  
     floralis, fortunata, fucina, fulges,  
     fulvicornis, fulvimaculata,  
     fulvipalpis, fumipennis, fuscipennis,  
     gamelia, genarum, gentilis, germana,

- gloriosa, groenlandica, hebes,  
 hederacea, himella, hortona, icela,  
 iejuna, ilerda, illustris, impavida,  
 incerta, inconclusa, inducta, infixa,  
 ingens, inscribens, insularis,  
 intermedia, ionia, jejuna, jejuna,  
 laemica, lagyra, laniaria, lanio,  
 laticornis, leonina, leucocephala,  
 leucoptera, ligurriens, lilaea, lunata,  
 luteola, lyrcea, macellaria, macularis,  
 malansi, marginalis, marginifera,  
 megacephala, melanoptera,  
 mesembrinoides, metilia, micans,  
 minima, minimus, mollis,  
 mortisequa, mortuorum, munda,  
 muralis, nana, nigriceps, nigrina,  
 nigromarginata, nitens, nitescens,  
 nobilis, obscoena, obscura, obscurata,  
 olivacea, opulenta, orientalis, orior,  
 pallens, pandia, parvula, patiens,  
 perlata, pervenio, phauda, phellia,  
 ponia, placida, polisma, porphyrina,  
 promittens, prospera, proxima,  
 pruinosa, puberula, pulchra, pumila,  
 punctulata, purpurascens, purpurea,  
 pusilla, putoria, putrida,  
 quadrilineata, quadrimaculata, recta,  
 recurvo, reflectens, regalis, regina,  
 remigro, remuria, reviso, rufis,  
 ruficeps, ruficornis, rufipalpis,  
 rufipes, sacra, sarcophagoides,  
 scutellata, segmentaria, selecta,  
 semicaerulea, sepulcralis,  
 serenissima, sericata, serva, silvarum,  
 solaia, sordida, spitzbergensis,  
 splendida, stygia, suffusa, taniaria,  
 tegularia, temperata, terminata,  
 thalassina, thuscia, tifata, torosa,  
 trifascia, turbida, umbrifera, unxia,  
 vagabunda, varensis, varia, varians,  
 varicolor, ventito, verena, vespillo,  
 vicina, violacea, viredaurea, virens,  
 virescens, viridaurea, viridescens,  
 viridi-caerulea, vivipara minor,  
 vomitoria.  
 Calobata: petronella.  
 Calocarcelia: cingulata.  
 Calythea: albicincta.  
 campestris: macrocephala.  
 campestris?: cephalotes.  
 canicularis: lateralis, minor, socio.  
 caninum: canina, chrysostoma, volets.  
 cannabina: alternata, balteata, elegans,  
     inaequalis, nectareus, nectarina,  
     scitule, scitulus.  
 Capnoptera: scutata.  
 Carcelia: processioneae.  
 Cardiacephala: longipes.  
 carnaria: vivipara major.  
 carnaria?: colossus.  
 Carpophthoromyia: vittata.  
 casei: atrata.  
 Catapicephala: ingens, micans.  
 caustum: cautus.  
 celsa: celsus, quadrimaculata,  
     quadrimaculella, signata.  
 cemiteriorum: caemeteriorum, lineolata.  
 Centroplebomyia: furcata.  
 ceratocera: cornuta.  
 Ceratolauxania: elevata.  
 Ceriana: adunata, conopoides, univalvis.  
 Ceromyia: flaviceps.  
 Cerotelion: lineaata, striata.  
 Ceroxys: hortulana, hyalinata, urticae.  
 chaerophylli: variegata.  
 Chaetotheresia: crassa.  
 Chalcosyrphus: femorata, longipes,  
     valga.  
 chalybata: chalybea, chalybeata,  
     sexdentata.  
 chalybeata: promano, promans.  
 chamaeleon: spatula.  
 Chamaemyiidae: atratula, gibbosa.  
 Chamaepsila: rosae.  
 Cheilosia: aenea, bardus, coryden,  
     corydon, flavicornis, fulva, funebres,  
     funebris, funibres, hortulana,  
     illustratus, merulina, prasina, tarditas,  
     tartidus.  
 Chirosia: flavipennis.  
 Chloromyia: aurata, cicur, formosa.  
 Chloropidae: avenae, calamitosa,  
     clavicrus, erro, frit, hordei, lepidus,  
     lineata, minuta, nasuta, nigra, pigra,  
     planifrons, pulchella, pumilionis,

- quadristrigata*, *saltatrix*, *scutata*,  
*strigula*, *umbelliferarum*, *umbraculata*. *Coelopa: frigida*.  
*Chloroprocta: purpurea*. *Coelopidae: frigida*, *relictus*, *simulator*.  
*Chlorops: lepidus*, *lineata*, *nasuta*,  
*pulchella*, *pumilionis*, *quadristrigata*,  
*saltatrix*, *strigula*, *umbelliferarum*. *Coenomyia: australis*, *crucis*, *ferruginea*,  
*chloropyga: bigemmis*, *hebes*, *putoria*. *grandis*, *olens*, *testacea*.  
*Chlororhinia: exempta*. *Coenosia: agromyzina*, *flavicornis*,  
*Cholomyia: acromion*, *longipes*. *geniculata*, *intermedia*, *mollicula*,  
*Choloroprocta: violacea*. *pedella*, *pumila*, *rapax*, *tigrina*.  
*chrysame: varia*. *coeruleifrons: trita*.  
*Chrysogaster: caemeteriorum*,  
*cemiteriorum*, *lineolata*, *lucida*,  
*speculifera*. *Comatacta: tricincta*.  
*Chrysomya: albiceps*, *bata*, *bibula*,  
*bigemmis*, *chloropyga*, *combrea*,  
*defixa*, *dux*, *elara*, *emoda*, *felix*,  
*flaviceps*, *hebes*, *himella*, *marginalis*,  
*megacephala*, *orientalis*, *putoria*,  
*remuria*. *Compsobata: cibaria*, *cothurnata*.  
*Chrysophilus: asiliformis*, *cingulata*,  
*cristata*, *tomentosa*. *Compsomyiops: caruca*, *gamelia*, *lyrcea*,  
*verena*. *Comptosia: sylvanus*.  
*Chrysotoxum: arcuata*, *bicincta*,  
*bicinctata*, *bipunctata*, *callosus*,  
*cautus*, *chrysozonias*, *fasciata*,  
*fasciolata*, *festiva*, *imbellis*. *conducens: kweilensis*, *lineata*, *lucens*,  
*praecox*, *pulla*, *scapularis*.  
*Chyliza: extenuata*, *leptogaster*. *Condylostylus: longicornis*.  
*Chyromya: flava*, *luteola*, *ochrapesus*,  
*oppidana*. *confiscata: fasciata*, *minuta*.  
*Chyromyidae: flava*, *luteola*, *ochrapesus*,  
*oppidana*. *conspersa: conspersus*.  
*cibaria: cothurnata*. *convexifrons: gibsoni*.  
*cincta: vacua*. *convexifrons?: latiparafrons*.  
*cinctus: cincta*, *succincta*. *Copestylum: brunnea*, *esuriens*,  
*cingulatus: pacifica*. *exinanita*, *inanis*, *vesiculos*.  
*clausa: aenea*. *Cordilura: puber*, *pubera*.  
*clavatus: clavata*, *filata*. *Cordyligaster: compressa*, *petiolata*.  
*clavicornis: univalvis*. *Coremacera: cincta*, *cribrata*, *limbata*,  
*clavipes: clauipes*, *curvipes*. *marginata*, *tristis*.  
*Clorismia: ardea*. *corni: pagana*, *princeps*.  
*Citellaria: coccinea*, *ephippium*, *inda*,  
*sellata*. *cornicina: amoena*, *caesarion*, *chloris*,  
*corriginata*, *cornicena*, *heraea*, *ingredior*,  
*corydon: coryden*. *connexa*, *puella*, *redeo*.  
*Corrigiolata: filiformis*, *strumosa*. *coryleti: coryletti*, *reticulata*.  
*Cosmina: punctulata*, *virens*. *Craticulina: rufifrons*, *tabaniformis*.  
*crassipennis: aurifrons*, *brachyptera*,  
*dimidiata*, *taeniata*. *cristatus: cingulata*, *cristata*.  
*crassipes: elephantopus*. *cruentatus: cruentata*.  
*Craticulina: rufifrons*, *tabaniformis*.  
*coalescens: abdominalis*. *crux: crax*.  
*Cochliomyia: certima*, *erythrocephala*,  
*fasciata*, *laniaria*, *macellaria*, *phauda*,  
*taniaria*, *turbida*. *Cuphocera: varia*.  
*Cuprarius: capraria*, *cupraria*, *violacea*.  
*cuprea: nitens*, *rutilo*. *cuprina: fucina*, *serenissima*, *temperata*,  
*varians*.

- Curtonotidae: gibba.  
 Curtonotum: gibba.  
 Cuterebra: americana.  
 Cuterebridae: americana, semiatra.  
*cyanella*: mano.  
*cyaneus*: agilitas, albimana, cyanea,  
     vivax.  
*cyanicolor*: occidentis, seplasia.  
*cyanophthalma*: annulata.  
*cylindricornis*: chrysoptera.  
*cylindricus*: cylindracea.  
 Cylindromyia: brassicaria, cylindrica.  
*cynipsea*: elongata, vibrans.  
 Cynomya: aurulans, chrysocephala,  
     compressa, ilerda, lilaea, mortisequa,  
     mortuorum.  
 Cyrtoneurina: grisea.  
 Dacus: oleae.  
 Dasyphora: pratorum, versicolor.  
 Dasysyrphus: consisto, pinastri, pyrorum.  
 delectans: conflagrans.  
 Delia: albula, coarctata, dubia, floralis,  
     liturariae, napobrassicae, radicum.  
 Delphinia: picta.  
 Dexia: germanica, plebeja, provenio,  
     rustica, vacua.  
 Dexiosoma: canina, chrysostoma, volets.  
 Dexopollenia: trifascia.  
 diabolus: ciliata, spinipes.  
 diadema: rostrata.  
 Dialineura: anilis, flavipes, ochropus.  
 diaphana: ludeus.  
 diaphanus: diaphana.  
 Dictya: umbrarum.  
 diffidens: optata, prefixa, refixa.  
 Dilophus: ciliata, novemmaculata,  
     tuberculata.  
 Dinera: carinifrons, ferina, grisescens.  
 Diplonevra: florea, florescens.  
 discalis: lateralis.  
 distincta: obscura.  
 Dolichopodidae: amicula, caudex,  
     ciliata, diadema, diaphana, equestris,  
     festiva, gibba, hebes, ingruo, ingrus,  
     joco, lacustris, longicornis, ludeus,  
     ludicrous, ludicus, nigripes,  
     nobilitata, notata, pallipes, plumipes,  
     quadrifasciata, rostrata, sanio, sannio,  
     semiargentea,  
     squama, squamifera, ungulata,  
     virens.  
 Dolichopus: gibba, plumipes, squamata,  
     squamifera, ungulata.  
 domestica: analis, antiquissima,  
     atirfrons, aurifacies, aurulans,  
     australia, australis, basilaris,  
     campicola, chilensis, consanguinea,  
     contigua, dawsoni, determinata,  
     divaricata, divisa, domestica major,  
     flavifacies, flavinervis, flavigennis,  
     frontalis, fulvescens, gymnosomea,  
     harpyia, hottentota, lateralis, minor,  
     multispina, nancauriensis, nebulo,  
     nigricans, obscura, oceanica,  
     pampasiana, pellucens, rivularis,  
     rufifrons, rufiventris,  
     sanctae-helenae, senegalensis,  
     sordidissima, soror, stomoxidea,  
     taitensis, tiberina, umbraculata,  
     vaccina, vagatoria, vicaria, vicina.  
 domestica?: conosoma.  
 domitor: albolineata.  
 doronici: melanoptera.  
 Doros: conopsea, profuges, profugus.  
 dorsalis: ferruginea.  
 Dorycera: aquatica, graminum.  
 Drosophila: cellaris, erythrophthalma,  
     funebris, oenopota.  
 Drosophilidae: cellaris, coleoptrata,  
     erythrophthalma, funebris, furtia,  
     oenopota.  
 Drymeia: decolor, hamata, mantes,  
     mantos, vicanus.  
 Dryomyza: flaveola.  
 Dryomyzidae: flaveola, melinus,  
     mellinus, mulinus.  
 dubia: luteola, placida.  
 Dufouria: promano, promans.  
 Ectophasia: aurifrons, crassipennis.  
 elata: alata, elater.  
 elegans: formosa.  
 Elgiva: cucularia, rufa, sollicitus.  
 eligans: elegans, interrupta.  
 Emmesomyia: socia.

- Empididae: arrogans, cimecoides,  
     cimicoides, cursitans, cursoria,  
     globulipes, grossipes.  
 Engistoneura: lugens, moerens.  
 Ensina: sonchi, subcutanea.  
 ephippium: coccinea, inda, sellata.  
 Ephydriidae: manicata, mantis.  
 Epistrophe: elegans, eligans, formosus,  
     interrupta.  
 Episyphus: alternata, balteata,  
     cannabina, elegans, inaequalis,  
     nectareus, nectarina, scitule, scitulus.  
 equestris: bombyliformis.  
 Eriothrix: demano, dimano, lateralis,  
     rufomaculata.  
 Eriozona: erratica.  
 Eristalinus: aruorum, arvorum, ater,  
     leucocephala, macrophtalma,  
     melanius, punctata, quadrilineatus,  
     quinquelineata, quinquestriata,  
     sepulchralis, tranquebarica.  
 Eristalis: americana, apiformis,  
     arbustorum, cinctus, cryptarum,  
     fuscus, horticola, interruptus,  
     intricaria, intricata, leneatus, lineatus,  
     lineolae, lunula, lyra, murorum,  
     nemorum, obfuscata, oestacea,  
     parallelis, parralleli, parrelleli, porcina,  
     tenax.  
 Ernestia: puparum, redambulo, torvis,  
     torvus.  
 errans: erratica.  
 Euboettcheria: tessellata.  
 Eudasypora: cyanella, mano,  
     occidentis, seplasia.  
 Euleia: centaureae, flauescens, heraclii,  
     onopordinis, subcutanea.  
 Eumerus: lineata, lunata.  
 Eumesembrinella: quadrilineata.  
 Euphranta: connexa, crax, crux.  
 Euphumosia: bivittata, calliphoroides,  
     eristaloides.  
 Eurychaeta: alni.  
 Eustalomysia: hilaris.  
 Euthycera: chaerophylli, fumigata,  
     rufifrons, variegata.  
 Eutrichota: praepotens.  
 Euxesta: annonae, costalis.  
     evecta: evectus, lucorum.  
     eximia: insularis.  
 Exoprosopa: capucina.  
 Exorista: larvarum, larvincola, major  
     larvarum, monachae.  
 fabricii: cincta.  
 fallax: ruficaudis, semirufa.  
 falleni: ruralis, vagans.  
 Fannia: alevo, allevo, canicularis,  
     clemens, fuscula, lateralis, lepida,  
     lustrator, minor, minor domestica,  
     scalaris, serena, socio.  
 Fanniidae: alevo, allevo, canicularis,  
     clemens, fuscula, lateralis, lepida,  
     lustrator, minor, minor domestica,  
     scalaris, serena, socio.  
 fasciata: cingulata.  
 fasciata?: palmipes, properans.  
 fasciatum: chrysozonias, fasciata.  
 fasciolatum: fasciolata.  
 femoralis: geniculata.  
 femoratus: femorata, longipes.  
 fenestralis: saltitans, senilis, spoliata,  
     tarda.  
 fenestratus: nigrita.  
 fera: reccumbo, rerumbo.  
 Ferdinandea: nitens, ruficornis, rutilo.  
 fernandica: leucocephala.  
 ferruginea: australis, crucis, grandis,  
     olens, testacea.  
 festiva: anteaimbulo, anteambulo,  
     citreofasciata, citrofasciata,  
     vespiformis.  
 fimetaria: flava, rufa.  
 flava: luteola, ochrapesus.  
 flavipennis: quadrilineata.  
 flavipes: chrysopus.  
 fletcheri: jacobsoni.  
 floralis: micans, sepulcralis.  
 florea: atropos.  
 florescens: florea.  
 forcipata: maura, morio.  
 formosa: aurata, cicur.  
 formosana: greeni, shanghaiensis.  
 Formosia: mirabilis.  
 frit: avenae, hordei, nigra.  
 fuciformis: fusiformis.  
 fulva: libatrix, ochracea.

- fulvicrura: caruca, gamelia, lyrcea.  
 fumigata: rufifrons.  
 funebres: funebris, funibres.  
 funebris?: cellaris, erythrophthalma,  
     oenopota.  
 fuscatus: fuscata, nigricans.  
 fuscipennis: punctulata.  
 gabónensis: natalensis.  
 gavisa: chalybea.  
 genarum: laticornis.  
 geniculata: urbanus.  
 gentilis: caerulea, cognata.  
 Geomyza: combinata.  
 glaucia: depressa.  
 globosum: globosa.  
 globula: lingens.  
 gloriosa: costalis, diffusa.  
 Gonia: capitata, crassicornis, piniperdae.  
 goniaeformis: angusticornis.  
 Goniocera: hartigii.  
 granditarsus: confusus, granditarsa,  
     ocymi.  
 Graphomya: adumbrata, compuncitor,  
     compunctus, eustolia, idessa,  
     leucomelas, maculata, praedicens,  
     stipata, vulpina.  
 grisella: grisea.  
 grossulariae: formosus.  
 guttata: longipes.  
 Gymnosoma: cerinus, costata, globosa,  
     hemisphaerica, rotundata.  
 Haematobia: irritans.  
 halterata: maura.  
 halterata?: nigrita.  
 Hebecnema: nigricolor, nigrita,  
     vespertina.  
 Heleomyza: latinarum, serrata.  
 Heleomyzidae: avarus, dejectus,  
     latinarum, sandaracha, serrata,  
     tuberis.  
 Helina: atricolor, celsus, concolor,  
     confinis, consimilis, cornuta, deduco,  
     depuncta, evectus, fector, impuncta,  
     interpuncta, lucorum, obtusipennis,  
     quadrimaculata, quadrimaculella,  
     quadrum, reversio, revolo,  
     sexmaculata, signata, subpuncta,  
     uliginosa, virescens.  
 Heliozeta: helluo.  
 Helophilus: cingulatus, pacifica,  
     parallelus, parrallelus, pendula,  
     seelandica, trelineata, trileniata,  
     trilenva, trilineata, trilineatus.  
 Hemilucilia: segmentaria.  
 Hemipenthes: bicolora, denigrata,  
     hirsuta, maura, morio.  
 hemiptera: affinis, tristis.  
 Hemipyrellia: fortunata, leucocephala,  
     ligurriens, phellia, pulchra, solaia.  
 hennigi: rosae.  
 Heptatoma: pellita.  
 heraclii: centaureae, onopordinis,  
     subcutanea.  
 Herina: frondescentiae.  
 Hermetia: illucens, leucopa.  
 Hilara: globulipes.  
 Hoplitimyia: mutabilis, variabilis.  
 Hoplocheiloma: fabricii, fasciata,  
     totiana.  
 hortensia: hortulana.  
 hortona: icela.  
 hortorum: importuna.  
 hortulanus: hortulana.  
 hugonis: longipes.  
 hyalinata: hortulana.  
 Hybos: grossipes.  
 Hydromya: dorsalis.  
 Hydromyza: livens.  
 Hydrophoria: conica, dubitata, lancifer,  
     lenio.  
 Hydrotaea: armipes, ciliata, constane,  
     constans, curvipes, dentipes,  
     diabolus, glabricula, hirticeps,  
     ignavus, irritans, meteorica, spinipes,  
     vaccarum.  
 Hylemya: strigosa, vagans, variata,  
     volito.  
 hyoscyami: cinerea, leontodontis.  
 Hypoderma: ascarides.  
 Hypopygiopsis: fumipennis, infixa.  
 iavana: varia.  
 ibis: atherix.  
 ichneumonea: aristata.  
 idioidea: purpurea, violacea.  
 ieuna: favillacea, jeiuna, jejuna.  
 ignava: ignavus, pigra.

- Ilione*: albiseta, aratoria, crocus, oratoria.  
*illucens*: leucopa.  
*illustrata*: fulva, illustratus.  
*illustris*: equestris, muralis, parvula.  
*impavida*: concinna.  
*impuncta*: interpuncta, revolo, virescens.  
*inanis*: annulatus, apivora, micans,  
 trifasciata.  
*incana*: nemorum.  
*incisilobata*: striata.  
*incisuralis*: cuprea.  
*inferior*: gurnei, gurneyi.  
*ingruo*: ingrus.  
*intermedium*: costata.  
*interrupta*: fuscus, interruptus, lineolae,  
 obfuscata.  
*interruptella*: interrupta.  
*intricaria*: fuscus, intricata.  
*iridatus*: caerulescens, indicus, iridata.  
*Ischiodon*: nigra.  
*Isomerocera*: quadrilineata.  
*Isomyia*: chrysoides, confixa,  
 conflagrans, delectans, ditissima,  
 dotata, munda, reflectens, selecta,  
 terminata, viredaurea, viridaurea.  
*Iteuthelaira*: esuriens.  
*jamaicensis*: nigrana, nigrita.  
*Keroplatidae*: lineata, longicornis,  
 macroceras, striata.  
*labialis*: depressa.  
*labiata*: pallens.  
*lacustre*: lacustris.  
*laesus*: nigripes.  
*lancifer*: conica, dubitata.  
*larvarum*: erucarum, larvincola, major  
 larvarum, monachae.  
*larvipara*: corvinoides, mesopotamiensis,  
 vivipara.  
*lasciva*: lasciuia.  
*Lasiopa*: villosa.  
*lasiophthalma*: polita.  
*latifrons*: obsoleta.  
*lauta*: polita, proerna.  
*Lauxania*: chrysoptera, cylindricornis.  
*Lauxaniidae*: caeruleophthalmica,  
 chrysoptera, cylindricornis, deduco,  
 elevata, glabrata, longipennis,  
 lupulina, mulsus, quadripunctata.  
*Lejogaster*: metallina, politus.  
*Lejops*: femorata, interpunctus, transfuga.  
*lepidia*: parcus.  
*Lespesia*: marmorata.  
*leucocephala*: semiargentata,  
 semiargentea.  
*leucomelaena*: leucomelas.  
*Leucomyia*: cinerea.  
*Leucophora*: buccata, grisea.  
*Leucopis*: atratula.  
*Leucostoma*: aterrima, brevis.  
*Leucozona*: asiliformis, depressa,  
 glaucia, laternaria.  
*Leucozona*: lucorum.  
*leucozona*: lucozona.  
*levida*: assimilis, borealis, levidus.  
*Liancalus*: virens.  
*ligurriens*: fortunata, solaia.  
*Ligyra*: satyrus, sphinx.  
*limbipennis*: virens.  
*Limnia*: flavifrons, unguicornis, varicus,  
 varieus.  
*Limnophora*: notata, riparia, triangula.  
*lincensis*: longicornis.  
*lineata*: cinctus, leneatus, lineatus.  
*lineatus*: femorata.  
*Linnaemya*: investigator.  
*linogrisea*: lenio.  
*Lispe*: tentaculata.  
*Lispococephala*: verna.  
*Litorhina*: lar.  
*Lomatia*: sabaeus.  
*Lonchaea*: chorea, vernicina.  
*Lonchaeidae*: chorea, vernicina.  
*longicornis*: strigata, tenebricus.  
*longipes*: atra.  
*longula*: parvula.  
*lopesi*: pusilla.  
*Lophosceles*: mutata.  
*Loxocera*: aristata, ichneumonea, vagus.  
*lucidula*: africana.  
*Lucilia*: azurea, cadaverina, caerulea,  
 caerulescens, caesar, caesarina,  
 caesarion, cluvia, cuprina, equestris,  
 eximia, fucina, fulges, illustris,  
 inducta, inscribens, insularis, ionia,  
 lagyra, marginifera, metilia, muralis,  
 nigriceps, nitescens, nobilis, parvula,

- ponia, porphyrina, purpurascens,  
regalis, ruficeps, serenissima,  
sericata, silvarum, splendida,  
tegularia, temperata, tifata, varians,  
viridi-caerulea.
- lucorum: asiliformis.  
lugubris: dirae, moesta.  
lunatus: lunata.  
lurida: hirsuta.  
lusoria: kihuris, spinohumera, teste.  
lustrator: clemens.  
lutaria: comito.  
macellaria: certima, erythrocephala,  
fasciata, laniaria, phauda, tanaria,  
turbida.
- Macquartia: pubiceps.  
Macrocerca: longicornis.  
macrocera: longicornis.  
Macrocerca: macroceras.  
Macrorchis: meditata.  
maculata: compuncor, compunctus,  
vulpina.
- maculatus: maculata.  
maculipennis: arcuata.  
magellanica: incerta.  
mallochi: squamata.  
Mallota: fuciformis, fusiformis.  
marginata: cribrata, limbata, tristis.  
marginifera: marginalis.  
marmorea: flauescens.  
marsya: chrysotelus.  
Masicera: crassiseta.  
maurus: denigrata, hirsuta, maura.  
Medetera: diadema, rostrata.  
Medinophyto: dilecta.  
meditabunda: aluta, incurvata,  
meditalnenda, nora.  
megacephala: bata, combrea, dux,  
flaviceps, remuria.  
meigenii: annulata.  
Melanomya: minima.  
Melanophora: grossificationis,  
interventum, roralis.  
Melanostoma: facultas, mellina.  
Melia: crassipennis, gangraenosia.  
Melinda: anthracina, caerulea, cognata,  
gentilis.  
mellinum: facultas, mellina.
- meridionalis: curvipes.  
Merodon: bombyliformis, calcarata,  
clauda, clauipes, crassipes, curvipes,  
elephantopus, funesta, melancholica,  
natans, segetum.
- Meromacrus: cincta, pinguis, pratorum.  
Meromyza: clavicrus, minuta, pigna,  
saltatrix.
- Mesembrina: aparia, bombinatrix,  
bombylina, bombylius, bombyloides,  
discolor, meridiana, mystacea.
- Mesembrinella: bicolor.  
Mesoblattina: lithophila.  
Metallea: cuprea.  
metallina: politus.
- Metasyrphus: corollae.  
meteorica: constane, constans, vaccarum.  
Metopia: frontalis, labiata, leucocephala.  
Microchrysa: caesia, parvulus, polita,  
vitreus.
- Microdon: apiformis, devia, mutabilis,  
nova, viridescens.
- microleon: nigricans.
- Micropeza: corrugolata, filiformis,  
strumosa.
- Micropezidae: annulata, cibaria,  
corrugolata, cothurnata, ephippium,  
fabricii, fasciata, filiformis, lascivua,  
longipes, pedo, peragro, petronella,  
strumosa, totliana.
- Microtropesa: sinuata.
- Milesia: crabroniformis, naevia,  
semiluctifera, virginiensis.
- miliaria: arcuata.
- Minettia: longipennis, lupulina.
- Mintho: cincta, compressa, praeceps,  
rufiventris, rutillus.
- minuta: minor, minutior, pusillus.
- Morellia: affixa, basalis, bipuncta,  
calisia, hortensia, hortorum,  
hortulana, importuna, nitida,  
ochricornis.
- Morinia: doronici, melanoptera.
- morio: bicolora.
- mortuorum: chrysocephala.
- muliebris: pulchella.
- Musca: aethiops, afra, africana, alba,  
albina, albipennis, albomaculata,

alpesa, amica, amita, analis,  
 angustifrons, antiquissima, asiatica,  
 atrifrons, aurifacies, aurulans,  
 australia, australis, autumnalis,  
 bakeri, basilaris, beckeri, bezzii,  
 biseta, bivittata, bovina, calleva,  
 campestris, campicola, capensis,  
 carnifex, cassara, chilensis, chui,  
 cingalaisina, conducens, confiscata,  
 congolensis, conosoma,  
 consanguinea, contigua, continua,  
 convexifrons, coruina, corvina,  
 corvinoides, craggi, crassirostris,  
 cuprea, curviforceps, curvinervis,  
 cuthbertsoni, dasyops, dawsoni,  
 determinata, dichotoma, divaricata,  
 divisa, domestica, domestica major,  
 dorsomaculata, efflatouni, elatior,  
 ethiopica, eutaeniata, exalbida,  
 fasciata, fergusoni, fieldi, flavifacies,  
 flavinervis, flavipennis, fletcheri,  
 floralis, formosana, freedmani,  
 frontalis, fulvescens, fulvithorax,  
 gabonensis, gibsoni, greeni, grisella,  
 gurnei, gurneyi, gymnosomea,  
 harpyia, heidiae, hervei, hilli, hoi,  
 hottentota, hugonis, humilis,  
 illingworthi, incerta, inconstans,  
 indica, inferior, insignis, interrupta,  
 interruptella, irwinrossi, jacobsoni,  
 kasauliensis, kihuris, kweilinensis,  
 larvipara, lasiopa, lasiophthalma,  
 lateralis, latifrons, latiparafrons,  
 liberia, lindneri, lineata, longipes,  
 lothari, lucens, lucidula, ludifacies,  
 ludifica, lusoria, malaisei, mallochi,  
 mediana, melanura, meruensis,  
 mesopotamiensis, minima, minor,  
 minuta, modesta, multispina, munroi,  
 nana, nancauriensis, natalensis,  
 nebulo, negriabdomina, nevilli,  
 nigricans, nigricruris, nigrithorax,  
 niveisquama, obscura, oceanica,  
 osiris, ovipara, pampasiana,  
 patersoni, pattoni, pellucens,  
 phasiaeformis, pilifacies, pilosa,  
 planiceps, polita, pollinosa, praecox,  
 prashadii, primitiva, prisca, promisca,

prostrata, pseudocorvina, pulla,  
 pumila, pungoana, pusilla, rau,  
 restituo, restituo, riparia, rivularis,  
 rufifrons, rufiventris, rustica,  
 sanctae-helenae, santoshi, scapularis,  
 scatophaga, senegalensis,  
 seniorwhitei, setigera, setulosa,  
 shanghaiensis, somalorum, sorbens,  
 sordidissima, soror, spangleri,  
 spectanda, speculifera, spinohumera,  
 spinosa, splendens, splendida,  
 squamata, stimulans, stomoxidea,  
 striacta, stuckenbergi, sugillatrix,  
 taitensis, tau, tempestatum,  
 tempestiva, terraereginae, teste,  
 tiberina, tibetana, tibiseta,  
 transvaalensis, ugandae, umbraculata,  
 unisetula, vaccina, vagatoria,  
 ventrosa, vetustissima, vicaria, vicina,  
 villeneuvii, violacea, vitripennis,  
 vivipara, wittweri, xanthomela,  
 xanthomelas, yerburyi.

*muscaria?*: alni.

*Muscidae*: adumbrata, aenea, aerea,  
 aethiops, affixa, afra, africana,  
 agromyzina, alba, albina, albipennis,  
 albolineata, albomaculata, alpesa,  
 aluta, amica, amita, amoena, analis,  
 angelicae, angustifrons, antiquissima,  
 apriaria, aricioides, armipes, asiatica,  
 assimilis, atra, atricolor, atrifrons,  
 aurifacies, aurulans, australis,  
 australis, autumnalis, bakeri, basalis,  
 basilaris, beckeri, benedicta, bezzii,  
 bipuncta, biseta, bivittata,  
 bombinatrix, bombylina, bombylius,  
 bombyloides, borealis, bovina,  
 braesia, caesarion, caesia, calcitrans,  
 calisia, calleva, campestris,  
 campicola, capensis, carnifex,  
 cassara, celsus, chalybea, chilensis,  
 chloris, chrysotelus, chui, ciliata,  
 cinerascens, cingalaisina, coeo,  
 coeruleata, coerulea, compuncitor,  
 compunctus, concolor, conducens,  
 confinis, confiscata, congolensis,  
 connexa, conosoma, consanguinea,  
 consimilis, constane, constans,

contigua, continua, convexifrons,  
corni, cornicena, cornicina, cornuta,  
coruina, corvina, corvinoides, craggi,  
crassirostris, cuprea, curviforceps,  
curvinervis, curvipes, cuthbertsoni,  
cyanea, cyanella, dasyops, dawsoni,  
decolor, deduco, dentipes, depuncta,  
determinata, diabolus, diaphana,  
dichotoma, diffidens, discolor,  
dispar, distincta, divaricata, divisa,  
domestica, domestica major, domitor,  
donysa, dorsomaculata, efflatouni,  
elatior, electa, erinacea, erratica,  
ethiopica, eustolia, eutaeniata,  
evectionis, exalbida, extranea, extrema,  
fasciata, fergusoni, fector, fieldi,  
flavicornis, flavifacies, flavinervis,  
flavipennis, fletcheri, floralis,  
formosana, freedmani, frontalis,  
fulvescens, fulvithorax, fuscata,  
fuscipennis, gabonensis, gavisa,  
geniculata, gibsoni, glabricula,  
greeni, grisea, grisella, guineensis,  
gurnei, gurneyi, gymnosomea,  
hamata, harpyia, heidiae, herea,  
hervei, hilli, hirticeps, hoi, hortensia,  
hortorum, hortulana, hottentota,  
hugonis, humilis, hyalinata, idessa,  
ignavus, illingworthi, importuna,  
impuncta, inanis, incerta, inconstans,  
incurvata, indica, inferior, ingredior,  
insignis, intermedia, interpuncta,  
interrupta, interruptella, intrahens,  
irritans, irwinrossi, jacobsoni,  
kasauliensis, kihuris, kweilinensis,  
laeta, lardaria, larvipara, lasiopa,  
lasiophthalma, lateralis, latifrons,  
latiparafrons, lauta, leucomelas,  
levidus, liberia, lindneri, lineata,  
litorea, longipes, lothari, lucens,  
lucidula, lucorum, ludifacies,  
ludifica, lusoria, luteipennis,  
maculata, malaisci, mallochi, mano,  
mantes, mantos, marginalis, maronea,  
marsya, mazaca, mediana,  
meditabunda, meditalnenda,  
meditata, meridiana, meruensis,  
mesopotamiensis, meteorica, minima,

minor, minuta, modesta, mollicula,  
multispina, munroi, mutata,  
mystacea, nana, nancauriensis,  
natalensis, nebulosa, negriabdomina,  
nemorum, nevilli, nigricans,  
nigricolor, nigricruris, nigrita,  
nigrithorax, nitida, niveisquama,  
nora, notata, nugater, nugator,  
obscura, obtrusa, obtusipennis,  
occidentis, oceanica, ochricornis,  
optata, ornata, osiris, ovipara,  
pabulorum, pagana, pallida,  
pampasiana, parviceps, pascuorum,  
patersoni, pattoni, pedella, pellucens,  
perfixa, phara, phasiaeformis,  
pilifacies, pilosa, planiceps, polita,  
pollinosa, praecox, praedicens,  
prashadii, pratorum, primitiva,  
princeps, prisca, prodeo, proerna,  
proferens, prolapsa, prolapsor,  
promisca, prostrata, pseudocorvina,  
puella, pulla, pumila, pungens,  
pungoana, pusilla, quadrimaculata,  
quadrimaculella, quadrum, racilia,  
rapax, rau, redeo, refixa, restituto,  
restituo, reversio, revolo, riparia,  
rivularis, rufifrons, rufiventris,  
ruralis, rustica, sanctae-helenae,  
santoshi, sarsina, scapularis,  
scatophaga, scutellaris, senegalensis,  
senilis, seniorwhitei, seplasia, serena,  
setigera, setulosa, sexmaculata,  
shanghaiensis, signata, somalorum,  
sorbens, sordidissima, soror,  
spangleri, spectanda, speculifera,  
sperata, spinipes, spinohumera,  
spinosa, splendens, splendida,  
squamata, stabulans, stimulans,  
stipata, stomoxidea, striacta,  
stuckenbergi, sublatus, subpuncta,  
subventus, sugillatrix, tahitiensis,  
taitensis, tau, tempestatum,  
tempestiva, tentaculata, terraereginae,  
testacea, teste, tiberina, tibetana,  
tibialis, tibiseta, tigrina,  
transvaalensis, triangula, triquetra,  
trita, tuguriorum, ugandae, uliginosa,  
umbraculata, unisetula, vaccarum,

vaccina, vagans, vagatoria, validus,	Neurigona: quadrifasciata.
variabilis, ventrosa, verna, versicolor,	niger: nigra, rugosa.
vespertina, vetustissima, vicanus,	nigricans: albicincta.
vicaria, vicina, vilis, villeneuvii,	nigricosta: nitida.
violacea, virescens, vitripennis,	nigrifrons: variabilis.
vivipara, vulpina, wittweri,	nigrina: lateralis, nigripes.
xanthomela, xanthomelas, yerburyi.	nigripes: pinivorae.
Muscina: assimilis, borealis, caesia,	nitidula: cylindrica.
cinerascens, levidus, publorum,	nobilitata: hirta, nobilis.
pascuorum, prodeo, prolapsa,	nobilitatus: ciliata, joco, ludicrous,
prolapsor, stabulans, tibialis.	ludicus, nobilitata.
mutabilis: variabilis.	noctiluca: erratica, tristor.
mutatus: mutata.	notata: nectarea, nectarivora.
Myathropa: atropos, florea.	notatus: notata.
Mydaea: corni, pagana, princeps, rustica.	Notogramma: stigma.
Mydas: clavata, filata.	novemmaculatus: novemmaculata,
Mydidae: clavata, filata.	tuberculata.
Myennis: octopunctata.	nubila?: cursoria.
Myiophaea: aricioides.	nubipennis: segmentaria.
Myiopharus: costalis.	nudibasis: resinellae.
Myoleja: caesio, lychnidis.	Nyctia: halterata, maura, nigrita.
Myolepta: luteola, potens.	obesa: nebulosa.
Myopina: myopina.	obscurata: mesembrinoides.
Myospila: aluta, incurvata, meditabunda,	Ochthera: manicata, mantis.
meditalnenda, nora.	Ocyptamus: cylindracea, cylindricus.
mystacea: aparia, bombinatrix,	Odontomyia: flavissima, hydroleon,
bombylina, bombylius, bombyloides,	microleon, nigricans.
discolor.	Oestridae: ascarides.
nana: minima.	Ogcodes: gibbosa.
natans: melancholica.	Onesia: germana, hederacea,
nebulosa: ornata.	mesembrinoides, micans, obscurata,
Nemopoda: cylindrica.	ruficornis, sepulcralis.
Nemotelus: albipes, marginata,	Oplodontha: jejuna, minor, minuta,
marginella, pantherina, uliginosa.	minutior, pusillus, viridula.
Neoascia: annexa, bifasciata, dizonias,	Opomyza: florum, germinationis.
meticulosa, molio, podagriva, tenur.	Opomyzidae: combinata, florum,
Neomyia: amoena, benedicta, braesia,	galpinus, germinationis.
caesarion, chalybea, chloris,	Opsdexia: serva.
coerulata, coerulea, connexa,	Opsozelia: lateralis.
cornicina, cornicina, cyanea,	Orellia: falcata, punctata, stictica.
diffidens, donysa, electa, gavisa,	Ormia: punctata.
guineensis, heraea, indica, ingredior,	Ornidia: obesa.
intrahens, lauta, maronea, mazaca,	Orthoceratium: lacustris.
obtrusa, optata, perfixa, phara, polita,	Orygma: palpator.
proerna, puella, racilia, redeo, refixa,	Oscinella: avenae, frit, hordei, nigra.
sarsina, sperata, trita.	Osiris: fulvithorax, sugillatrix.
Neria: ephippium.	Oties: formosa, lamed, porca.
Neta: chilensis.	

- Otitidae: *aenea*, *alceae*, *annonae*,  
*aquatica*, *costalis*, *crassipennis*,  
*demandata*, *formosa*, *frondescentiae*,  
*gangraenosa*, *graminum*, *hortulana*,  
*hyalinata*, *lamed*, *myennis*,  
*octopunctata*, *picta*, *porca*, *stigma*,  
*urticae*, *vibrans*.
- Oxycera: *graeca*, *hypoleon*, *muscaria*,  
*rara*, *tardigradus*, *trilineata*.
- Oxyna: *parietina*.
- Oxysarcodexia: *auriflua*.
- Palloptera: *muliebris*, *pulchella*, *saltuum*,  
*umbellatarum*, *unicolor*.
- Pallopteridae: *minutus*, *muliebris*,  
*pulchella*, *saltuum*, *umbellatarum*,  
*unicolor*.
- Palpada: *agrorum*, *hortorum*,  
*surinamensis*.
- pandia: *unxia*.
- pantherinus: *albipes*, *marginata*,  
*marginella*, *pantherina*.
- panzeri: *conica*.
- papua: *calliphoroides*.
- papuensis: *azurea*, *inducta*, *inscribens*,  
*marginifera*, *metilia*, *nitescens*, *tifata*.
- Paragus: *cruentata*, *pusilla*, *thymastri*.
- parallelus: *parrallelus*.
- Paramenia: *macularis*.
- Paraprissopoda: *gulo*.
- Parasarcophaga: *ruficornis*.
- Paregle: *audaculus*.
- Parhelophilus: *frutetorum*, *versicolor*,  
*versicolorata*.
- Paroxyna: *cinereus*, *cinerius*.
- pattoni: *incerta*, *spinosa*, *yerburyi*.
- pedissequum: *pedissequus*.
- Pegomya: *auctus*, *bicolor*, *conformis*,  
*flavipes*, *fungorum*, *laticornis*, *rufina*,  
*rufipes*, *ruminicis*, *testacea*, *transversa*.
- Pelidnoptera: *nigripennis*.
- pellucens: *pellita*.
- pellucens?: *deceptoria*.
- peltatus: *timeo*.
- pendulus: *pendula*, *trelineata*, *trileniata*,  
*trilenva*, *trilineata*.
- penicillata: *versicolor*.
- perexilis: *perexillis*.
- Pericallimyia: *perlata*.
- Periscepsia: *carbonaria*.
- pernix: *haemorrhoidalis*, *striata*.
- peronii: *braesia*, *cyanea*, *mazaca*, *phara*,  
*sarsina*.
- personata: *apiformis*, *personatus*.
- pertinax: *lunula*.
- Peteina: *erinaceus*.
- petiolata: *compressa*.
- Phaeomyiidae: *nigripennis*.
- Phaonia: *angelicae*, *erinacea*, *erratica*,  
*extranea*, *fuscata*, *laeta*, *nemorum*,  
*nugater*, *nugator*, *pallida*, *rufiventris*,  
*scutellaris*, *sublatus*, *subventus*,  
*testacea*, *tuguriorum*, *vagans*, *validus*.
- Phasia: *brachyptera*, *dimidiata*, *taeniata*.
- Pherbellia: *punctata*.
- Pherbina: *coryleti*, *coryletti*, *reticulata*.
- Phlebotomus: *papatasi*.
- Phora: *aterrima*, *festinans*.
- Phoridae: *aterrima*, *festinans*, *florea*,  
*florescens*, *subsultans*.
- Phormia: *accincta*, *bicolor*, *mollis*,  
*proxima*, *regina*, *thalassina*.
- Phumosia: *dioclea*, *promittens*.
- Phyllomya: *volvulus*.
- Phyllomyia: *cylindrica*.
- Physiphora: *aenea*, *alceae*, *demandata*.
- Phytomia: *crassa*, *errans*, *sinensis*,  
*zonalis*, *zonatus*.
- picea: *piniperdae*.
- Pierretia: *sexpunctata*.
- pilifacies: *hoi*.
- pilosus: *hystrix*, *pilosa*.
- pinastri: *consisto*, *pyrorum*.
- pinguis: *cincta*.
- Piophila: *atrata*, *casei*.
- Piophilidae: *atrata*, *casei*, *cynophila*,  
*furcata*.
- Piophilidae?: *ornithovora*.
- pipiens: *pipines*.
- Pipiza: *dirae*, *erratica*, *lugubris*, *moesta*,  
*noctiluca*, *tristor*.
- Pipizella: *viduata*.
- Pipunculidae: *cephalotes*.
- Pipunculus: *cephalotes*.
- planiceps: *chui*, *cingalaisina*, *indica*,  
*pollinosa*.
- Platycephala: *planifrons*, *umbraculata*.

- Platycheirus*: agilitas, albimana,  
 confusus, cyanea, dexter, granditarsa,  
 navus, ocymi, rosarum, timeo, vivax.  
*Platypalpus*: cursitans.  
*Platyparea*: discoidea, poeciloptera.  
*Platypeza*: palmipes, properans.  
*Platypezidae*: palmipes, properans.  
*Platystoma*: fulviventris, seminationis,  
 variolosa.  
*Platystomatidae*: fulviventris, lugens,  
 moerens, naevosus, seminationis,  
 singenesiae, syngenesiae, variolosa.  
*Platypteresa*: opulenta.  
*plebeja*: fasciata.  
*plumicornis*: lineata.  
*plumipes*: gibba.  
*pluvialis*: litus.  
*Pocota*: apiformis, personatus.  
*podagrifica*: molio.  
*Poecilanthrax*: lucifer.  
*Poecilobothrus*: ciliata, joco, ludicrous,  
 ludicus, nobilitata.  
*Poecilometopa*: spilogaster.  
*Polistes*: albolineata, domitor, lardaria.  
*polita*: caesia, parvulus, vitreus.  
*Pollenia*: amentaria, atramentaria,  
 bicolor, caerulescens, cinerea,  
 degener, depressa, familiaris,  
 flavigans, floralis, fulvicornis,  
 fulvipalpis, intermedia, lanio,  
 leucoptera, macellaria, nana, nigrina,  
 nitens, obscura, olivacea, pruinosa,  
 pumila, recta, recurvo, remigro, rufis,  
 ruficeps, ruficornis, rufipalpis,  
 umbrifera, vagabunda, varensis,  
 varia, varians, virescens.  
*Polleniopsis*: inconclusa, patiens,  
 sarcophagooides.  
*pomeroyi*: fasciata.  
*postica*: heraclei.  
*posticus*: elongata.  
*praeceps*: cincta, rutillus.  
*profuges*: conopsea, profugus.  
*prolapsa*: caesia, pabulorum, prolapsor.  
*Protocalliphora*: azurea, carnarida,  
 chrysorrhoea, pervenio, reviso,  
 sordida, violacea.  
*Protophormia*: boganidae, groenlandica,  
 orior.  
*Psarus*: abdominalis, coalescens.  
*Pseudogametes*: semiatra.  
*Psila*: fimetaria, flava, rufa.  
*Psilidae*: aristata, extenuata, fimetaria,  
 flava, histrionica, ichneumonea,  
 leptogaster, longerro, longervo,  
 luteus, percussus, rosae, rufa, vagus.  
*Psilidae?*: rufa.  
*Psychodidae*: papatasi.  
*Ptecticus*: elongata, japonica, testaceus.  
*pubera*: puber.  
*pubescens*: ruficornis.  
*pubicornis*: dolens, puberula.  
*pulchra*: phellia.  
*pulchripes*: flavipes.  
*pumilionis*: lineata, nasuta, saltatrix,  
 umbelliferarum.  
*punctatus*: bilineata.  
*punctella*: cinereus, cinerius.  
*punctulata*: hieracii.  
*punctum*: stigma.  
*purmundus*: permundus.  
*putris*: conssecnis, cossenico, fmeti.  
*Pycnomalla*: splendens.  
*pyrastri*: mellina, rosae.  
*Pyrellia*: aenea, coeo, extrema,  
 fuscipennis, luteipennis, parviceps,  
 proferens, rapax, serena.  
*Pyrellina*: chrysotelus, distincta, marsya,  
 obscura.  
*quadrimaculata*: fulvimaculata, sacra,  
 violacea.  
*quadrup*: subpuncta.  
*quinquelineatus*: quinquelineata.  
*quinquestriatus*: quinquestriata.  
*radicum*: napobrassicae.  
*rapax*: aenea, coeo, fuscipennis,  
 luteipennis, parviceps, serena.  
*rara*: tardigradus.  
*Ravinia*: haemorrhoidalis, moneta,  
 pernix, striata.  
*regina*: accincta, bicolor, mollis,  
 proxima, thalassina.  
*Rhagio*: annulata, lineola, maculata,  
 scolopacea, tringaria, vanella.

- Rhagionidae: annulata, asiliformis, cingulata, cristata, fasciatus, lineola, maculata, norwegica, scolopacea, tomentosa, tringaria, vanella.
- Rhagionidae?: acuminata.
- Rhagoletis: cerasi.
- Rhamphinina: pica.
- Rhingia: macrocephala, nasatus, nosatus, rostrata.
- Rhinophora: parcus.
- Rhinophoridae: atra, grossificationis, interventum, parcus, roralis.
- Rhyncomya: collecta, concinna, divisa, felina, impavida, ruficeps.
- ribesii: blandus, vacua.
- Rivellia: singenesiae, syngenesiae.
- rognesi: carnarida, chrysorrhoea, sordida.
- rohrelliformis: inanis.
- rroralis: grossificationis, interventum.
- rotundatum: cerinus, hemisphaerica, rotundata.
- rudis: caerulescens, familiaris, floralis, fulvicornis, fulvipalpis, nana, obscura, olivacea, pruinosa, pumila, recta, remigro, rufipalpis, varia.
- rudis?: torvis, torvus.
- rufifacies: orientalis.
- rufiventris: nugater, nugator, scutellaris, testacea.
- rufomaculata: demano, dimano, lateralis.
- rustica: plebeja, provenio.
- rustica?: germanica.
- rutilans: valida.
- Rutilia: formosa, liris, regalis, retusa, splendida.
- saltatrix: clavicus, minuta, pigra.
- Saltella: spondylii.
- saltuum: unicolor.
- Sapromyza: caeruleophthalmica, quadripunctata.
- Sarconesia: incerta.
- Sarcophaga: carnaria, colossus, prolabor, reverto, serpo, valens, variegata, vivipara major.
- Sarcophagidae: affinis, africa, auriflua, carnaria, cinerea, colossus, frontalis, gulo, haemorrhoa, haemorrhoidalalis, halterata, hauriens, labiata, latifrons, leucocephala, lineata, martia, maura, moneta, nigrita, obsoleta, occidua, pernix, prolabor, pusilla, reverta, ruficornis, rufifrons, ruralis, serpo, sexpunctata, spilogaster, striata, tabaniformis, taenionota, tessellata, valens, variegata, vivipara major.
- Sarcophila: latifrons, obsoleta.
- Sargus: bipunctata, caerulescens, capraria, cupraria, indicus, iridata, violacea.
- Scaeva: mellina, pyrastri, rosae.
- scalaris: alevo, allevo, minor domestica.
- Scathophaga: comito, exilis, lucophaeus, lucopheus, lataria, merdaria, scybalaria, stercoraria, suilla.
- Scathophagidae: abbreviata, cloacaris, comito, exilis, livens, lucophaeus, lucopheus, lataria, merdaria, paganus, puber, pubera, scybalaria, stercoraria, suilla, tipularia.
- Scatopse: nectarea, nectarivora.
- Scatopsidae: nectarea, nectarivora.
- Scellus: notata.
- Scenopinidae: albicincta, fenestratalis, nigra, rugosa, saltitans, senilis, spoliata, tarda.
- Scenopinus: albicincta, fenestratalis, nigra, rugosa, saltitans, senilis, spoliata, tarda.
- schoenherri: punctata.
- Sciapus: ingruo, ingrus.
- Sciaridae: maura, morio.
- Sciomyzidae: alata, albisetata, aratoria, chaerophylli, cincta, coryletti, coryletti, cibrata, crocus, cucularia, delicate, dorsalis, elata, elater, flavifrons, flavipes, fumigata, hieracii, limbata, marginata, oratoria, punctata, punctulata, reticulata, rufa, rufifrons, rufipes, sollicitus, sphegea, spinipes, tener, tenur, tristis, umbrarum, unguicornis, varicus, variegata, varieus.
- scita: scitus.
- scolopaceus: scolopacea.
- Scotiaptera: venatoria.
- scripta: inviso, molitus.

- Scutellomusca*: *scutellaris*.  
*scybalaria*: *lucophaeus*, *lucopheus*.  
*seelandicus*: *seelandica*, *trilineatus*.  
*segnis*: *contracta*, *fucatus*, *maritima*,  
*melanochrysa*.  
*Scioptera*: *urticæ*, *vibrans*.  
*seminicinereus*: *hyalinata*.  
*semiluctifera*: *naevia*.  
*seminationis*: *fulviventris*, *variolosa*.  
*senegalensis*: *luteola*.  
*seniorwhitei*: *tibiseta*.  
*Senogaster*: *dentipes*.  
*Sepedon*: *flavipes*, *rufipes*, *sphegea*,  
*spinipes*.  
*Sepsidae*: *cosssencis*, *cossenico*,  
*cylindrica*, *cynipsea*, *elongata*, *fimeti*,  
*minimus*, *palpator*, *punctum*, *putris*,  
*spondylii*, *stigma*, *vibrans*.  
*Sepsis*: *cynipsea*, *elongata*, *punctum*,  
*stigma*, *vibrans*.  
*sepulchralis*: *ater*, *melanius*.  
*seriata*: *rufifrons*.  
*sericata*: *caerulescens*, *lagyra*, *nobilis*,  
*tegularia*.  
*sericea*: *buccata*.  
*Sericomyia*: *lappona*, *silentis*,  
*subtomentosa*.  
*serrata*: *latrinarum*.  
*serratulae*: *serratula*.  
*Servillia*: *argyrostoma*, *hirsuta*, *lurida*.  
*setipyga*??: *collecta*.  
*setiventris*: *cornuta*.  
*sexmaculata*: *deduco*, *uliginosa*.  
*Silbomyia*: *fuscipennis*, *prospera*.  
*silentis*: *subtomentosa*.  
*silvatica*: *cinerea*.  
*Simuliidae*: *colombaschenensis*.  
*Simulium*: *colombaschenensis*.  
*singularior*: *singularius*.  
*Siphona*: *geniculata*, *urbanus*.  
*solennis*: *rumicis*.  
*solicita*: *rufa*, *solicitus*.  
*solivaga*: *solivagus*.  
*solstitialis*: *arctii*, *dauci*.  
*sonchi*: *subcutanea*.  
*sorbens*: *alba*, *bivittata*, *dichotoma*,  
*efflatouni*, *eutaeniata*, *exalbida*,  
*humilis*, *mediana*, *niveisquama*,  
*primitiva*, *promisca*, *pusilla*,  
*spectanda*, *stuckenbergi*.  
*spерата*: *electa*.  
*Sphaeroceridae*: *saniosa*.  
*Sphaerophoria*: *invisito*, *menthastri*,  
*molitus*, *scripta*.  
*sphegeus*: *flavipes*, *rufipes*, *sphegea*.  
*sphingivora*: *crassiseta*.  
*sphondylii*: *spondylii*.  
*Spilogona*: *aerea*, *dispar*, *litorea*,  
*marginalis*.  
*Spilomyia*: *diophthalma*.  
*spiloptera*: *grisea*.  
*spissa*: *aricioides*.  
*splendens*: *splendida*.  
*stabulans*: *cinerascens*, *prodeo*, *tibialis*.  
*Stegana*: *coleoptrata*, *furta*.  
*steini*: *fungorum*.  
*stercoraria*: *exilis*, *merdaria*.  
*Sterphus*: *jamaicensis*, *nigrana*, *nigrita*.  
*stictica*: *punctata*.  
*stigma*??: *placida*.  
*Stilbomyella*: *costalis*, *diffusa*, *gloriosa*.  
*Stomina*: *tachinoides*.  
*Stomorrhina*: *discolor*, *lunata*.  
*Stomoxys*: *calcitrans*, *occidentis*,  
*pungens*.  
*Stratiomyidae*: *aequinoctialis*, *albipes*,  
*aurata*, *bipunctata*, *caerulescens*,  
*caesia*, *capraria*, *chalybata*, *chalybea*,  
*chalybeata*, *chamaeleon*, *cicur*,  
*clavipes*, *coccinea*, *cupraria*, *elongata*,  
*ephippium*, *flavissima*, *formosa*,  
*graeca*, *hydroleon*, *hypoleon*,  
*illucens*, *inda*, *indicus*, *iridata*,  
*japonica*, *jejuna*, *leucopa*, *marginata*,  
*marginella*, *melanopis*, *microleon*,  
*minor*, *minuta*, *minutior*, *muscaria*,  
*mutabilis*, *nigricans*, *pallipes*,  
*pantherina*, *parvulus*, *polita*, *pusillus*,  
*quadrilineata*, *rara*, *rex*, *sellata*,  
*sexdentata*, *similis*, *singularius*,  
*spatula*, *splendens*, *strigata*,  
*tardigradus*, *tenebricus*, *testaceus*,  
*trilineata*, *uliginosa*, *vallata*,  
*variabilis*, *villosa*, *violacea*, *viridula*,  
*vitreus*.

- Stratiomys*: *chamaeleon*, *pallipes*,  
*singularius*, *spatula*, *strigata*,  
*tenebricus*.  
*striatus*: *lineata*, *striata*.  
*strigatus*: *lineata*.  
*Strongygaster*: *lingens*.  
*stygia*: *laemica*.  
*stylata*: *iacobaeae*, *jacobaeae*.  
*Subhylemyia*: *longula*, *parvula*.  
*subventa*: *subventus*.  
*Suillia*: *tuberis*.  
*superbiens*: *fulvus*, *mussitanus*.  
*sylvana*: *sylvanus*.  
*sylvarum*: *longisco*.  
*Sylvicola*: *bilineata*, *cinctus*, *fuscata*,  
*nigricans*, *punctatus*, *succincta*.  
*syngenesiae*: *singenesiae*.  
*Syntormon*: *pallipes*.  
*Syritta*: *pipiens*, *pipines*.  
*Syrphidae*: *abdominalis*, *ablectus*,  
*adunata*, *aenea*, *agilitas*, *agrorum*,  
*albibimana*, *alternata*, *americana*,  
*annexa*, *annulatus*, *anteaimbulo*,  
*anteambulo*, *aphidivora*, *apiformis*,  
*apivora*, *arbustorum*, *arcuata*,  
*aruorum*, *arvorum*, *asiliformis*, *ater*,  
*atropos*, *balteata*, *bardus*, *bicincta*,  
*bicinctata*, *bifasciata*, *bipunctata*,  
*blandus*, *bombylans*, *bombyliformis*,  
*brunnea*, *caemeteriorum*, *calcarata*,  
*callosus*, *cannabina*, *cautus*,  
*cemiteriorum*, *chrysorrhaea*,  
*chrysozonias*, *cincta*, *cinctus*,  
*cingulatus*, *citrofasciata*,  
*citrofasciata*, *clauda*, *clauipes*,  
*coalescens*, *comitus*, *confusus*, *conica*,  
*conopsea*, *conopsoides*, *consisto*,  
*contracta*, *corollae*, *coryden*, *corydon*,  
*crabroniformis*, *crassa*, *crassipes*,  
*cruentata*, *cryptarum*, *curvipes*,  
*cyanea*, *cylindracea*, *cylindricus*,  
*dentipes*, *depressa*, *devia*, *dexter*,  
*diophthalma*, *dirae*, *dispar*, *dizonias*,  
*elegans*, *elephantopus*, *eligans*,  
*elongata*, *ericotorum*, *errans*, *erratica*,  
*esuriens*, *exinanita*, *facultas*, *fallax*,  
*fasciata*, *fasciolata*, *femorata*, *festiva*,  
*flavescens*, *flavicornis*, *florea*,  
*formosus*, *frutetorum*, *fucatus*,  
*fuciformis*, *fulva*, *fulvus*, *funebres*,  
*funebris*, *funesta*, *funibres*, *fucus*,  
*fusiformis*, *glaucia*, *granditarsa*,  
*hirsutissima*, *horticola*, *hortorum*,  
*hortulana*, *illustratus*, *imbellis*,  
*inaequalis*, *inanis*, *inflata*,  
*interpunctus*, *interrupta*, *interruptus*,  
*intersecta*, *intricaria*, *intricata*,  
*invisito*, *jamaicensis*, *laeta*, *lappona*,  
*laternaria*, *leneatus*, *leucocephala*,  
*libatrix*, *lineata*, *lineatus*, *lineolae*,  
*lineolata*, *longipes*, *longisco*, *lucida*,  
*lucorum*, *lugubris*, *lunata*, *lunula*,  
*luteola*, *lyra*, *macrocephala*,  
*macrophthalma*, *maritima*,  
*melancholica*, *melanius*,  
*melanochrysa*, *melanopyrrha*,  
*mellina*, *menthastri*, *merulina*,  
*metallina*, *meticulosa*, *micans*,  
*moesta*, *molio*, *molitus*, *monoculus*,  
*murina*, *murorum*, *mussitanus*,  
*mutabilis*, *naevia*, *nasatus*, *natans*,  
*navus*, *nectareus*, *nectarina*,  
*nemorum*, *nigra*, *nigrana*, *nigrita*,  
*nitens*, *noctiluca*, *nosatus*, *nova*,  
*obesa*, *obfuscata*, *ochroleuca*, *ocymi*,  
*oestracea*, *pacifica*, *palustris*,  
*paralleli*, *parallelus*, *parralleli*,  
*parrallelus*, *parrellei*, *pedissequus*,  
*pellucens*, *pendula*, *perexlimis*,  
*pereillis*, *personatus*, *pigra*, *pinastri*,  
*pinguis*, *pipiens*, *pipines*, *plumata*,  
*plumosa*, *podagraca*, *politus*, *porcina*,  
*potens*, *prasina*, *pratorum*, *profuges*,  
*profugus*, *pulla*, *punctata*, *pusilla*,  
*pyrastri*, *pyrorum*, *quadrilineatus*,  
*quinquelineata*, *quinquestriata*,  
*ribesii*, *rosae*, *rosarum*, *rostrata*,  
*rubiginea*, *rubiginosa*, *ruficaudis*,  
*ruficornis*, *rutilo*, *salviae*, *scitule*,  
*scitulus*, *scitus*, *scripta*, *seelandica*,  
*segetum*, *segnis*, *semiluctifera*,  
*semirufa*, *semulater*, *semulator*,  
*sepulchralis*, *silentis*, *simulator*,  
*sinensis*, *speculifera*, *subtomentosa*,  
*surinamensis*, *sylvarum*, *tarditas*,  
*tartidus*, *tenax*, *tenur*, *tetragona*,

thymastri, timeo, tranquebarica,  
transfuga, trelineata, trifasciata,  
trileniata, trilenva, trilineata,  
trilineatus, tristis, tristor, univalvis,  
vacua, valentina, valga, versicolor,  
versicolorata, vesiculosa,  
vespaesimilis, vespiformis, viduata,  
virginiensis, viridescens, vivax,  
vorax, zonalis, zonaria, zonatus.

*Syrphus*: blandus, ribesii, vacua.

*Tabanidae*: pellita.

*Tachina*: erucarum, fera, grossa,  
neustriae, reccumbo, rerumbo,  
tessella, tessellata.

*Tachinidae*: acromion, affinis, agilis,  
albifrons, aldrichi, altratus, anceps,  
angusticornis, argyrostoma, aterrima,  
atratus, atrifrons, aurifrons, ausus,  
bifasciata, bohemica, brachyptera,  
brassicaria, brevis, canina, canus,  
capitata, carbonaria, carinifrons,  
cerasina, cerinus, chrysostoma,  
cincta, cingulata, compressa, conflujo,  
conspersus, continua, convenio,  
convolo, costalis, costata, crassa,  
crassicornis, crassipennis, crassiseta,  
cylindrica, deceptoria, demano,  
diaphana, dilecta, dimano, dimidiata,  
erinaceus, erucarum, esuriens,  
evonymellae, fasciata, fera, ferina,  
flaviceps, formosa, gastrodes,  
geniculata, germanica, globosa,  
grisescens, grossa, hartigii, helluo,  
hemiptera, hemisphaerica, hirsuta,  
hirta, hystrix, iliaca, interpunctionis,  
interventus, investigator, larvarum,  
larvincola, lateralis, leucozona,  
libatrix, lingens, liris, longipes,  
lucozona, lupina, lurida, major  
larvarum, marmorata, metallina,  
metus, minor larvarum, mirabilis,  
moesta, monachae, nebulosa,  
neustriae, nigricornis, nigrina,  
nigripes, obsidianus, ochracea,  
parasitica, pavida, pennipes, petiolata,  
phalaenarum, pica, pilosa, pinguis,  
piniperdae, pinivorae, plebeja,  
praeceps, procedo, processionae,

promano, promans, provenio,  
pubiceps, punctata, puparum,  
quadripustulata, reccumbo, recedans,  
redambulo, redeans, redians, regalis,  
repens, rerumbo, resinellae, retusa,  
rotundata, ruficrus, rufiventris,  
rufomaculata, rustica, rutilans,  
rutillus, sensifera, sericea, sinuata,  
solivagus, splendida, subcoleoptrata,  
tachinoides, taeniata, temula, tessella,  
tessellata, torvis, torvus, tremula,  
tricincta, tristis, urbanus, vacua,  
valida, varia, venatoria, villica,  
volets, volvulus.

*Tachinidae*?: nigripes, rustica.

*Tachydromia*: arrogans.

*Tachypeza*: cursoria.

*Taenaptera*: annulata, lasciua.

*taeniatum*: gibba.

*tasmaniae*: extrema.

*Temnostoma*: vespaesimilis, vespiformis.  
*tempesta*: albipennis, carnifex, cuprea,  
curvinervis, minima, nana,  
phasiaeformis, pusilla, stimulans,  
unisetula, violacea.

*temula*: obsidianus, tremula.

*tenax*: apiformis, porcina.

*Tephritidae*: amica, arctii, arcuata,  
arnicae, artemisiae, caesio, cardui,  
centaureae, cerasi, cinerea, cinereus,  
cinerius, connexa, cornuta, coryli,  
crax, crux, dauci, discoidea,  
erythrocephala, falcata, ferruginea,  
flauescens, florescentiae, fucata,  
furcata, haemorrhoidalis, helianthi,  
heraclei, heraclii, hyoscyami,  
iacobeae, jacobae, lappae,  
leontodontis, limatus, lincensis,  
linzensis, longicornis, lychnidis,  
miliaria, oleae, onopordinis, parietina,  
perelegand, perelegans, permundus,  
placida, poeciloptera, punctata,  
purmundus, radiata, ruficauda,  
scabiosae, serratula, serratulae,  
solstitialis, sonchi, stellata, stictica,  
stylata, subcutanea, tussilaginis,  
vecors, vinulus, viridescens, vittata.

- Tephritis: arnicae, cinerea, heraclei,  
hyoscyami, leontodontis.
- Terellia: cornuta, lappae, ruficauda,  
serratula, serratulae, tussilaginis.
- terraenovae: groenlandica, orior.
- terraereginae: hilli.
- tessariae: fucata, furcata.
- tessella: tessellata.
- testaceus: elongata.
- Tetanocera: alata, elata, elater.
- Thelaira: lateralis, leucozona, lucozona,  
nigrina, nigripes, solivagus.
- Themira: consencis, cossenico, fimetis,  
putris.
- Theresia: valida.
- Thereva: fasciata, hirta, lugubris, nobilis,  
nobilitata, plebeja.
- Therevidae: anilis, ardea, bilineata,  
fasciata, flavipes, hirta, lugubris,  
nobilis, nobilitata, ochropus, plebeja.
- Thoracites: abdominalis.
- Thricops: atra, diaphana, hyalinata,  
variabilis.
- Thyreophora: cynophila.
- Thyridanthrax: nigrita.
- Thysocnema: striata.
- tigrina: notata, rapax.
- timorensis: benedicta, intrahens, obtrusa.
- totliana: fabricii, fasciata.
- Toxomerus: dispar.
- Trachydromia: cimecoides, cimicooides.
- transfugus: transfuga.
- Tricharaea: occidua.
- Trichodura: anceps.
- Trichopoda: pennipes.
- trilineata: graeca, hypoleon.
- tringaria: vanella.
- tringarius: tringaria.
- Trixia: conspersus.
- Tromodesiopsis: atrifrons.
- Tropidia: scitus.
- Trupanea: radiata, stellata.
- Trypetia: artemisiae, florescentiae,  
perelegand, perelegans.
- Trypetoptera: hieracii, punctulata.
- tuguriorum: extranea.
- uliginosus: uliginosa.
- unguicornis: flavifrons, varicus, varieus.
- ungulata: squamata, squamifera.
- ungulatus: unguulata.
- UNKNOWN: abrupta, adiaphora,  
adspersus, aestiuia, aestivationis,  
aestuans, agilis, albomicans, allii,  
alterabilis, ambulans, annulata,  
aphidioides, argentata, argentea,  
aselliformis, atra, aurata, bibosa,  
bicolor, bilineata, bimaculata,  
brunnea, buccata, buphthalmi, capito,  
caracteristica, carinata, castanea,  
caudata, cerasi, chrysocephala,  
chrysostoma, cineraria, cinerascens,  
cinerea, cingulata, clandestina,  
clauata, clavicornis, claviformis,  
claviventris, concinna, conferta,  
conica, contigua, costata, crassipes,  
cruciata, cuprea, cyanophthalma,  
cylindrica, depressa, despecta,  
divaricata, doclea, doliata, duplicata,  
elongata, erratica, erythrocephala,  
fallax, familiaris, fervida, fimetorum,  
flauicans, flavescentis, flavicans,  
flavipes, fulvipes, fungorum, fusca,  
fuscipes, galeata, geographica, gibba,  
gibbosa, gigantea, glaucoptera,  
grisea, guttata, helenii, helvola,  
hemipterus, hirticollis, hortensis,  
hydrophila, hydropica,  
ichneumonoides, larvata, lateritia,  
latipes, latro, leprae, leuconyx,  
leucophaeus, leucopis, leucoptera,  
leucosticta, libatrix, liturata, longipes,  
luctuosa, lutea, macrocephala,  
mactans, maculata, maculatus,  
marginalis, marginella, marmorata,  
melanochrysa, melanura, mesoleuca,  
minimus, minutissima, muraria,  
nigra, nigripilis, nivalis, nobilis,  
notata, novemmaculata,  
novempunctata, ochroptera,  
orbiculata, pallida, pennipes,  
personata, pertinax, picata, picea,  
piceae, plombea, plumbea,  
potentillae, praedo, praeusta,  
prodromaria, psyllia, pulicaria,  
pullata, punctata, punctifera, putrida,  
putris, quadrata, quadricolor,

- quadrifasciata, quadrilineata,  
 quadrimaculata, quinquefasciata,  
 quinquepunctata, resinosa, reticulata,  
 rubripes, rufa, ruficapilla, rufipes,  
 rufiventris, sacalis, scolopacina,  
 secalis, segmentaria, semiflava,  
 sericea, setaria, seticornis, setosa,  
 smaragdina, solstitialis, sordida,  
 spinifera, squalida, stellata, strigosa,  
 stuans, subcutanea, subulata, suecica,  
 syrphus, teres, terrestris, tessellata,  
 tibialis, tipulaea, tipuliformis,  
 topiaria, tricolor, tridens,  
 trigrammica, trimaculata, trinervis,  
 tripunctata, tristis, truncata,  
 tuberculata, turcica, varia, velox,  
 venosa, versicolor, versipellis,  
 vinculata, violacea, vires, viridana,  
 vitrata, vivipara.  
 urbana: rustica.  
**Urophora:** arctii, cardui, dauci, iacobae,  
 jacobaeae, placida, solstitialis, stylata.  
 ursina?: argyrostoma.  
 vagabunda: varians.  
 vagans: strigosa, volito.  
 valgus: valga.  
 valida: validus.  
 ventrosa: kasauliensis, nigrithorax,  
     pungoana, xanthomela.  
**Vermileo:** vermileo.  
**Vermileonidae:** vermileo.  
 versicolor: hartigii, versicolorata.  
 vertiginosa: marmorata.  
 vesiculosum: vesiculosa.  
 vespertina: nigrita.  
 vespiformis: vespaesimilis.  
 vespillo: ruficornis, virescens.  
 vetustissima: angustifrons, fieldi,  
     latifrons, nigrirurus, prisca, pumila.  
 viarum: pusilla.  
 viatica: fulvicornis.  
 vibrans: urticae.  
 vicana: decolor, mantes, mantos, vicanus.
- vicina: aucta, carnivora, erythrocephala,  
     scutellata, spitzbergensis, thuscia.  
**Vidalia:** cornuta.  
**Villa:** troglodyta.  
**vinetorum:** surinamensis.  
**violacea:** infixa.  
**viredaurea:** munda, reflectens, viridaurea.  
**viridicyanea:** anthracina.  
**viridula:** jejuna.  
**Volucella:** annulatus, apivora,  
     bombylans, hirsutissima, inanis,  
     inflata, melanopyrrha, micans,  
     pellucens, plumata, plumosa,  
     trifasciata, valentina, zonaria.  
**volvulus:** cylindrica.  
**vomitoria:** affinis, carnaria, coerulea,  
     minimus, obscoena.  
**vulgaris:** agilis.  
**vulpina?:** investigator.  
**Winthemia:** pinguis, quadripustulata.  
**Xanthandrus:** comtus.  
**Xanthogramma:** anteaimbulo,  
     anteambulo, citreofasciata,  
     citrofasciata, festiva, laeta,  
     pedissequus, vespiformis.  
**xanthomelaena:** albomaculata,  
     dorsomaculata, irwinrossi, lothari,  
     rufiventris, setigera, xanthomelas.  
**Xenocalliphora:** hortona, icela.  
**Xylophagidae:** australis, cincta, crucis,  
     ferruginea, grandis, olens, testacea.  
**Xylophagus:** cincta.  
**Xylota:** contracta, fucatus, longisco,  
     maritima, melanochrysa, pigra,  
     segnis, semulater, simulator,  
     simulator, sylvarum.  
**Xyphosia:** arcuata, miliaria.  
**Zaphne:** ambigua.  
**Zenillia:** libatrix, ochracea.  
**zoe:** perelegand, perelegans.  
**zonaria:** valentina.  
**zonata:** rufipes, sinensis, zonalis,  
     zonatus.

## Summary of nomenclatural changes

Names are arranged by family and alphabetically by genus and species within family. Lectotype entries and junior synonyms are listed under the original combination of the species group name, others are under the present combination.

### New names

**Calliphoridae:** *Protocalliphora rognesi* Thompson & Pont = *chrysorrhoea* Meigen 1826, not Scopoli 1763; **Muscidae:** *Musca mallochi* Thompson & Pont = *squamata* Malloch 1934, not Müller 1764; **Psilidae:** *Chamaepsila hennigi* Thompson & Pont = *rosae* Fabricius 1794, not DeGeer 1776; **Sarcophagidae:** *Anapunophyto lopesi* Thompson & Pont = *pusilla* Macquart 1848, not Gmelin 1790.

### New lectotype designations

**Calliphoridae:** *Musca laemica* Walker; **Syrphidae:** *Musca pinastri* De Geer, *Musca pyrorum* Schrank.

### Changes in taxonomic status (New synonymy)

**Anisopodidae:** *Musca nigricans* de Villers = *Sylvicola fuscatus* (Fabricius); **Bombyliidae:** *Musca hirsuta* de Villers = *Hemipenthes maurus* (Linnaeus); **Calliphoridae:** *Musca laemica* Walker = *Calliphora stygia* (Fabricius), *Musca melanoptera* Fallén = *Morinia doronici* (Scopoli), *Musca bicolor* Walker = *Phormia regina* Meigen, *Musca sordida* Zetterstedt = *Protocoliphora rognesi* Thompson & Pont, *Musca felix* Walker = *Chrysomya albiceps* (Wiedemann), *Musca virens* Wiedemann = *Cosmina limbipennis* (Macquart), *Musca quadrilineata* Fabricius = *Eumesembrinella flavipennis* (Macquart), *Musca segmentaria* Fabricius = *Hemilucilia nubipennis* (Rondani), *Musca gentilis* Meigen = *Melinda gentilis* (Robineau-Desvoidy); **Chloropidae:** *Musca nigra* Linnaeus = *Oscinella frit* (Linnaeus), *Musca pigra* Geoffroy = *Meromyza saltatrix* Linnaeus, *Musca pumilionis* Banks = *Chlorops pumilionis* (Bjerkander), *Musca saltatrix* Geoffroy = *Chlorops pumilionis* (Bjerkander); **Curtonotidae:** *Musca gibba* Fabricius = *Curtonotum taeniatum* Hendel; **Dolichopodidae:** *Musca squamata* Müller = *Dolichopus unguarius* (Linnaeus), *Musca ciliata* Müller = *Poecilobothrus nobilitatus* (Linnaeus), *Musca nigripes* Fabricius = *Chrysotus laesus* (Wiedemann); **Keroplatidae:** *Tipula lineata* Fabricius = *Cerotelion striatum* (Gmelin); **Micropezidae:** *Hoplocheiloma fabricii* Steyskal = *Hoplocheiloma totiana* (Gmelin); **Otitidae:** *Musca hortulana* Rossi = *Ceroxys hyalinata* (Panzer), *Musca graminum* Fabricius 1794 = *Dorycera aquatica* (Geoffroy), *Musca aenea* Fabricius = *Physiphora clausa* (Macquart); **Phoridae:** *Bibio*

*florea* Fabricius = *Diplonevra florescens* (Turton); **Sarcophagidae:** *Musca cinerea* Fabricius = *Leucomyia alba* (Schiner); **Sciomyzidae:** *Musca cribrata* de Villers = *Coremacera marginata* (Fabricius); **Sepsidae:** *Musca spondylii* Schrank 1837 = *Saltella sphondyliae* (Schrank 1803); **Stratiomyidae:** *Musca sellata* Fuesslin = *Psecticus posticus* (Wiedemann); **Syrphidae:** *Musca apiformis* Geoffroy = *Eristalis tenax* (Linnaeus), *Musca balteata* De Geer = *Episyphus cannabina* (Scopoli), *Musca bipunctata* Müller = *Chrysotoxum arcuatum* (Linnaeus), *Musca bombyliformis* Geoffroy = *Merodon equestris* (Fabricius), *Musca chrysozonias* Gmelin = *Chrysotoxum fasciatum* (Müller), *Musca consisto* Harris = *Dasysyrphus pinastri* (De Geer), *Musca contracta* Geoffroy = *Xylota segnis* (Linnaeus), *Musca depressa* Swederus = *Leucozona glaucia* (Linnaeus), *Musca elegans* de Villers = *Episyphus cannabina* (Scopoli), *Musca festiva* Geoffroy = *Chrysotoxum arcuatum* (Linnaeus), *Musca fulva* Gmelin = *Cheilosia illustrata* (Harris), *Musca hirsutissima* de Razoumowsky = *Volucella bombylans* (Linnaeus), *Musca imbellis* Harris = *Chrysotoxum arcuatum* (Linnaeus), *Musca inaequalis* Geoffroy = *Episyphus cannabina* (Scopoli), *Musca leucocephala* Gmelin = *Eristalinus aeneus* (Scopoli), *Musca lineata* Geoffroy = *Eumerus strigatus* (Fallén), *Musca lineolae* Harris = *Eristalis interrupta* (Poda), *Musca lineolata* Gmelin = *Chrysogaster cemiteriorum* (Linnaeus), *Musca longipes* Thunberg = *Chalcosyrphus femoratus* (Linnaeus), *Musca lunula* de Villers = *Eristalis pertinax* (Scopoli), *Musca luteola* Gmelin = *Myolepta dubia* (Fabricius), *Musca maritima* Scopoli = *Xylota segnis* (Linnaeus), *Musca melanochrysa* Gmelin = *Xylota segnis* (Linnaeus), *Musca naevia* Gmelin = *Milesia semiluctifera* (de Villers), *Musca nectarina* Gmelin = *Episyphus cannabina* (Scopoli), *Musca pyrorum* Schrank = *Dasysyrphus pinastri* (De Geer), *Musca vespiformis* Geoffroy = *Xanthogramma festiva* (Linnaeus).

#### New and revised combinations

**Bibionidae:** *Bibio bidentatus* (de Villers), *Dilophus novemmaculatus* (de Villers); **Calliphoridae:** *Isomyia selecta* (Walker), *Onesia hederacea* (Geoffroy), *Phumosia dioclea* (Walker), *Pollenia umbrifera* (Walker); **Chloropidae:** *Chlorops pulchella* (Müller), *C. quadristrigata* (Donovan), *C. lepidus* (Harris); **Empididae:** *Hilara globulipes* (Geoffroy); **Heleomyzidae:** *Suillia tuberis* (Vallot); **Keroplatidae:** *Macroceras longicornis* (Fabricius), *M. macroceras* (Gmelin); **Muscidae:** *Morellia nigricosta* Hough; **Otitidae:** *Doryceria aquatica* (Geoffroy); **Stratiomyidae:** *Psecticus japonica* (Thunberg); **Sarcophagidae:** *Agria affinis* (Fallén); **Syrphidae:** *Cheilosia aenea* Scopoli.

## New status

**Calliphoridae:** *Morinia doronici* (Scopoli); **Curtonotidae:** *Curtonotum taeniatum* Hendel; **Dolichopodidae:** *Chrysotus laesus* (Wiedemann); **Keroplatidae:** *Cerotelion striatus* (Gmelin); **Micropezidae:** *Hoplocheiloma totiana* (Gmelin); **Otitidae:** *Ceroxys hyalinata* (Panzer), *Melieria gangraenosa* (Panzer), *Otites elegans* (Latreille), *Physiphora clausa* (Fabricius); **Phoridae:** *Diplonevra florescens* (Turton); **Sarcophagidae:** *Leucomyia alba* (Schiner); **Stratiomyidae:** *Psecticus posticus* Wiedemann; **Syrphidae:** *Baccha pereelixis* Harris, *Dasyzyrus pinastri* (De Geer), *Epi-syrphus cannabina* (Scopoli), *Eristalis lineatus* (Harris), *Eristalis lyra* (Harris), *Ischiodon nigra* (Forskål), *Psarus coalescens* (Geoffroy); **Tachinidae:** *Tachina tessella* (Turton), *Thelaira nigrina* (Fallén), *Zenillia fulva* (Fallén).

## Overlooked names (names not in recent Diptera catalogs)

**Anisopodidae:** *nigricans* de Villers; **Athericidae:** *atherix* Donovan; **Bibionidae:** *bidentata* de Villers, *ciliata* de Villers, *novemmaculata* de Villers, *tuberculata* de Villers; **Calliphoridae:** *bicolor* Walker, *caerulea* Brullé, *caesarina* Scopoli, *caesarion* Scharfenberg, *chalybaea* Gravenhorst, *dioclea* Walker, *dolens* Zetterstedt, *felix* Walker, *flavicans* Doleschall, *fulvimaculata* Manuel, *ionia* Meigen, *micans* Fabricius, *minimus* Harris, *polisma* Walker, *remigro* Harris, *reviso* Harris, *selecta* Walker, *suffusa* Walker, *umbrifera* Walker, *violacea* Fabricius, *virens* Wiedemann, *viridicaerulea* De Geer; **Chamaemyiidae:** *gibbosa* De Geer; **Chloropidae:** *erro* Harris, *nigra* Linnaeus, *pigra* Geoffroy, *pulchella* Müller, *pumilionis* Banks, *saltatrix* Geoffroy; **Coelopidae:** *relictus* Harris, *simulator* Harris; **Dolichopodidae:** *amicula* Harris, *caudex* Harris, *ciliata* Müller, *festiva* Donovan, *gibba* Müller, *hebes* Harris, *ingruo* Harris, *sannio* Harris, *semiargenteata* Donovan, *semiargentea* Turton, *squamata* Müller, *squamifera* Müller; **Dryomyzidae:** *mellinus* Harris; **Empididae:** *globulipes* Geoffroy; **Fanniidae:** *minor* Harris; **Heleomyzidae:** *avarus* Harris, *dejectus* Harris, *sandaracha* Harris; **Keroplatidae:** *longicornis* Fabricius, *macroceras* Gmelin; **Lauxaniidae:** *caeruleo-phthalmica* Scopoli, *mulsus* Harris; **Lonchaeidae:** *vernicina* Scopoli; **Micropezidae:** *ephippium* Walckenaer, *pedo* Harris, *peragro* Harris, *totliana* Gmelin; **Opomyzidae:** *galpinus* Harris; **Otitidae:** *alceae* Preyssler, *aquatica* Geoffroy, *myennis* Coquebert, *porca* Latreille; **Phaeomyiidae:** *nigripennis* Fabricius; **Pallopteridae:** *minutus* Harris; **Phoridae:** *florescens* Turton; **Piophilidae:** *ornithovora* Scopoli; **Pipunculidae:** *cephalotes* Bosc; **Platystomatidae:** *naevosus* Harris, *singenesiae* Manuel, *variolosa* Schrank; **Psilidae:** *histrionica* Fabricius, *longerro* Harris, *luteus* Harris, *percussus* Harris, *rufa* De Geer, *vagus* Harris; **Rhagionidae:** *cingulatus* Donovan, *norwegica* Gmelin; **Sarcophagidae:** *frontalis* Latreille, *haemorrhoa* Gravenhorst, *labiata* Fabricius, *leucocephala* de Villers, *martia* Scopoli, *moneta* Walker, *nigrita* Scopoli, *prolabor* Harris, *pusilla* Macquart, *reverta* Harris,

*sexpunctata* Fabricius, *taenionota* Wiedemann, *valens* Harris; **Scathophagidae**: *abbreviata* Gravenhorst, *exilis* Harris, *lucophaeus* Harris, *paganus* Harris; **Scatopsidae**: *nectarivora* Turton; **Scenopinidae**: *tarda* Linnaeus; **Sciaridae**: *maura* Turton; **Sciomyzidae**: *cibrata* de Villers, *delicate* Harris, *oratoria* Turton, *tener* Harris, *varicus* Harris; **Sepsidae**: *elongata* Müller, *palpator* Harris, *spondylii* Schrank; **Stratiomyidae**: *aequinoctialis* Linnaeus, *japonica* Thunberg, *melanopis* Gmelin, *minor* Turton, *minutior* Turton, *nigricans* Gmelin, *sellata* Fuesslin; **Syrphidae**: *apiformis* Geoffroy, *bombyliformis* Geoffroy, *chrysorrhaea* Scopoli, *contracta* Geoffroy, *depressa* Swederus, *elegans* de Villers, *fusiformis* Turton, *hirsutissima* de Razumowsky, *hortulana* Scopoli, *inaequalis* Geoffroy, *interrupta* Geoffroy, *intricata* Turton, *lineata* Geoffroy, *longipes* Thunberg, *merulina* Scopoli, *murina* Schrank, *nectarina* Gmelin, *nigra* Forskål, *pulla* Gmelin, *univalvis* Turton, *versicolorata* Turton, *vespaesimilis* Barbut; **Therevidae**: *ochropus* Turton; **Xylophagidae**: *olens* Fuesslin.

## Acknowledgements

We have been grateful to receive the help of the following, who have reviewed the manuscript, assisted in its preparation or provided invaluable additional advice and information: Drs. T. J. Henry, D. R. Miller, A. L. Norrbom, Brian Brown, and M. E. Schauff, Systematic Entomology Laboratory, USDA, Washington; J. W. Ismay, Hope Entomological Collections, University Museum, Oxford; L. Matile, Entomology, Muséum National d'Histoire Naturelle, Paris; W. N. Mathis, Department of Entomology, Smithsonian Institution, Washington; Amnon Freidberg, Department of Zoology, Tel-Aviv University, Tel-Aviv; J. P. O'Connor, National Museum of Ireland, Dublin; J. Clary, Musée Guimet d'Histoire Naturelle, Lyon; I. C. Beavis, Tunbridge Wells Museum and Art Gallery, Royal Tunbridge Wells, Kent; T. Pape, V. Michelsen, Entomology, Universitets Zoologiske Museum, Copenhagen; Sarah Mascherini, Museo Zoológico de "La Specola," Florence; Knut Rognes, Stavanger Laererhøgskole, Stavanger; G. Bächli, Zoologisches Museum, Zürich; D. J. Bickel, Entomology, Australian Museum, Sydney; and Rudolf Rozkosny, Department of Environmental Studies, Masaryk University, Brno. We are grateful to the Keeper of Entomology at the Natural History Museum, London, for permission to quote from an unpublished list of Diptera in the Linnaean Collection, compiled by J. P. Dear. Special thanks are due to two of our colleagues who have continually shared data, "tested" our database, and proofed numerous drafts of the final manuscript: Drs. Curtis W. Sabrosky, Systematic Entomology Laboratory,

USDA, Washington and Neal L. Evenhuis, Department of Entomology,  
Bishop Museum, Honolulu.

Building a database of scientific names requires a tremendous amount of bibliographic work. Our database could not have been built without the assistance of a number of skilled and dedicated librarians, without whose help we would not have been able to see all the reference herein cited. So, we are especially grateful to the following: Ms Pamela Gilbert, General Library, The Natural History Museum (formerly BMNH), London; Julie Harvey, Entomology Library, The Natural History Museum, London; Stella Newton, Hope Library, University Museum, Oxford; Jack Macquardt (now retired) and Carol Hahn, Natural History Library, Smithsonian Institution, Washington; Leslie Overstreet and Ellen Wells, Special Collections, Smithsonian Institution, Washington and Ruth Schallert, Botany Library, Smithsonian Institution, Washington. Again, especial thanks are due to Dr. Neal Evenhuis, who provided us with most of the precise publication dates here listed for the first time.

## References Cited.

**Ackland, D. M. & E. Taylor**

- 1972 Notes on the Verrall-Collin collection of Diptera in the Hope Department of Entomology, University Museum, Oxford. *Ent. mon. Mag.* 108: 12-15. (27 September)

**Aldrich, J. M.**

- 1929 Further studies of types of American muscoid flies in the collection of the Vienna Natural History Museum. *Proc. U. S. natl Mus.* 74(19), 34 pp.

**Aldrovandus, U.**

- 1602 *De animalibus insectis libri septem cum singylorum iconibus ad viuum expressis.* [10+] 767 [+ 43] pp. Bellagamba, Bonona. [=Bologna]

**Allioni, C.**

- 1766 Manipulus Insectorum Taurinensium. *Mélanges Soc. R. Turin* 3: 185-198.

Aim Stein — see Stein, J. R. Am

**Ander, K.**

- 1952 Mitteilung über die im Entomologischen Museum zu Lund befindlichen Typen. *Opusc. ent.* 17: 77-79. (12 March)

**Andersen, S.**

- 1982 Revision of European species of *Siphona* Meigen (Diptera: Tachinidae). *Ent. Scand.* 13: 149-172. (5 July)

**Anker, J.**

- 1950 Otto Friderich Müller's *Zoologia Danica. Libr. Res. Monogr., Copenhagen* 1, xiv + 108 pp. (21 June)

**Arntfield, P. W.**

- 1975 A revision of *Graphomya* Robineau-Desvoidy (Diptera: Muscidae) from North America. *Canad. Ent.* 107: 257-302. (27 February)

**Austen, E. E.**

- 1909 New genera and species of blood-sucking Muscidae from the Ethiopian and Oriental Regions, in the British Museum (Natural History). *Ann. Mag. nat. Hist.* (8) 3: 285-299. (March)

- 1910 A new Indian species of *Musca*. *Ann. Mag. nat. Hist.* (8) 5: 114-117. (January)

Avelin, G. E. — see Linnaeus 1764b

**Awati, P. R.**

- 1915 Studies in flies. I. Chaetotaxy and pilotaxy of Muscidae and range of their variability in the same species. *Indian J. med. Res.* 3: 135-148, pls 12-14. (July)

- 1916a Studies in flies. - II. Contributions to the study of specific differences in the genus *Musca*. 1. - Genitalia. *Indian J. med. Res.* 3: 510-529, 2 tables, pls 32-50. (January)

- 1916b Studies in flies. Contributions to the study of specific differences in the genus *Musca*. 2. - Structures other than genitalia. *Indian J. Med. Res.* 4: 123-139. (July)
- 1917 Studies in flies. III. Classification of the genus *Musca* and descriptions of the Indian species. *Indian J. med. Res.* 5: 160-191, pls 20-29. (July)
- 1918 A new larviparous *Philaematomyia* (*Philaematomyia indica*, N. Sp.). *Indian J. med. Res.* 5: 529-539, pls 50-55. (January)
- 1920 A note on the genitalia of Portschinsky's species *M. corvina* (*vivipara*) and *M. corvina* (*ovipara*). *Indian J. med. Res.* 8: 89-92, 2 tables, pl. 13. (July)
- Baccetti, B.**
- 1962 Pietro Rossi, Naturalista Toscano del '700. *Frustula ent.* 5(3), 30 pp.
- Bächli, G.**
- 1982 On the type material of Palearctic species of Drosophilidae (Diptera). *Beitr. Ent., Berlin* 32: 289-301.
- 1984 Die Drosophiliden-Typen der Dipterensammlung des Zoologischen Museums in Berlin. *Mitt. zool. Mus. Berlin* 60: 229-261.
- 1990 Type specimens of Drosophilidae (Diptera) described by Linnaeus, Fallén, Wahlberg and Zetterstedt. *Rev. suisse Zool.* 97: 843-849.
- Baeckner, M. A. — see Linnaeus 1764c
- Banks, J.**
- 1791 On the *Musca pumilionis*. *Ann. Agric.* 16(2): 176-177, 1 pl.
- Barbut, A.**
- 1781 *Les Genres des Insectes de Linné; constatés par divers échantillons d'Insectes d'Angleterre, copiés d'Après Nature.* xx + 371 [+ 20] pp., 22 pls. J. Sewell, London. (after 9 February)  
[With a second engraved English pictorial title page: *The genera insectorum of Linnaeus, exemplified by various specimens of English insects drawn from nature.* Both title pages state that the work was published by the author. The french title page gives Dixwell as the printer and J. Sewell as publisher.]
- Baumgartner, D. L. & B. Greenberg**
- 1984 The genus *Chrysomya* (Diptera: Calliphoridae) in the New World. *J. med. Ent., Honolulu* 21: 105-113.
- Beavis, I. C.**
- 1988 *Insects and other invertebrates in classical antiquity.* xv + 269 pp. University of Exeter, Exeter.
- Bechstein, J. M. & G. L. Scharfenberg**
- 1805 *Vollständige Naturgeschichte der schädlichen Forstinsekten. Ein Handbuch für Forstmänner, Cameralisten und Oekonomen. Dritte und letzte Abtheilung. Nebst dem Anhang von den nützlichen Forstinsekten.* [Vol. 3], pp. 605-1042, pls. 10-13. C. F. E. Richter, Leipzig.
- Becker, T.**
- 1902- Aegyptischen Dipteren. *Mitt. zool. Mus. Berlin* 2: 1-66 (1902, 12 August); 67-195 (1903, 23 September)

**Beier, M.**

- 1973 The early naturalists and anatomists during the Renaissance and seventeenth century. Pp. 81-94. In Smith, R. F., T. E. Mittler & C. N. Smith [eds], *History of Entomology*, vii + 517 pp. Annual Reviews Inc., Palo Alto, California.

**Bezzi, M.**

- 1892 Di alcuni ditteri raccolti nel paese dei Somali dall'ingegnere L. Bricchetti-Robecchi. *Ann. Mus. Civ. Stor. Nat. Genova* 32: 181-196. (March)
- 1906 Noch einige neue Namen für Dipterengattungen. *Ztschr. System. Hymenopt. Dipt.* 6: 49-55. (1 January)
- 1907 Nomenklatorisches über Dipteren. II. *Wien. ent. Ztg* 26: 292-296. (20 August)
- 1908a Nomenklatorisches über Dipteren. III. *Wien. ent. Ztg* 27: 74-84. (20 February)
- 1908b Nomenklatorisches über Dipteren. IV. *Wien. ent. Ztg* 27: 291-295. (1 December)
- 1908c Ditteri eritrei raccolti dal Dott. Andreini e dal Prof. Tellini. Parte seconda. Diptera cyclorrhapha. *Bull. Soc. Ent. Ital.* 39: 3-199 (1907). (9 November)
- 1911 Miodarii superiori raccolti dal signor C. W. Howard nell'Africa australe orientale. *Boll. Lab. Zool. R. Scuola Agric. Portici* 6: 45-104 (1912)  
[The separate is dated as published on 14 November 1911 (see Smith et alia 1980: 917)]
- 1921 *Musca inferior* (Stein), type of a new genus of philaematomyine flies (Diptera). *Ann. Trop. Med. Parasit.* 14: 333-340. (8 February)
- 1923 Les mâles de *Musca albina* Wied. et de *Musca lucidula* Loew (Dipt.). *Bull. Soc. r. ent. Égypte* 7 [1922-1923]: 108-118.

**Bezzi, M. & P. Stein**

- 1907 Cyclorrhapha Aschiza. Cyclorrhapha Schizophora: Schizometopa. Pp. 1-189, 190-749. In Becker, T., M. Bezzi, K. Kertész & P. Stein (eds.), *Katalog der paläarktischen Dipteren*. Vol. 3, 828 pp. Budapest. (1 December)

**Bickel, D. J.**

- In The Australian Sciapodinae (Diptera: Dolichopodidae), with a review  
 Press of the Oriental and Australasian faunas, and a world conspectus of the  
 subfamily. *Rec. Australian Mus. Suppl.*

**Bigot, J. M. F.**

- 1887 [Diagnoses abrégées de quelques Diptères nouveaux, provenant de l'Amérique du nord, dont les descriptions détaillées seront publiées ultérieurement, et qui tous appartiennent à sa collection]. *Bull. Bimens. Soc. ent. Fr.* 1887 (21): clxxii-clxxxiv. (19 November)

[Due to an error in the printing of Bulletin No. 21, the pagination is different from the reprinted version in 1888b.]

1888a Diptères nouveaux ou peu connus. *Muscidi* (J.B.) *Bull. Soc. Zool. France* 12: 581-617 (1887). (15 January)

1888b [Diagnoses abrégées de quelques Diptères nouveaux, provenant de l'Amérique du nord, dont les descriptions détaillées seront publiées ultérieurement, et qui tous appartiennent sa collection]. *Bull. Soc. ent. Fr.* (6) 7: clxxx-clxxxii. (11 April)

**Bjerkander, C.**

1777 Rön Om Rot-masken. *K. Vetenskapsakad. Handl.* 38: 29-43.  
[A German synopsis of this article is in Bjerkander 1783.]

1778 Rag-Dvergs Masken. *K. Vetenskapsakad. Handl.* 39: 240-241.

1780 Beskrifning pa tvänne slags Maskar, som äro Larver af härtils okände Insector och funnits pa Ror-Kal *Napo-Brassica*. *K. Vetenskapsakad. Nya Handl.* 1: 194-196.

1781 Hafre Masken. *K. Vetenskapsakad. Nya Handl.* 2: 171-172.

1783 Über das Wurzelinsekt. *Lichtenberg, Mag. neueste Phys. Naturgesch.*, Gotha 2: 101-104.

1793 *Musca subcutanea*, eller en ny och obeskrefven fluga uti korn-bladen. *K. Vetenskapsakad. Nya Handl.* 14: 57-60, pl. 2.

**Blackwelder, R. E.**

1947 The dates and editions of Curtis' British Entomology. *Smithson. Misc. Coll.* 107 (5), 27 pp., 4 pl. (12 June)

**Blanchard, E.**

1840 *Histoire Naturelle des Insectes. Orthoptères, Nevropières, Hémiptères, Lépidoptères et Diptères*. 672 pp., 71 pls, Duméril, Paris. (December)

**Blankaart, S.**

1688 *Schou-burg der rupsen, wormen, ma'den, en vliegende dierkens daar uit voortkomende*. 4 + 232 [+ 2] pp., 16 coloured pls. J. ten Hoorn, Amsterdam.

**Bloch, M. E.**

1776 Beitrag zur Naturgeschichte des Kopals. *Beschäft. Berlin. Gesellsch. naturf. Fr.* 2: 91-224, pls. III-V.

**Bodenheimer, F. S.**

1928 *Materialien zur Geschichte der Entomologie bis Linné*. Band I. x + 498 pp., 24 pls. Junk, Berlin.

1929 *Materialien zur Geschichte der Entomologie bis Linné*. Band II. vi + 486 pp., 4 pls. Junk, Berlin.

**Boisduval, J. B. A. D. de**

1835 Faune entomologique de l'Océan Pacifique, avec l'illustration des Insectes nouveaux recueillis pendant le voyage . Deuxième Partie. Coléoptères et autres ordres. vii + 716 pp., plates 6-12. *In his Voyage de découvertes de l'Astrolabe, exécuté par ordre du Roi, pendant les années 1826-1827-1828-1829, sous le commandement de M. J. Dumont d'Urville*. J. Tastu, Paris. (9 May)

**Bosc d'Antic, L. A. G.**

1792 Description de deux Mouches. *Bull. Sci. Soc. Philom.*, Paris 1: 18.

**Bouché, P. F.**

1834 *Naturgeschichte der Insekten, besonders in Hinsicht ihrer ersten Zustände als Larven und Puppen. Erste Lieferung* vi + 216 pp., 10 pls. Nicolai, Berlin.

**Brauer, F. M.**

1899 Beiträge zur Kenntnis der *Muscaria schizometopa*. Bemerkungen zu den Originalexemplaren der von Bigot, Macquart und Robineau-Desvoidy beschriebenen *Muscaria schizometopa* aus der Sammlung des Herrn G. H. Verrall. Dritte Folge. *Sber. Akad. Wiss. Wien* 108, Abt. 1: 495-529.

**Brauer, F. M. & J. von Bergenstamm**

1889 Die Zweiflügler des Kaiserlichen Museums zu Wien. IV. Vorarbeiten zu einer Monographie der *Muscaria Schizometopa* (exclusive Anthomyidae). Pars I. *Denkschr. Akad. Wiss., Wien*, 56: 69-180, 11 pls. (March)

1891 Die Zweiflügler des Kaiserlichen Museums zu Wien. V. Vorarbeiten zu einer Monographie der *Muscaria Schizometopa* (exclusive Anthomyidae). Pars II. *Denkschr. Akad. Wiss., Wien*, 58: 305-446.

1893 Die Zweiflügler des Kaiserlichen Museums zu Wien. VI. Vorarbeiten zu einer Monographie der *Muscaria Schizometopa* (exclusive Anthomyidae). Pars III. *Denkschr. Akad. Wiss., Wien*, 60: 89-240.

**Brullé, G. A.**

1832 IV<sup>e</sup> Classe. Insectes. Pp. 64-395. In Bory de Saint-Vincent (ed.), *Expédition scientifique de Morée. Section des sciences physiques* 3(1) (Zool. 2), 400 pp. Levrault, Paris & Strasbourg.

**Brunetti, E.**

1910 Revision of the Oriental bloodsucking Muscidae (Stomoxinae, *Philaematomyia* Aust., and *Pristirhynchomyia*, gen. nov.). *Rec. Indian Mus.* 4(4): 59-93, pls. 7-8. (5 July)

**Brünnich, M. T.**

1764 *Entomologia, sistens insectorum tabulas systematicas, cum introduktione et iconibus. Insektaere, indeholdende Insekternes systematiske tavler, samt indledning og figurer.* [7+] 88 pp., 1 pl. Godiche, Hafniae, Kiøbenhavn [= Copenhagen].

**Bryk, F.**

1949 De Geers Insekten Sammlung eine Fundgrube von unbekannten Linnéischen Typen! *Opusc. Ent.* 14: 77-80.

**Burger, J. F. & F. C. Thompson**

1981 The *Tabanus striatus* complex (Diptera; Tabanidae): A revision of some Oriental horse fly vectors of surra. *Proc. ent. Soc. Washington* 83: 339-358. (30 August)

**Cederhielm, I.**

- 1798 *Favnae Ingricae Prodromus exhibens methodicam descriptionem Insectorum agri Petropolensis praemissa Mammalium, Avium, Amphibiorum et Piscium envmeratione.* xviii + 348 pp., 3 pls. Iohann. Freid. Hartknoch, Lipsiae [=Leipzig].

[Name given by Hagen as Cederhielm, J.]

**Chalmers-Hunt, J. M.**

- 1976a Entomological sales. Pp. 3-14. In his *Natural History auctions. 1700-1972. A register of sales in the British Isles.* xii + 189 pp. Sotheby Parke Bernet, London.

**Chalmers-Hunt, J. M.**

- 1976b The register of natural history sales. Pp. 53-189. In his *Natural History auctions. 1700-1972. A register of sales in the British Isles.* xii + 189 pp. Sotheby Parke Bernet, London.

**Clausen, P. J.**

- 1977 A revision of the Nearctic, Neotropical, and Palearctic species of the genus *Ochthera*, including one Ethiopian species, and one new species from India. *Trans. Amer. ent. Soc.* 103: 451-530. (1 November)

**Cobres, J. P.**

- 1781- *Deliciae Cobresianae. Büchersammlung zur Naturgeschichte.* [I 1782 Theil] [iii +] xxviii + 1-470 pp. (1781); II Theil, 471-956 [+ 1 errata] pp.(1782) J. F. de Göz, [Augsburg].

[A facsimile reprint was issued by A. Asber & Co., Amsterdam, 1966.]

**Collin, J. E.**

- 1926 Supplementary notes on the genus *Lucilia*, Desv. (Diptera). *Trans. ent. Soc. London* 1926: 258-260. (21 December)
- 1948 On the classification of the genera allied to *Musca* L. (Diptera). *Proc. R. ent. Soc. London (B)* 17: 125-127. (15 October)

**Coquebert de Montbret, A. J.**

- 1798 Sur une nouvelle espèce de Mouche. *Bull. Sci. Soc. Philom.*, Paris 2: 145 ("an VII")
- 1804 *Illustratio Iconographica Insectorum quae in Musaeis parisiniis observavit et in lucem edidit Joh. Christ. Fabricius, praemissa ejusdem descriptionibus; accedunt species plurimae, vel minus aut nondum cognitae.* Vol. 3, [4] + pp. 91-142. Petri Didot Natu Majoris, Parisiis [=Paris]. ["anno XII"]

[The whole work is in 3 parts ("decads"): 1799 [An VII], pp. 1-44, pl. 1-10; 1801 [An X], [iv] + pp. 47-90, pl. 11-20; and 1804 [An XIII], [iv] + pp. 91-142, pl. 21-30.]

**Coulson, J. R., C. W. Sabrosky & I. Muller**

- 1965 Selected bibliography of North American Diptera. Pp. 1117-1547. In Stone, A., C. W. Sabrosky, W. W. Wirth, R. H. Foote & J. R. Coulson (eds.), A catalog of the Diptera of America north of Mexico. U.S. Dept. Agric., Agric. Handb. 276, iv + 1696 pp. (23 August)

**Cowan, C. F.**

- 1969 Notes on Griffith's *Animal Kingdom of Cuvier* (1824-1835). *J. Soc. Biblphy nat. Hist.* 5: 137-140.
- 1970 *Annotationes Rhopalocerologicae*. 70 p. Clunbury Press, Berkhamsted.
- 1971 On Guerin's *Iconographie*: particularly the insects. *J. Soc. Biblphy nat. Hist.* 6: 18-29.

**Crosskey, R. W.**

- 1965 A systematic revision of the Ameniinae (Diptera: Calliphoridae). *Bull. Br. Mus. nat. Hist. (Ent.)* 16: 33-140. (23 August)
- 1971 The type-material of Australasian, Oriental and Ethiopian Tachinidae (Diptera) described by Macquart and Bigot. *Bull. Br. Mus. nat. Hist. (Ent.)* 25: 251-305, 1 pl. (2 March)
- 1973a A revisionary classification of the Rutiliini (Diptera: Tachinidae), with keys to the described species. *Bull. Br. Mus. nat. Hist. (Ent.)*, suppl. 19, 167 pp. (22 February)
- 1973b A conspectus of the Tachinidae (Diptera) of Australia, including keys to the supraspecific taxa and taxonomic and host catalogues. *Bull. Br. Mus. nat. Hist. (Ent.)*, suppl. 21, 221 pp. (7 December)
- 1974 The British Tachinidae of Walker and Stephens (Diptera). *Bull. Br. Mus. nat. Hist. (Ent.)* 30: 267-308. (12 June)
- 1977 A revision of the Rhinophoridae (Diptera) and a revision of the Afrotropical species. *Bull. Br. Mus. nat. Hist. (Ent.)* 36: 1-66. (29 September)

**Curtis, J.**

- 1824- *British Entomology: Being illustrations and descriptions of the genera of insects found in Great Britain and Ireland: containing coloured figures from nature of the most rare and beautiful species, and in many instances of the plants upon which they are found*. Vol. 1, pls. 1-50 (1824); vol. 2, pls. 51-98 (1825); vol. 3, pls. 99-146 (1826); vol. 4, pls. 147-194 (1827); vol. 5, pls. 195-241 (1828); vol. 6, pls. 242-289 (1829); vol. 7, pls. 290-337 (1830); vol. 8, pls. 338-383 (1831); vol. 9, pls. 384-433 (1832); vol. 10, pls. 434-481 (1833); vol. 11, pls. 482-529 (1834); vol. 12, pls. 530-577 (1835); vol. 13, pls. 578-625 (1836); vol. 14, pls. 626-673 (1837); vol. 15, pls. 674-721 (1838); vol. 16, pls. 722-769 (1839). "Printed for the author" [by Richard and John E. Taylor], London.
- 1829- *A guide to an arrangement of British insects; being a catalogue of all the named species hitherto discovered in Great Britain and Ireland*.
- 1831 vi + 256 columns (128 pages) [+ 2 pp]. "Printed for the author" [by Richard and John E. Taylor], London.
- [1 June 1829: columns 1-128; 1830: columns 129-160; 1831: columns 161-256.]
- 1837 *A guide to an arrangement of British insects; being a catalogue of all the named species hitherto discovered in Great Britain and Ireland*.

Second edition, greatly enlarged. vi + 282 columns (141 pages) + 283-294 full pp. "Printed for the author" [by Richard and John E. Taylor], London.

**Day, M. C. & M. G. Fitton**

- 1977 Discovery in the Linnaean collection of type-material of insects described by Johann Reinhold Forster, with notes on the Hymenoptera. *Biol. J. Linnean Soc.* 9: 31-43.
- 1978 Re-curation of the Linnaean Hymenoptera (Insecta), with a reassessment of the taxonomic importance of the collection. *Biol. J. Linnean Soc.* 10: 181-198.

**De Geer, C.**

- 1773 *Mémoires pour servir à l'histoire des insectes*. Vol. 3, viii + 696 pp. P. Hesselberg, Stockholm.
- 1776 *Mémoires pour servir à l'histoire des insectes*. Vol. 6, viii + 523 pp., 30 pls. P. Hesselberg, Stockholm.

[Earlier volumes are considered non-binomial, and the names are available either from the German translation by Goeze (1782) or from the synopsis by Retzius (1783).]

**de Meijere, J. C. H.**

- 1904 Neue und bekannte Süd-Asiatische Dipteren. *Bijdr. Dierkd.* 17/18: 83-118, pl. 8, fig. 22-23. (31 December+)

**de Razoumowsky, G.**

- 1789 *Histoire naturelle du Jorat et de ses environs; et celle des trois lacs de Neufchatel, Morat et Bienne; précédées d'un essai sur le climat, les productions, le commerce, les animaux de la partie du Pays de Vaud ou de la Suisse Romande, qui entre dans le plan de cet ouvrage*. Vol. 1, xvi + 322 pp., 3 pls.; vol. 2, 238 pp., 3 pls. Mourer, Lausanne.  
[Insects, vol. 1: 289-301 & 133-239 (catalog & new species)]

**de Villers, C. J.**

- 1789a *Caroli Linnaei entomologia, faunae suecicae descriptionibus aucta; DD. Scopoli, Geoffroy, de Geer, Fabricii, Schrank, &c. speciebus vel in systemate non enumeratis, vel nuperrime detectis, vel speciebus Galliae Australis locupletata, generum specierumque rariorū iconibus ornata*. Vol. 3, 657 pp., 4 pls. Piestre & Delamollière, Lugduni [=Lyon].

- 1789b *Caroli Linnaei entomologia, faunae suecicae descriptionibus aucta; DD. Scopoli, Geoffroy, de Geer, Fabricii, Schrank, &c. speciebus vel in systemate non enumeratis, vel nuperrime detectis, vel speciebus Galliae Australis locupletata, generum specierumque rariorū iconibus ornata*. Vol. 4, 556 + ccxiii pp., 1 pl. Piestre & Delamollière, Lugduni [=Lyon].

**Dear, J. P.**

- 1977 A revision of Australian Rhiniinae (Diptera: Calliphoridae). *Aust. J. Zool.* 25: 779-826. (October)

1979 A revision of the Toxotarsinae (Diptera: Calliphoridae). *Pap. Avuls. Zool.* 32: 145-182. (20 February)

1985a Calliphoridae (Insecta: Diptera). *Fauna of New Zealand* 8, 86 pp. (24 February)

1985b A revision of the New World Chrysomyini (Diptera: Calliphoridae). *Revta bras. Zool.* 3: 109-169. (28 June)

**DerkSEN, W. & U. Scheiding [Scheiding-Göllner]**

1963- *Index Litteraturae Entomologicae. Serie II: Die Welt-Literatur über die gesamte Entomologie von 1864 bis 1900.* Band I A-E, xii + 697 pp. (1963); II F-L, 678 pp. (1965); III M-R, 528 pp. (1968); IV S-Z, 482 pp. (1972); V Register [by R. Gaedike], 238 pp. (1975). Deutsche Akademie der Landwirtschaftswissenschaften, Berlin.

**Descourtiz, T.**

1826 Description de quelques Insectes nouveaux découverts en France en 1825. *Mém. Soc. Linn. Paris* 5: 156-163, 1 pl.

**Disney, R. H. L.**

1982 Confirmation of a suspected synonym in the genus *Borophaga* (Diptera: Phoridae). *Ent. mon. Mag.* 119: 115. (9 August)

**Doleschall, C. L.**

1859 Derde bijdrage tot de kennis der dipteren fauna van Nederlandsch Indië. *Natuurkd. Tijdschr. Ned.-Indië* [1858] 17: 73-128. (14 April)

[The 14 April 1859 date is recorded in the Society's Proceedings (Evenhuis, *in litt.*.)]

**Donovan, E.**

1792- *The natural history of British Insects; explaining them in their several*

1813 *states, with the periods of their transformations, their food, oeconomy &c. Together with the history of such minute Insects as require investigation by the microscope. The whole illustrated by coloured figures, designed and executed from living specimens.* Vol. 1, 80 + 20

pp., pls. 1-36 (1792); 2, 96 + 6, pls. 37-72 (1793); 3, 98 + 7 pp., pls. 73-108 (1794); 4, 96 + 6 pp., pls. 109 - 144 (1795); 5, 110 + 6 pp., pls.

145 - 180 (1796); 6, 86 + 6 pp., pls. 181 - 216 (1797); 7, 96 + 6 pp., pls. 217 - 252 (1798); 8, 88 + 6 pp., pls. 253 - 288 (1799); 9, 78 + 4

pp., pls. 289 - 324 (1800); 10, 95 + 14 pp., pls. 325 - 360 (1801); 11, ix + 100 + 7 pp., pls. 361 - 396 (1806); 12, 102 + 5 pp., pls. 397 - 432

(1807); 13, 74 + 4 pp., pls. 433 - 468 (1808); 14, 90 + 5 pp., pls. 469 - 504 (1810); 15, 83 + 5 pp., pls. 505 - 540 (1811); and 16, 91 + 10

pp., pls. 541 - 576 (1813). Rivington, London.

1805 *An epitome of the natural history of the insects of New Holland, New Zealand, New Guinea, Otaheite, and other islands in the Indian, Southern, and Pacific oceans: including the figures and descriptions of one hundred and fifty-three species of the more splendid, beautiful, and interesting insects, hitherto discovered in those countries, and which for the most part have not appeared in the works of any preceding author. The figures are correctly delineated from speci-*

*mens of the insects; and with the descriptions are arranged according to the Linnaean system, with reference to the writings of Fabricius and other entomologists.* Rivington, London. iv + [167] pp. + [41] pls. (after 20 July)

[This work is unpaginated and the plates and figures are unnumbered. It appears that copies of the printing proof were distributed as there are no paginated copies known. The figures are distinguished in each plate and associated legend by a series of diamond-shaped marks in place of figure numbers. The Diptera are described on the 166th and 167th pages and are figured on the 41st plate. The dedication page is dated 20 July and, noting that the work was apparently not finalized with pagination, it is assumed here that the issue date was soon after the 20 July date.]

### Drury, D.

- 1770- *Illustrations of natural history. Wherein are exhibited upwards of two hundred and forty figures of exotic insects, according to their different genera; very few of which have hitherto been figured by any author, being engraved and coloured from nature, with the greatest accuracy, and under the author's own inspection, on fifty copper-plates. With a particular description of each insect: interspersed with remarks and reflections on the nature and properties of many of them. To which is added, a translation into French.* Vol. 1, xxviii + 130 pp., 4 figs., 50 pls. (1770); vol. 2, vii + 90 pp., 50 pls., and index to Vols. 1-2 (4 unnumbered pages) (1773). "Printed for the author; sold by White," London.

[The scientific names are found only in the index.]

### Dryander, J.

- 1796- *Catalogus bibliothecae historico-naturalis Josephi Banks regi a consiliis intimis, Baroneti, Balnei equitis, Regiae Societatis praesidis, caet.* Tomus I, *Scriptores Generales*, vii + 309 [+14 index] pp. (1798); II, *Zoologi*, xx + 578 [+ 30 index] pp. (1796); III, *Botanici*, xxiii + 656 [+38 index] pp. (9 November 1797); IV, *Mineralogi*, ix + 390 [+26 index] pp. (1799); V, *Supplementum et Index Auctorum*, 514 pp. (6 November 1800). Gul. Bulmer et Soc., Londini [=London]

### Dupuis, C.

- 1986 Dates de publication de l'"Histoire naturelle générale et particulière des crustacés et des insectes" (1802-1805) par Latreille dans le "Buffon de Sonnini". *Ann. Soc. Ent. Fr. (N.S.)* 22: 205-210. (30 June)

### Dušek, J. & P. Láska

- 1967 Versuch zum Aufbau eines natürlichen Systems mitteleuropäischer Arten der Unterfamilie Syrphinae (Diptera). *Acta Sci. Nat. Brno* 1: 349-390. (16 March)

### Emden, F. I. van

- 1939 Muscidae: A. — Muscinae and Stomoxydinae. *Ruwenzori Exped. 1934-35* 2: 49-89. British Museum (Natural History), London. (22 April)
- 1965 Diptera. Vol. 7. Muscidae, part 1. xiv + 647 pp., 2 pls. In Sewell, R.B.S. and M. L. Roonwal (eds.), *The fauna of India and the adjacent countries*. Government of India, Delhi. (June)

**Erichson, W. F.**

- 1851 Hymenoptera, Diptera and Neuroptera, pp. 60-69. In E. Ménétriés, Insecten, pp. 43-76, plate 3. In Middendorf, A. T. von. *Reise in den düssersten Norden und Osten Sibiriens während der Jahre 1843 und 1844 mit allerhöchster Genehmigung auf Veranstaltung der Kaiserlichen Akademie der Wissenschaften zu St. Petersburg ausgeführt und in Verbindung mit vielen Gelehrten herausgegeben (von Dr. A. Th. v. Middendorf)*. Band II, Zoologie. Theil 1. Wirbellose Thiere: Annulaten. Echinodermen. Insecten. Krebse. Mollusken. Parasiten. Bearbeitet von F. Brandt, W. F. Erichson, Seb. Fischer, E. Grube, E. Ménétriés, A. Th. v. Middendorf. 516 pp., 32 plates. St. Petersburg.

**Eschscholtz, J. F.**

- 1822 *Entomographien*. Erste Lieferung. Reimer, Berlin. 128 + III pp., 2 pls.  
 [This was a preprint issued in advance of the article that appeared in *Naturwiss. Abhandl. Dorpat*; see Eschscholtz, 1823. The preface is dated 24 May.]

- 1823 Entomographien. *Naturwiss. Abhandl. Dorpat*. 1: 57-186, 2 pls.

**Estlund, O.**

- 1796 Entomologiske Anmärkningar hörande till Fauna Suecica. *K. Vetenskapsakad. Nyd Handl.* 17: 126-132.

**Evenhuis, N. L.**

- 1988 Localities of Fabrician types collected by Labillardière on the voyage of the *Recherche* and *Esperance*, with special reference to the Diptera. *Arch. Nat. Hist.* 15: 185-196.

**Evenhuis, N. L., F. C. Thompson, A. C. Pont & B. L. Pyle**

- 1989 Literature cited. Pp. 809-991. In Evenhuis, N. L. (ed.), *Catalog of the Diptera of the Australasian and Oceanian Regions*. 1155 pp. Bishop Museum Press & E. J. Brill, Honolulu & Leiden. (23 August)  
 [Bishop Museum Special Publication 86.]

**Eversmann, E.**

- 1834 Diptera Wolgam fluvium inter et montes Uralenses observata. *Bull. Soc. imp. Nat. Moscou* 7: 420-432.

**Fabricius, J. C.**

- 1775 *Systema entomologiae, sistens insectorum classes, ordines, genera, species, adiectis synonymis, locis, descriptionibus, observationibus*. [32] + 832 pp. Kortii, Flensbvrgi et Lipsiae [=Flensburg and Leipzig]. (30 April)  
 [The I.C.Z.N. (1958) determined the publication for this work as 17 April on the assumption that it first appeared at the Leipzig Easter book fair with the 17th being the Monday after Easter. However, the Leipzig book fair was held two weeks after Easter (Sayre 1959: 4).]

- 1777 *Genera insectorum eorumque characteres natvrales secundvm nvmervm, figvram, sitvm et proportionem omnivm partivm oris adiecta mantissa specierum nuper detectarvm*. [16+] 310 pp. M. F. Bartschii, Chilonii [=Kiel].

[Horn & Schlenkling (1928: 334) gave 1776 as the date without comment. Hagen (1862: 220) gave 1777 following Engelmann. The title page bears no date. The preface is dated 26 December 1776. Given that late date, we follow Hagen and date this work as appearing in 1777.]

- 1779 *Reise nach Norwegen mit Bemerkungen aus der Naturhistorie und Oekonomie.* Ixiv + 388 pp. [+ 12] pp. Bohn, Hamburg.
- 1781 *Species insectorum exhibentes eorum differentias specificas, synonyma, avctorum, loca natalia, metamorphosin adiectis observationibus, descriptionibus.* Vol. 2, 517 pp. C. E. Bohnii, Hambvrgi et Kilonii [=Hamburg and Kiel]. (March)  
 [Shown at the Leipzig Easter book fair (29 April) (Cobres 1782: 820). Also listed in *Allgemeines Verzeichniss Neuer Bücher 6: 181* (Evenhuis *in litt.*).]
- 1787 *Mantissa insectorum sistens species non per detectas adiectis synonymis, observationibus, descriptionibus, emendationibus.* Vol. 2, 382 pp. C. G. Proft, Hafniae [=Copenhagen]. (27 December)  
 [The titles given for volumes 1 and 2 differ. The above is the title given for volume 2. The title given in Coulson *et al.* (1965: 1222) is incomplete; that in Evenhuis *et al.* (1989: 863) is volume 1, not 2. The date is from *Minerva* 10: 440 (Evenhuis *in litt.*).]
- 1794 *Entomologia systematica emendata et aucta. Secundum classes, ordines, genera, species adiectis synonymis, locis, observationibus, descriptionibus.* Vol. 4, [6] + 472 + [5] pp. C. G. Proft, Hafniae [=Copenhagen].
- 1798 *Supplementum entomologiae systematicae.* ii + 572 pp. Storch, Hafniae [= Copenhagen]. (20 May)
- 1805 *Systema antiatorum secundum ordines, genera, species adiecta synonymis, locis, observationibus, descriptionibus.* 373 pp., + 30 pp. Reichard, Brunsvigae [=Brunswick].

**Fabricius, O.**

- 1780 *Favna Groenlandica, systematice sistens animalia groenlandiae occidentalis hactenus indagata, quoad nomen specificum, triviale, vernaculaeque; synonyma avctorum plurim, descriptionem, locum, vicum, generationem, mores, usum, capturamque singuli, prout degendi occasio fuit, maximaque parte secundum proprias observationes.* xvi + 452 pp., 1 plate. Ioannis Gottlob Rothe, Hafniae et Lipsiae [=Copenhagen & Leipzig]. (before 13 August)  
 [The date is from Dawson (1958: 333, *The Banks letter ...*) (Evenhuis *in litt.*).]

**Fallén, C. F.**

- 1810 *Specim. entomolog. novam Diptera disponendi methodum exhibens.* 26 pp., 1 pl. Berlingianis, Lundae [=Lund]. (9 June)
- 1817 Beskrifning öfver de i Sverige funna fluge arter, som kunna föras till slägget *Musca*. Första afdelningen. *K. Svenska. Vetenskapsakad. Handl.* (3) 1816: 226-254.
- 1820- *Monographia Muscidum Sveciae.* Pp. 1-12 (13 June 1820), 13-24 (13 June 1820), 25-40 (14 June 1820); Pp. 41-48 (6 June 1821); pp. 49-56 (26 April 1823), 57-64 (31 May 1823); pp. 65-72 (9 December 1824); pp. 73-80 (23 April 1825), 81-94 (18 June 1825). Berlingianis, Lundae [=Lund]

**Fan, Zi-de**

[Correct form of first name is "Zi-de" according to Dr. Fan, but it also appears as "C.T.," "Tse-Teh," "Tze-Teh," or "Zide".]

- 1965 Key to species of common synanthropic flies in China. xv + 330 pp., 40 pls. Peking. [In Chinese.] (September)

- 1978 A new fly of the genus *Musca* from Tibet, China (Diptera: Muscidae). *Acta ent. sin.* 21: 329-331. [In Chinese with English summary.] (August)

**Ferrar, P.**

- 1987 A guide to the breeding habits and immature stages of Diptera Cyclorrhapha. *Entomonograph* 8, 907 pp. E. J. Brill/Scandinavian Science Press, Leiden & Copenhagen.

[Issued in two volumes: volume 1 (text), pp. 1-478, volume 2 (figures), pp. 479-907.]

**Forskål, P.**

- 1775 *Descriptiones animalium avium, amphibiorum, piscium, insectorum, vermium; quae in itinere orientali observavit Petrus Forskål. Post mortem auctoris edidit Carsten Niebuhr. Adjuncta est materia medica kahirina atque tabula maris rubri geographica.* 19 + 34 + 164 pp. Möller, Havniae [=Copenhagen]. (3 June)

Forsskål, J. P.— see Linnaeus 1764a

**Forster, J. R.**

- 1770 *Catalogue of British Insects.* 16 pp. Eyres, Warrington.

- 1771 *Novae species insectorum. Centuria I.* viii + 100 pp. Davies & White, London. (1 December)

- 1788 *Enchiridion historiae naturali inserviens, quo termini et delineationes ad avium, piscium, insectorum et plantarum adumbrationes intelligendas et concinnandas, secundum methodum systematis Linnaeani continentur.* [15 +] 224 pp. Hemmerde & Schwetschke, Halae [= Halle]. (27 March)

- 1794 *Enchiridion historiae naturali inserviens, quo termini et delineationes ad avium, piscium, insectorum et plantarum adumbrationes intelligendas et concinnandas, secundum methodum systematis Linnaeani continentur. Editio altera emendatior [2nd edition].* 8 + 224 pp. Stewart, Edinburgi [= Edinburgh].

**Fourcroy, A. F. de**

- 1785 *Entomologia parisiensis; sive Catalogus Insectorum quae in Agro Parisiensi reperiuntur; Secundum methodum Geoffroeanam in sectio-nes, genera & species distributus: Cui addita sunt nomina trivialia & fere trecentae novae Species.* viii + 544 pp. Aedibus Serpentineis, Parisiis [=Paris]. (before 1 August)

[As two volumes, pp. 1-231 & 233-544. The introduction written by Fourcroy makes it clear that all the species in this work should be properly attributed to Geoffroy who provided the names and descriptions (also in Hagen (1862: 246)).]

**Frisch, J. L.**

- 1728 *Beschreibung von allerley Insecten in Teutsch-Land, nebst nützlichen Anmerckungen und nöthigen Abbildungen von diesem kriechenden und fliegenden inländischen Gewürme, zur Bestätigung und Fortsetzung der gründlichen Entdeckung, so einige von der Natur dieser*

- Creaturen heraus gegeben, und zur Ergänzung und Verbesserung der andern.* Volume 7. [10+] 31 pp., 22 pls. Nicolai, Berlin.
- 1732 *Beschreibung von allerley Insecten in Teutsch-Land, nebst nützlichen Anmerckungen und nöthigen Abbildungen von diesem kriechenden und fliegenden inländischen Gewürme, zur Bestätigung und Fortsetzung der gründlichen Entdeckung, so einige von der Natur dieser Creaturen heraus gegeben, und zur Ergänzung und Verbesserung der andern.* Volume 10. [7+] 25 [+ 5] pp., 21 pls. Nicolai, Berlin.
- Fuesslin, J. C.**
- 1775 *Verzeichniss der ihm bekannten Schweizerischen Inseckten mit einer ausgemahlten Kupfertafel: nebst der Ankündigung eines neuen Insecten Werks.* xii + 62 pp., 1 pl. Fuesslin, Zürich & Steiner, Winterthur. (11 November)
- [We have used the spelling "Fuesslin" as that is how the name appears on the title page and preface. Fuessly is used by Percheron (1837: 121) and Sherborn (1902: xxv). Gilbert (1977: 130) used Füssley and Horn & Schenkling (1928: 390) used Flüssly but gave Füssly, Fuessly, Flüssli and Fuesslin as alternative spellings. Hagen (1862: 257) used Flüssly, but wrote "The author is always called Fuessly, only in Archiv he wrote Füssly and alternatively Fuessly!"]
- 1778 Dr. Sulzers abgekürzter Geschichte der Insecten Etc. Nachricht davon, nebst Anmerkungen und Zusätzen. *Mag. Liebhaber. Ent.* 1: 141-242.
- Gaedike, R. & O. Smetana**
- 1978 Ergänzungen und Berichtigungen zu Walter Horn und Sigmund Schenkling: Index Litteraturae Entomologicae, Serie I, die Welt-Literatur über die gesamte Entomologie bis inklusive 1863. Teil I: A—K. *Beitr. Ent., Berlin* 28: 329-436.
- 1984 Ergänzungen und Berichtigungen zu Walter Horn und Sigmund Schenkling: Index Litteraturae Entomologicae, Serie I, die Welt-Literatur über die gesamte Entomologie bis inklusive 1863. Teil I: L—Z. *Beitr. Ent., Berlin* 34: 167-291.
- Gagné, R. J.**
- 1981 *Chrysomya* spp., old world blow flies (Diptera: Calliphoridae) recently established in the Americas. *Bull. Ent. Soc. Amer.* 27: 21-22.
- Geoffroy, E. L.**
- 1762 *Histoire abrégée des insectes qui se trouvent aux environs de Paris; dans laquelle ces animaux sont rangés suivant un ordre méthodique.* Volume 2. 690 pp., pls 11-22. Durand, Paris.  
[First edition was published anonymously.]
- 1785 [new species in Fourcroy 1785 (q.v.)]
- Germar, E. F.**
- 1825 *Fauna Insectorum Europae.* Fasc. 10, 25 pls. Kümmelii, Halae [=Halle].  
[See Rautenberg (1957) for details on this series.]
- 1839 Die versteinerten Insecten Solenhofens. *Nova Acta Acad. Caes. Leop. Car.* 19: 187-222, pl. 21-23. (June)

**Gilbert, P.**

- 1977 *A compendium of the biographical literature on deceased entomologists.* [viii +] 455 pp. British Museum (Natural History), London.

**Gimmerthal, B. A.**

- 1834 Observations de quelques nouvelles espèces de Diptères, accompagnées de recherches sur la métamorphose de quelques autres. *Bull. Soc. Imp. Moscou* 7: 98-121, pl. 1-2.

**Girschner, E.**

- 1894 Beitrag zur Systematik der Musciden. *Berl. ent. Ztschr.* 38 [1893]: 297-312. (January)

- 1896 Ein neues Musciden-System auf Grund der Thoracalbeborstung und der Segmentierung des Hinterleibes. *Illustr. Wochenschr. Entomol.* 1: 12-16, 30-32, 61-64, 105-112.

- 1897 Über die Postalar-Membran (Schüppchen, Squamulae) der Dipteren. *Illustr. Wochenschr. Entomol.* 2: 534-539, 553-559, 567-571, 586-589, 603-607, 641-645, 666-670.

**Gistl, J. N. F. X.**

- 1835 Neue Arten von Diptern aus der Umgegend von München, benannt und beschrieben von Meigen, aufgefunden von Dr. J. Waltl, Professor der Naturgeschichte in Passau. *Faunus, Zeitschrift für Zoologie und vergleichende Anatomie* [1], 2: 66-72. (May)

- 1837a Kritische Revisionen und Ergänzungen zu Schrank's "Enumeratio Insectorum Austriae, Fauna boica, u.s.w." Aus dem bisher noch ungedruckten literarischen Nachlass des sel. Schrank mitgetheilt von Gistl. *Faunus, Zeitschrift für Zoologie und vergleichende Anatomie* (2), 1 (1): 5-19. (May)

- 1837b Franz von Paula Schrank. Gallerie denkwürdiger Naturforscher. *Faunus, Zeitschrift für Zoologie und vergleichende Anatomie* (2), 1 (1), Beiblatt 1: 1-4; 1 (2), Beiblatt 2: 5-8.

[This also includes, on page 8, the start of an unfinished bibliography of Schrank's works.]

- 1856 *Die Mysterien der europäischen Insectenwelt. Ein geheimer Schlüssel für Sammler aller Insecten-Ordnungen und Stände behufs des Fangs, des Aufenthalts-Orts, der Wohnung, Täg- und Jahreszeit u.s.w. Zum ersten Male nach 25jährigen eigenen Erfahrungen zusammengestellt und herausgegeben.* 12 + 532 pp. Dannheimer, Kempten. (18 February)

**Gmelin, J. F.**

- 1790 *Caroli a Linné, Systema naturae per regna tria naturae, secundum classes, ordines, genera, species; cum caracteribus, differentiis, synonymis, locis.* Editio decima tertia, aucta, reformata [= Ed. 13.] Vol. 1: Regnum Animale, Pt. 5, Pp. 2225-3020. G. E. Beer, Lipsiae [=Leipzig]. (6 December)

**Goedart, J.**

- 1662a *Metamorphosis et historia naturalis insectorum*. [Volume 1.] [29+] 236 pp. Fierensium, Medioburgi [=Middleburg].
- 1662b *Metamorphosis et historia naturalis insectorum*. [Volume 2.] [30+] 259 pp. Fierensium, Medioburgi [=Middleburg].
- 1662c *Metamorphosis et historia naturalis insectorum*. [Volume 3.] [14+] 159 pp. Fierensium, Medioburgi [=Middleburg].  
[See Lister for English translation.]

**Goeze, J. A. E.**

- 1782 *Des Herrn Baron Karl Degeer Königlichen Hofmarschalls & c. & c. Abhandlungen zur Geschichte der Insekten aus dem Französischen übersetzt und mit Anmerkungen herausgegeben*. Volume 6. [6+] 200 pp., 30 pls. Raspe, Nürnberg. (31 August)

**Goffe, E. R.**

- 1946 The Syrphidae (Diptera) of Moses Harris, 1776. *Ent. mon. Mag.* 82: 67-86 (30 March)
- 1951 The Syrphidae (Diptera) described by Moses Harris in 1776. *Entomologist* 84: 195-204.

**Graham, M. W. R. de V.**

- 1979 "Ambulator": Francis Walker, English entomologist (1809-1874). *Entomologist's Gaz.* 30: 7-20, pls 2-3. (24 February)

**Gravenhorst, J. L. C.**

- 1807 *Vergleichende Uebersicht des Linneischen und einiger neuern zoologischen Systeme nebst dem eingeschalteten Verzeichnisse der zoologischen Sammlung des Verfassers und den Beschreibungen neuer Thierarten, die in derselben vorhanden sind*. xvi + 476 pp. Dieterich, Goettingen. (19 March)

**Greenberg, B.**

- 1971 *Flies and Disease*. [Volume 1:] *Ecology, classification and biotic associations*. xii + 856 pp. Princeton University Press, Princeton.

**Griffin, F. J.**

- 1938 On the date of publication of Latreille (*in Sonnini's Buffon*), An X [sic], Hist. Nat. Gén. Partic. Crust. Ins. 3. *J. Soc. Biblphy nat. Hist.* 1: 157. (25 May)

**Griffith, E. & E. Pidgeon**

- 1832 *The animal kingdom arranged in conformity with its organization, by the Baron Cuvier, member of the Institute of France, & c. & c. & c. with supplementary additions to each order*. Volume 15. The class insecta arranged by the Baron Cuvier, with supplementary additions to each order, by Edward Griffith, F.L.S., A.S., & c. and Edward Pidgeon, Esq. and notices of new genera and species by George Gray, Esq. Volume 2. 796 pp., many plates. Whittaker, Treacher and Co., London. (December)

[Diptera are treated on pages 683-780 and on plates 62, 114, 122, 125 & 128. For dating information, see Cowan 1969.]

### Grossinger, J. B.

- 1794 *Universa historia physica regni hungariae secundum tria regna naturae digesta*. Tomus IV. Regni animalis Pars IV. Entomologia, sive historia insectorum. xiv + 416 + xix pp., 1 pl. Weber, Posonii et Comaromii [=Posen & Komárno].

### Guérin-Méneville, F. E.

- 1831 Insects, plates No. 1-No. 21. In Duperrey, L. I. (ed.), *Voyage autour du monde, exécuté par ordre du Roi, sur la corvette de sa majesté, La Coquille, pendant les années 1822, 1823, 1824 et 1825, sous le Ministère et conformément aux instructions de S. E. M. Le Marquis de Clermont-Tonnerre, Ministre de la Marine; et publié sous les auspices de son excellence M<sup>gr</sup> Le C<sup>te</sup> de Charbrol, Ministre de la Marine et des Colonies. Histoire naturelle, Zoologie. Atlas*. A Bertrand, Paris. '1826'.

[The Diptera are on plates 20 (25 July 1831) and 21 (5 September 1831).]

- 1838 Histoire naturelle des crustacés, arachnides et insectes, recueillis dans le voyage autour du monde de la corvette de sa Majesté, La Coquille, exécuté pendant les années 1822, 1823, 1824 et 1825, sous le commandement du Capitaine Duperrey. Première division. Crustacés, arachnides et insectes. xii + 319 pp. In Duperrey, L. I. (ed.), *Voyage autour du monde, exécuté par ordre du Roi, sur la corvette de sa Majesté, La Coquille, pendant les années 1822, 1823, 1824 et 1825, sous le Ministère et conformément aux instructions de S. E. M. Le Marquis de Clermont-Tonnerre, Ministre de la Marine; et publié sous les auspices de son excellence M<sup>gr</sup> Le C<sup>te</sup> de Charbrol, Ministre de la Marine et des Colonies. Histoire naturelle, Zoologie*. Vol. 2, Part 2. H. Bertrand, Paris. '1826'.

### Guiglia, D.

- 1957 I tipi di Ditteri di Camillo Rondani e di altri autori appartenenti alle collezioni del Museo Civico di Storia Naturale di Genova. *Annali Mus. civ. Stor. nat. Giacomo Doria* 69: 193-207.

Guillou — see Le Guillou

### Gurney, A. B.

- 1956 On the rendering of Charles De Geer's surname. *Bull. Brooklyn ent. Soc.* 51: 127-136.

### Hagen, H.

- 1844 Nachricht über die entomologischen Sammlungen in Norwegen, Schweden und Dänemark. *Stettin. ent. Ztg* 5: 68-70, 75-80, 130-131. (February, March, April)

- 1861 On Moses Harris' work. *Ent. weekly Intellig.* 10: 181-183. (7 September)

- 1862 *Bibliotheca Entomologica. Die Litteratur über das ganze Gebiet der Entomologie bis zum Jahre 1862. Erster Band, A—M.* xii + 566 pp. W. Engelmann, Leipzig

- 1863 *Bibliotheca Entomologica. Die Litteratur über das ganze Gebiet der Entomologie bis zum Jahre 1862. Zweiter Band, N—Z mit einem systematischen Sachregister.* 512 pp. W. Engelmann, Leipzig

**Haliday, A. H.**

- 1833 Catalogue of the Diptera occurring about Holywood in Downshire. *Ent. Mag. (London)* 1: 147-180. (January)
- 1836 Notes, &c. upon Diptera. 2. Characters of some undescribed species of the Family Muscidae. *Ent. Mag. (London)* 4: 149-152. (October)
- 1851 Sendschreiben von Alexis H. Haliday an C. A. Dohrn über die Dipteren der in London befindlichen Linnéischen Sammlung. *Stettin. ent. Ztg* 12: 131-145.

**Harpaz, I.**

- 1973 Early entomology in the Middle East. Pp. 21-36. In Smith, R. F., T. E. Mittler & C. N. Smith [eds], *History of Entomology*. vii + 517 pp. Annual Reviews Inc., Palo Alto, California.

**Harris, M.**

- 1776- *An exposition of English insects, with curious observations and remarks, wherein each insect is particularly described; its parts and properties considered; the different sexes distinguished, and the natural history faithfully related. The whole illustrated with copper plates, drawn, engraved, and coloured, by the author.* Decad I, pp. 1-40, 2 pls. + pls. 1-10 (1776a); Decad II, pp. 41-72, pls. 11-20 (?1776b); Decads III, IV, V, pp. 73-99, 100-138, 139-166, pls. 21-30, 31-40, 41-50 + 1 pl. (?1780). Robson Co., London "1776".
- 1782 *An exposition of English insects including the several classes of Neuroptera, Hymenoptera, & Diptera, or bees, flies, & libellulae. Exhibiting on 51 copper plates near 500 figures, accurately drawn, & highly finished in colours, from Nature. The whole minutely described, arranged, & named, according to the Linnean system, with remarks. The figures of a great number of moths, not in the Aurelian collection, formerly published by the same author, and a plate with an explanation of colours, are likewise given in the work.* viii + 166 pp. + index (4 unnumbered pages), 50 + 2 pls. White & Robson, London.

**Harris, T. W.**

- 1833 VIII. Insects [of Part IV. Catalogues of Animals and Plants]. Pp. 566-595. In Hitchcock, E., *Report on the geology, mineralogy, botany, and zoology of Massachusetts. Made and published by order of the government of that state: In four parts: Part I. Economical Geology. Part II. Topographical Geology. Part III. Scientific Geology. Part IV. Catalogues of animals and plants. With a descriptive list of the specimens of rocks and minerals collected for the government. Illus-*

*trated by numerous wood cuts and an atlas of plates.* xii + 700 pp. + Atlas, 17 pls. J. S. & C. Adams, Amherst, Massachusetts.

- 1835 VIII. Insects [of Part IV. Catalogues of Animals and Plants]. Pp. 553-602. *In Hitchcock, E., Report on the geology, mineralogy, botany, and zoology of Massachusetts. Made and published by order of the government of that state: In four parts: Part I. Economical Geology. Part II. Topographical Geology. Part III. Scientific Geology. Part IV. Catalogues of animals and plants. With a descriptive list of the specimens of rocks and minerals collected for the government. Illustrated by numerous wood cuts and an atlas of plates.* Ed. 2, [iv] + xii + 702 pp. + Atlas, 17 pls. J. S. & C. Adams, Amherst, Massachusetts.
- 1841 *A report on the insects of Massachusetts, injurious to vegetation.* 459 pp. Cambridge, Massachusetts.

- 1869 Descriptions of insects selected from the manuscripts of Dr. Harris. Pp. 325-336. *In Scudder, S. H. (ed.), Entomological correspondence of Thaddeus William Harris, M.D. Boston Soc. Nat. Hist. Occas. Pprs 1: 1-375, 4 pls.*

### Hartig, T.

- 1838 Ueber die parasitischen Zweiflügler des Waldes. *[Hartig's] Jahressber. Fortschr. Forstwiss. Naturk.* 1(2): 275-306.

### Hayek, C. M. F. von

- 1985 On the type of material of the species of Coleoptera described from the Drury Collection by D. Drury and J. C. Fabricius with notes on some Coleoptera from the Milne collection preserved in the British Museum (Natural History). *Arch. Nat. Hist.* 12: 143-152.

### Hennig, W.

- 1955- Muscidae [Fam.] 63b. *In Lindner, E. (ed.), Die Fliegen der palaearktischen Region* 7(2), 1110 pp., 33 pls. Schweizerbart, Stuttgart.

[The dates and lieferung numbers are: 182, pp. 1-48, pls. 1-3 (9 September 1955); 185, pp. 49-96, pls. 4-6 (17 October 1955); 194, pp. 97-144 (24 October 1956); 197, pp. 145-192, pls. 7-9 (7 August 1957); 199, pp. 193-232 (24 January 1958); 204, pp. 233-286, pls. 10-12 (19 February 1959); 205, pp. 289-336, pls. 13-15 (3 June 1959); 207, pp. 337-384, pls. 16-18 (28 September 1959); 209, pp. 385-432, pls. 19-20 (30 March 1960); 213, pp. 433-480, pls. 21-22 (9 December 1960); 215, pp. 481-528, pls. 23-26 (14 March 1961a); 217, pp. 529-576, pls. 27-30 (12 May 1961c); 223, pp. 577-624, pls. 31 (1 February 1962); 225, pp. 625-672 (26 January 1962); 227, pp. 673-720, pls. 32-33 (6 June 1962); 229, pp. 721-768 (15 June 1962); 233, pp. 769-816 (25 January 1963); 234, pp. 817-864 (10 July 1963); 241, pp. 865-912 (19 November 1963); 242, pp. 913-960 (19 November 1963); 248, pp. 961-1008 (24 June 1964); 249, pp. 1009-1056 (24 June 1964); 253, pp. 1057-1100 (30 November 1964).]

- 1961b Bericht über die Untersuchung der Typen einiger der von Rondani beschriebenen Arten aus der Familie Muscidae. *Beitr. Ent.* 11: 225-229. (March)

- 1966- Anthomyiidae [Fam.] 63a. *In Lindner, E. (ed.), Die Fliegen der palaearktischen Region* 7(1), lxxvii + 974 pp., 114 pls. Schweizerbart, Stuttgart.

[The dates and lieferung numbers are: 315, lxxvii pp. (4 March 1976); 262, pp. 1-48, pls. 1-3 (31 January 1966); 268, pp. 49-96 (16 December 1966); 271, pp. 97-144, pls. 7-9 (17 April 1967); 272, pp. 145-192, pls. 10-14 (12 October 1967); 276, pp. 193-240, pls. 15-19 (28 November 1968); 278, pp. 241-288, pls. 20-26 (30 July 1969); 280, pp. 289-328, pls. 27-33 (24 November 1969); 282, pp. 329-376, pls. 34-44 (30 December

1970); 283, pp. 377-424, pls. 45-55 (30 December 1970); 294, pp. 425-472, pls. 56-61 (22 December 1972); 295, pp. 473-512, pls. 62-69 (24 July 1973); 296, pp. 513-592, pls. 70-77 (24 July 1973); 297, pp. 593-680, pls. 78-85 (24 July 1973); 306, pp. 681-776 (30 September 1974); 307, pp. 777-872 (30 September 1974); 308, pp. 873-920, pls. 86-108 (30 September 1974); 314, pp. 921-974, pls. 109-114 (4 March 1976).]

### **Henriksen, K. L.**

- 1922 *Oversigt over Dansk entomologis historie. [Part.] Ent. Meddr* 15: 49-96. (7 November)

- 1940 Frederiksdal og Otto Friedrich Müller. *Ent. Meddr* 22: 5-6. (October)

### **Herbst, J. F. W.**

- 1787 *Kurze Einleitung zur Kenntniss der Insekten, für Ungeübte und Anfänger. Vol. 3*, 200 pp., pls. 54-87. Lange, Berlin & Stralsund.

[Also issued as vol. 8 of Borowski, G. H. (ed.), *Gemeinnützige Naturgeschichte des Thierreichs*. Lange, Berlin & Stralsund. Hagen gives separate authorship to the plates as: Sotzmann, D. F., *Natürliche Abbildungen der merkwürdigsten Insecten*.]

### **Herting, B.**

- 1972 Die Typenexemplare der von Meigen (1824-1838) beschriebenen Raupenfliegen (Dipt. Tachinidae). *Stuttgart. Beitr. Naturk.* 243, 15 pp.

- 1984 Catalogue of Palearctic Tachinidae (Diptera). *Stuttgart. Beitr. Naturk. (A, Biol.)* 369, 228 pp.

### **Higgins, L. G.**

- 1963 *Entomologia Carniolica: J. A. Scopoli, 1763. J. Soc. Biblphy nat. Hist.* 4: 167-169.

### **Ho, Ch'i**

- 1938 The significance of the female terminalia of house-flies as a grouping character. *Ann. trop. Med. Parasit.* 32: 287-312. (12 October)

### **Horn, W. & I. Kahle**

- 1935- Über entomologische Sammlungen, Entomologen & Entomo-

- 1937 Museologie (Ein Beitrag zur Geschichte der Entomologie). *Ent. Beihefte* 2: 1-160 + 12 pp., pls. 1-16 (14 December 1935), 3: 161-296, pls. 17-26 (4 October 1936), 4: 297-536, pls. 27-38, i-vi (1 August 1937)

[The title given above is that from the title page and of part III, the titles for parts I-II lacked the words "Entomologen & Entomo-Museologie."]

### **Horn, W., I. Kahle, G. Friese & R. Gaedike**

- 1990 *Collectiones entomologicae. Ein Kompendium über den Verbleib entomologischer Sammlungen der Welt bis 1960. Teil I: A bis K; Teil II: L bis Z.* 573 pp. Akademie der Landwirtschaftswissenschaften der Deutschen Demokratischen Republik, Berlin.

### **Horn, W. & S. Schenkling**

- 1928- *Index Litteraturae Entomologie. Serie I: Die Welt-Literatur über die gesamte Entomologie bis inklusive 1863.* Band I Aalborg - Ferrière,

- 1929 pp. 1-352, pl. 1 (May 1928); II Ferrill - Leconte, pp. 353-704, pl. 2 (August 1928); III Leconte - Schaum, pp. 705-1056, pl. 3 (November 1928); IV Schaum - Zwinger und Nachtrag, xxi + pp. 1057-1426, pl. 4 (February 1929). W. Horn, Berlin-Dahlem.

**Horning, W.**

- 1988 Entomology. Chap. 6, pp. 126-130. In Stanbury, P & J. Holland (eds.), *Mr Macleay's celebrated Cabinet. The history of the Macleays and their museum*. The Macleay Museum, University of Sydney, Sydney. (October)

**Hough, G. de N.**

- 1898 The Muscidae collected by Dr. A. Donaldson Smith in Somaliland. *Proc. Acad. nat. Sci. Philadelphia* 1898: 165-187. (3 May)

**Houttuyn, M.**

- 1761- *Natuurlyke historie of uitvoerige Beschryving der Dieren, Planten en Mineraalen, volgens het samenstel van Heer Linnaeus met nauwkeurige afbeeldingen*. Deel 1, Dieren (18 stukken). Houttuyn, Amsterdam.

[Vol. 12, [8+] 624 pp., pls. 93-98 (1768)]

**Huckett, H. C.**

- 1968 A note on the identity of *Hylemya variata* (Fallén) and *H. variabilis* Stein (Diptera: Anthomyiidae). *Proc. ent. Soc. Wash.* 70: 346-247. (18 December)

**Illiger, J. K. W.**

- 1807 Vol. 2, viii + 511 pp., 11 pls. In [Hellwig, J. C. L., & J. K. W. Illiger], *Favna Etrvsca sistens Insecta qvae in provinciis Florentina et Pisana praesertim collegit Petrus Rossivs. Iterum edita et annotatis perpetvis avcta*. 2 vols. Fleckeisen, Helmstadii [=Helmstadt]. (19 December)

**International Commission on Zoological Nomenclature**

- 1914 Opinion 57. Names dating from Hasselquist's "Iter Palaestinum," 1757, and the translation, 1762, are untenable. Pp. 131-134. In their Opinions rendered by the International Commission on Zoological Nomenclature, Opinions 57 to 65. *Smithson. Inst. Pub.* 2256: 131-169.

[Also published in Hemming, F., *Opinions and declarations rendered by the International Commission on Zoological Nomenclature*, Vol. 1, Sect. B, pp. 131-134, 1958.]

- 1925 Opinion 82. Suspension of rules for *Musca* Linnaeus, 1758A, type *Musca domestica*. Pp. 1-7. In their Opinions rendered by the International Commission on Zoological Nomenclature, Opinions 82 - 90. *Smithson. Inst.*, *Smithson. Misc. Collect.* 73(3), 40 pp. (16 December)

[Also published in Hemming, F., *Opinions and declarations rendered by the International Commission on Zoological Nomenclature*, Vol. 1, Sect. B, pp. 295-301, 1958.]

- 1954 Opinion 228. Rejection for nomenclatorial purposes of Geoffroy, 1762, *Histoire abrégée des Insectes qui se trouvent aux environs de Paris*. *Opin. Decl. Int. Comm. Zool. Nom.* 4: 209-220. (21 April)

- 1957 Opinion 441. Validation under the plenary powers of the names for five genera in the order Diptera (class Insecta) published in 1762 by Geoffroy (E. L.) in the work entitled *Histoire abrégée des Insectes qui se trouvent aux environs de Paris*. (Opinion supplementary to Opinion 228.) *Opin. Decl. int. Comm. zool. Nom.* 15: 83-120. (29 January)

- 1958 Opinion 516. Determination under the Plenary Powers of the relative precedence to be assigned to certain works on the Order Lepidoptera (Class Insecta) published in 1775 by Pieter Cramer, Michael Denis & Ignaz Schiffermüller, Johann Christian Fabricius, Johann Casper Fuessly, and S. A. von Rottemburg respectively. *Opin. Decl. int. Comm. zool. Nom.* 19: 1-44. (16 May)
- 1969 Opinion 884. *Musca lateralis* Linnaeus, 1758 (Insecta Diptera): suppressed under the plenary powers. *Bull. zool. Nom.* 26: 123-124. (24 October)
- 1987 Opinion 1444. *Musca trilineata* Linnaeus, 1767 (Currently *Oxycera trilineata*; Insecta, Diptera): Specific name conserved. *Bull. zool. Nom.* 44: 149. (25 June).
- 1988 Opinion 1507. *Musca marginalis* Wiedemann, 1830 (currently *Chrysomya marginalis*; Insecta, Diptera): specific name conserved. *Bull. zool. Nom.* 45: 236. (23 September)
- 1990 Opinion 1618. *Protocalliphora* Hough, 1899 (Insecta, Diptera) and its type species *Musca azurea* Fallén, 1817: Usage conserved by the designation of a replacement lectotype. *Bull. zool. Nom.* 47: 235-236. (28 September)
- 1992 Opinion 1670. *Calliphora vicina* Robineau-Desvoidy, 1830 (Insecta, Diptera): specific name conserved. *Bull. zool. Nom.* 49: 90. (26 March)
- Jackson, B. D.**
- 1890 History of the Linnean collections, prepared for the centenary anniversary of the Linnean Society. *Proc. Linn. Soc. Lond.* 1887-1888: 18-34.
- Joblot, L.**
- 1754 *Observations d'histoire naturelle, faites avec le microscope. Première partie.* 38 pp., 14 pls. Briasson, Paris.
- Johnson, C. W.**
- 1925 Diptera of the Harris collection. *Proc. Boston Soc. nat. Hist.* 38: 57-99. (November)
- Johnston, T.H. & M.J. Bancroft**
- 1920a The life histories of *Musca australis* Macq., and *M. vetustissima* Walker. *Proc. R. Soc. Queensl.* 31 [1919]: 181-203. (10 January)
- 1920b Notes on the chalcid parasites of muscoid flies in Australia. *Proc. R. Soc. Queensl.* 32: 19-30. (28 April)
- 1920c Experiments with certain Diptera as possible transmitters of bovine onchocerciasis. *Proc. R. Soc. Queensl.* 32: 31-57. (30 April)
- 1920d The life history of *Habronema* in relation to *Musca domestica* and native flies in Queensland. *Proc. R. Soc. Queensl.* 32: 61-88. (1 June)
- 1920e Notes on the biology of some Queensland flies. *Mem. Queensl. Mus.* 7: 31-43. (30 June)

**Jördens, J. H.**

- 1801- *Entomologie und Helminthologie des menschlichen Körpers, oder Beschreibung und Abbildung der Bewohner und Feinde desselben unter den Insekten und Würmern*. Vol. 1, 28 + 319 pp., 15 pls. (1801); 2, 2 + 153 pp., 7 pls (1802). G. A. Grau, Hof.

**Joseph, A. N. T. & P. Parui**

- 1972 A new subgenus of *Musca* from India (Diptera: Muscidae). *Zool. Anz.* 189: 179-181.

**Kabat, A. R. & R. E. Petit**

- 1988 The two printings of J. F. Gmelin's *Systema Naturae*, 13th Edition (1788-1796). *Nautilus* 102: 164-166.

**Karsholt, O. E. & E. Schmidt Nielsen**

- 1986 The Lepidoptera described by C. P. Thunberg. *Ent. Scand.* 16: 433-463. (10 March)

**Karsch, F. A.**

- 1886- Dipteren von Pungo-Andongo, gesammelt von Herrn Major Alexander von Homeyer. *Ent. Nachr.* 12: 49-58 (February), 257-264 (September), 337-342 (November); 13: 4-10 (January 1887)

**Kinel, J.**

- 1958 [After three years work in the Zoological Museum of the University of Wroclaw]. *Przeglad Zoologiczny* 1(4) [1957]: 305-312.  
[In Polish, with detailed English summary (pp. 310-312).]

**Kirby, W.**

- 1837 The Insects. Pt. 4, 325 pp., 8 pls. In Richardson, J., *Fauna Boreo-American, or the zoology of the northern parts of British America*. 4 pts. Fletcher, Norwich.

**Kleynhans, K. P. N.**

- 1987 *Musca nevilli* sp. nov. (Diptera: Muscidae), a dung-breeding fly from South Africa. *Onderstepoort J. vet. Res.* 54: 115-118. (June)

**Kowarz, F.**

- 1873 Beitrag zur Dipteren-Fauna Ungarns. *Verh. zool.-bot. Ges. Wien* 23: 453-464.

**Krastin, N. I.**

- 1950 Decipherment of the life-cycle of the nematode *Thelazia gulosa* (Railliet et Henry, 1910), a parasite of the eyes of cattle. [In Russian.] *Dokl. Akad. Nauk SSSR (n.s.)* 70: 549-551.

**Kuhn, T. S.**

- 1970 *The structure of scientific revolutions*. 2nd edition, enlarged. xii + 210 pp. Chicago University Press, Chicago.

**Kurahashi, H.**

- 1989 109. Family Calliphoridae. Pp. 702-718. In Evenhuis, N. L. (ed.), *Catalog of the Diptera of the Australasian and Oceanian Regions*. 1155 pp. Bishop Museum Press & E. J. Brill, Honolulu & Leiden. (23 August)

[Bishop Museum Special Publication 86.]

**Lamarck, J. B. P. A. de Monet**

- 1801 *Système des animaux sans vertèbres ou tableau général des classes, des ordres et des genres de ces animaux; présentant leurs caractères essentiels et leur distribution, d'après la considération de leurs rapports naturels et de leur organisation, et suivant l'arrangement établi dans les galeries du Muséum d'Hist. Naturelle, parmi leurs dépouilles conservées; précédé du discours d'ouverture du cours de Zoologie, donné dans le Muséum National d'Histoire Naturelle l'an 8 de la République.* viii + 432 pp. Déterville, Paris ("an IX"). (January)
- 1816 *Histoire naturelle des animaux sans vertèbres, présentant les caractères généraux et particuliers de ces animaux, leur distribution, leurs classes, leurs familles, leurs genres, et la citation des principales espèces qui s'y rapportent; précédée d'une introduction offrant la détermination des caractères essentiels de l'animal, sa distinction du végétal et des autres corps naturels, enfin, l'exposition des principes fondamentaux de la zoologie.* Tome troisième. 586 pp. Verdière, Paris. (August)
- Latreille, P. A.**
- 1796 *Précis des caractères génériques des insectes, disposés dans un ordre naturel.* xiv + 201 [+ 7] pp., 1 folding chart. Prévôt, Paris, & Brive, Bourdeaux, "an 5 de la R." (after 22 September)  
[Also, a facsimile edition by A. Hermann, 1947 [Paris?].]
- 1802 *Histoire naturelle, générale et particulière, des Crustacés et des Insectes.* Tome troisième. Familles naturelles des genres. [Vol. 95], 468 pp. In Sonnini, C.S. (ed.), *Histoire naturelle par Buffon, nouvelle édition accompagnée de notes.* Dufart, Paris, "An X". (6 November)  
[The year on the title page is incorrect. The volume was actually published in "an XI" of the Republic, which was 22 September 1802 until 21 September 1803 (Griffin 1938; also Dupuis 1986).]
- 1809 *Genera crustaceorum et insectorum secundum ordinem naturalem in familias disposita, iconibus exemplisque plurimis explicata.* Volume 4. 399 pp. Koenig, Parisiis et Argentorati [= Paris & Strasbourg]. (20 March)
- 1810 *Considérations générales sur l'ordre naturel des animaux composant les classes des Crustacés, des Arachnides, et des Insectes; avec un tableau méthodique de leurs genres, disposés en familles.* 444 pp. Schoell, Paris. (23 May)
- 1818 Mouche, *Musca.* Pp. 437-442. In Société de Naturalistes et d'Agriculteurs. *Nouveau Dictionnaire d'Histoire Naturelle, appliquée aux arts, à l'agriculture, à l'économie rurale et domestique, à la médecine, etc.* Nouvelle édition, presqu'entièrement refondue et considérablement augmentée; avec des figures tirées des trois règnes de la nature. Volume 21. 612 pp., plates M18, G26-29. Déterville, Paris. (5 September)

1825 *Familles naturelles du règne animal, exposées succinctement et dans un ordre analytique, avec l'indication de leurs genres.* 570 pp. Baillière, Paris. (16 May)

1829 Suite et fin des insectes. Tome 5. xxiv + 556 pp. In Cuvier, G. L. C. F. D., *Le règne animal distribué d'après son organisation, pour servir de base à l'histoire naturelle des animaux et d'introduction à l'anatomie comparée. Nouvelle édition, revue et augmentée.* Déterville & Crochard, Paris. (11 April)

**Leeuwenhoek, A. van**

1719 *Epistolae physiologicae super compluribus naturae arcanis, ubi variorum animalium atque plantarum fabrica, conformatio, proprietates atque operationes, novis & hactenus inobservatis experimentis illustrantur & oculis exhibentur; item peculiares & hactenus incognitae rerum quarundam qualitates explicantur; ut sequens pagina docet; hactenus numquam editae. Cum figuris aeneis, & indice locupletissimo.* [20+] 446 [+ 26] pp., many plates, Beman, Delphis [=Delft].

**Le Guillou, E. J. F.**

1842 Description de sept Diptères nouveaux, recueillis pendant le voyage autour du monde de l'Astrolabe et la Zélée. *Revue zool. Soc. Cuvier.* 5: 314-316. (October)

**Lepechin, I. I.**

1774 *Tagebuch der Reise durch verschiedene Provinzen des Russischen Reiches in den Jahren 1768 und 1769. Aus dem Russischen übersetzt von M. Christian Heinrich Hase.* [viii] + 332 pp., 23 pls. Richter, Altenburg. (April)

1775 *Tagebuch der Reise durch verschiedene Provinzen des Russischen Reiches im Jahr 1770. Aus dem Russischen übersetzt von M. Christian Heinrich Hase.* [ii] + 211 pp., 11 pls. Richter, Altenburg.

1783 *Tagebuch der Reise durch verschiedene Provinzen des Russischen Reiches im Jahr 1771. Aus dem Russischen übersetzt von M. Christian Heinrich Hase.* [ii] + 234 pp., 17 pls. Richter, Altenburg.

**Lichtenstein, A. A. H.**

1796 *Catalogus musei zoologici ditissimi Hamburgi, d III. Februar 1796. Auctionis lege distrahendi. Sectio Tertia. Continens Insecta.* 223 pp. Hamburg. (3 February)

[For full title and particulars, see Burger & Thompson 1981: 356.]

**Lindroth, C. H.**

1957 The Linnaean species of carabid beetles. *Zool. J. Linn. Soc.* 43: 325-341. (February)

1973 Systematics specializes between Fabricius and Darwin; 1800-1859. Pp. 119-154. In Smith, R. F., T. E. Mittler & C. N. Smith [eds], *History of Entomology.* vii + 517 pp. Annual Reviews Inc., Palo Alto, California.

**Linnaeus, C.**

- 1746 *Fauna Svecica sistens Animalia Sveciae Regni: Quadrupedia, Aves, Amphibia, Pisces, Insecta, Vermes, distributa per classes & ordines, genera & species. Cum differentiis specierum, synonymis auctorum, nominibus incolarum, locis habitationum, descriptionibus Insectorum.* [xxviii] + 411 pp., 2 pls. L. Salvii, Stockholmiae [=Stockholm]. (20 October)
- 1758a *Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum caracteribus, differentiis, synonymis, locis.* Ed. 10, Vol. 1. 824 pp. L. Salvii, Holmia [=Stockholm]. (1 January)
- 1758b *Pandora insectorum, quam sub Praesido D. D. Car. Linnaei, proposuit Ericus Ol. Rydbeck, Ostro-Gothus. Upsaliae 1758. Julii 15.* [i] + 31 pp., 1 pl. (15 July)  
[Also reprinted in his *Amoenitates Academicae; seu Dissertationes variae Physicae, Medicae, Botanicae antehac seorsim editae, nunc collectae et auctae, cum tabulis aeneis.* [Ed. 1] 4: 232-252. Laurentii Salvii, Holmia [=Stockholm] 1760.]
- 1761 *Fauna Svecica sistens Animalia Sveciae Regni: Mammalia, Aves, Amphibia, Pisces, Insecta, Vermes. Distributa per classes & ordines, genera & species, cum differentiis specierum, synonymis auctorum, nominibus incolarum, locis natalium, descriptionibus insectorum.* Editio altera, auctior [49] + 578 pp., 2 pls. L. Salvii, Stockholmiae [=Stockholm]. (after 28 July, date of preface)
- 1762 Zweyter Theil, enthält Beschreibungen verschiedener wichtiger Naturalien. Pp. 267-606. In Hasselquist, F., *Reise nach Palästina in den Jahren von 1749 bis 1752.* viii + 606 pp. J. C. Koppe, Rostock. (July)  
[This is the German translation by T. H. Gadebusch of Hasselquist, F., *Iter Palaestinum, eller resa til heliga landet, foerrättad ifran ar 1749 til 1752 med beskrifningar, rön annerkningar äfver maerkwaerdigste naturalier, utgjifven af C. Linnaeus.* 619 pp. Holmia [=Stockholm] 1757. Suppressed by I. C. Z. N. 1914: 131.]
- 1764a XLIII. *Hospita insectorum flora, quam praeside D:n. Doct. Carolo Linnaeo, proposuit alumnus degeerianus Jonas Gustav Forsskål Uplandus.* Upsaliae 1752. Novemb. 4. Vol. 3: 171-312 in his *Amoenitates Academicae; seu Dissertationes variae Physicae, Medicae, Botanicae antehac seorsim editae, nunc collectae et auctae, cum tabulis aeneis.* [Ed. 1, reprint]. Laurentii Salvii, Holmia [=Stockholm].  
[While the author of this dissertation was J. G. Forsskål the names are attributed to Linnaeus.]
- 1764b XLIV. D. D. *Miracula insectorum, sub Praesido D:ni Doct. Caroli Linnaei, s:ae r:ae M:tis archiatri, proposuit Gabriel Emanuel Avelin, Dalekarlus,* Upsaliae 1752. Novemb. 19. Vol. 3: 313-334 in his *Amoenitates Academicae; seu Dissertationes variae Physicae, Medicae, Botanicae antehac seorsim editae, nunc collectae et auctae, cum tabulis aeneis.* [Ed. 1, reprint]. Laurentii Salvii, Holmia [=Stockholm].
- 1764c XLV. D. D. *Noxa insectorum sub Praesidio D:n Doct. Caroli Linnaei, proposita a Michael A. Baeckner, Helsingo,* Upsaliae, 1752, Decemb. 18. Vol. 3: 335-362 in his *Amoenitates Academicae; seu Dissertationes variae Physicae, Medicae, Botanicae antehac seorsim editae, nunc collectae et auctae, cum tabulis aeneis.* [Ed. 1, reprint]. Laurentii Salvii, Holmia [=Stockholm].

*nes variae Physicae, Medicae, Botanicae antehac seorsim editae, nunc collectae et auctae, cum tabulis aeneis.* [Ed. 1, reprint]. Laurentii Salvii, Holmiae [=Stockholm].

- 1767 *Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum caracteribus, differentiis, synonymis, locis.* Ed. 12 (revised.). Vol. 1, Pt. 2, pp. 533-1327. L. Salvii, Holmiae [=Stockholm]. (14 June)

- 1771 *Pandora et flora Rybyensis quam praeside D. D. Car. v. Linné proposuit Daniel Henr. Söderberg Junecopia Smolandus. Upsaliae 1771. Junii 26.*

[This dissertation was reprinted in his *Amoenitates Academicae seu Dissertationes variae Physicae, Medicae Botanicae antehac seorsim editae nunc collectae et auctae cum tabulis aeneis*, Ed. 2 [Schreber], 8: 75-106, Erlangae [=Erlangen], 1785. While the author of this dissertation was D. H. Söderberg the names are attributed to Linnaeus.]

### Lioy, P.

- 1864 I Ditteri distribuiti secondo un nuovo metodo di classificazione naturale. [Part.] *Atti R. Ist. veneto Sci.* (3) 9: 879-910. (20 April)

### Lisney, A. A.

- 1960 *A bibliography of British Lepidoptera 1608-1799.* xviii + 315 pp. The Chiswick Press, London.

### Lister, M.

- 1682 *Johannes Godartius of Insects. Done into English, and methodized, with the addition of notes.* 4 + 140 pp. John White, York.

### Loew, H.

- 1856 Neue Beiträge zur Kenntniss der Dipteren. Vierter Beitrag. *Programm K. Realschule Meseritz* 1856: 1-57. (17 March)

[Also issued as a separate with same title and pagination under the imprint "Mittler, Berlin, 1856".]

### Lopes, H. de Souza

- 1975 On female holotypes of some American species described by Francis Walker and J. Macquart (Diptera, Sarcophagidae, Calliphoridae). *Revta bras. Biol.* 34: 535-549. (20 June)

### Lundbeck, W.

- 1916 *Diptera Danica. Genera and species of flies hitherto found in Denmark.* Part V. Lonchopteridae, Syrphidae. 603 pp., G. E. C. Gad, Copenhagen. (1 July)

### Macquart, J.

- 1834 Insectes Diptères du nord de la France. Athéricères: Créophiles, Oestrides, Myopaires, Conopsaires, Scénopiniens, Céphalopsides. *Mém. Soc. R. Sci. Agric. Arts, Lille* 1833: 137-368, 6 pls. (before 1 October)

[Also published separately as his "Insectes Diptères du nord de la France, vol. 5, 232 pp., 6 pls., Lille, 1834.]

- 1835 Histoire naturelle des Insectes.--Diptères. Vol. 2. 703 pp., pls. 13-24. In [Roret, N. E., ed.], *Nouvelles suites à Buffon, formant, avec les œuvres de cet auteur, un cours complet d'Histoire Naturelle. Collection accompagnée de planches.* Roret, Paris. (before 1 April)

- 1843 Diptères exotiques nouveaux ou peu connus. Vol. 2, 3rd part. *Mém. Soc. R. Sci. Agric. Arts, Lille* 1842: 162-460, 36 pls.  
 [Published separately with pagination, 5-304, 36 pls. Roret, Paris, 1843. There is the possibility that the journal version was actually published in 1844; Guérin-Méneville reviews it in the February 1844 issue of *Rev. Zool. Soc. Cuvier.; Bibliogr. Fr.* mentions it in the 22 June 1844 issue; Macquart presented it to the Société Entomologique de la France at the 3 April 1844 meeting.]
- 1846 Diptères exotiques nouveaux ou peu connus. Supplément. *Mém. Soc. R. Sci. Agric. Arts, Lille* 1844: 133-364, 20 pls. (before 22 July)  
 [The date on the wrapper of the journal is 1846, the date on the title page is 1845. The last 4 pages of this paper in the journal (containing the index) are numbered 363, 364, 363, 364. The next article in the journal begins on page 365. The pagination of the reprint is continuous, thus the reprint appears to contain 2 additional pages. Reprinted with pagination 5-238, 20 pls. Roret, Paris, with "1846" on both the wrapper and title page. Reprint recorded in the 7 November 1846 issue of *Bibliogr. Fr.* The *Bibliogr. Fr.* mentions the journal version in the 22 August 1846 issue; it was presented by Macquart at the 22 July 1846 meeting of the Société Entomologique de la France.]
- 1847a *Diptères exotiques nouveaux ou peu connus*. 2<sup>e</sup> supplément. Roret, Paris. Pp. 5-104, 6 pls. (before 11 September)  
 [This is the reprint version issued in advance of the journal version, Macquart, 1847b. Another version of this reprint is bound with Macquart, 1848. Its wrapper has "2.<sup>e</sup> et 3.<sup>e</sup> supplements." and the dates "1847 et 1848."] [Mentioned in the 25 December 1847 issue of *Bibliogr. Fr.*]
- 1847b Diptères exotiques nouveaux ou peu connus. 2<sup>e</sup> supplément. *Mém. Soc. R. Sci. Agric. Arts, Lille* 1846: 21-120, 6 pls. (before 25 December)  
 [Presented by Macquart at the 22 November meeting of the Société Entomologique de la France. Reprint version with separate pagination 1-77, 7 pls. Roret, Paris. See also annotation under Macquart, 1847a.]
- 1850a Diptères exotiques nouveaux ou peu connus. 4<sup>e</sup> supplément [part]. *Mém. Soc. R. Sci. Agric. Arts, Lille* 1849: 309-479, pl. 1-14. (before 6 July)  
 [Reprinted with pagination 5-161, 14 pls. Roret, Paris. This includes only p. 309-465 of the original journal article. The legends to the plates and the index (p. 466-79 of the journal version) are not found in all copies of the reprint. A reprinted version seen by us has the dates "1850 et 1851" on the wrapper and contains the plate legends and index on p. 337-64. See also annotation under Macquart, 1851.]
- 1850b Animaux invertébrés. Pp. 1-634 pp. In *Catalogue du Musée d'Histoire naturelle de la Ville de Lille*. Vol. 2. iv + 634 pp. L. Danel, Lille.
- 1851 Diptères exotiques nouveaux ou peu connus. Suite du 4<sup>e</sup> supplément publié dans les mémoires de 1849. *Mém. Soc. R. Sci. Agric. Arts, Lille* 1850: 134-294, pl. 15-28. (before 5 April)  
 [Reprinted by Roret, Paris (without title, continuing from the end of the 1850 reprint without a break in the text or arrangement) with separate pagination 161-336, pls. 15-28. The plate legends and index to both the 1850 and 1851 reprinted versions is present on p. 337-64 in a copy seen by us.]
- 1855 Diptères exotiques nouveaux ou peu connus. 5<sup>e</sup> supplément. *Mém. Soc. Sci. Agric. Lille* (2) 1: 25-156, 7 pls. (before 18 August)  
 [Recorded in the 18 August issue of *Bibliogr. Fr.* Reprinted with separate pagination, 5-136, 7 pls. Roret, Paris, 1855.]
- Magpayo, F. R., S. Shinonaga & R. Kano**
- 1987 Studies on the calypterate muscoid flies in Philippines. 1. Report on species belonging to the genus *Musca* Linnaeus (Diptera, Muscidae). *Jap. J. sanit. Zool.* 38: 257-269. (15 December)

**Malloch, J. R.**

- 1924 Exotic Muscaridae (Diptera). XV. *Ann. Mag. nat. Hist.* (9) 15 [1925]: 131-142. (31 December; date received at BMNH)
- 1925 Exotic Muscaridae (Diptera). XVII. *Ann. Mag. nat. Hist.* (9) 16: 361-377. (1 October; date received at BMNH)
- 1928 Fauna sumatrensis (Beitrag No. 56). Family Muscidae (Dipt.) *Entomol. Mitt.* 17: 290-303 (5 July), 310-336 (20 September)
- 1929a Exotic Muscaridae (Diptera). XXIV. *Ann. Mag. nat. Hist.* (10) 3: 249-280 (March)
- 1929b Exotic Muscaridae (Diptera). XXVI. *Ann. Mag. nat. Hist.* (10) 4: 97-120. (July)
- 1929c Exotic Muscaridae (Diptera). XXVII. *Ann. Mag. nat. Hist.* (10) 4: 249-257. (September)
- 1932 Exotic Muscaridae (Diptera). XXXVII. *Ann. Mag. nat. Hist.* (10) 10: 297-330. (October)
- 1934 Diptera II. Cyclorrhapha: Muscidae, Calliphoridae and Tachinidae. In van Straelen, V. (ed.), *Résultats scientifiques du voyage aux Indes Orientales Néerlandaises de LL. AA. RR. le Prince et la Princesse Léopold de Belgique. Mém. Mus. r. Hist. nat. Belg.* (hors série) 4 (10): 1-24. (30 June)

**Manuel, B. E.**

- 1797 Mouche. [Part.] Pp. 741-827. In Olivier, G. A., *Histoire Naturelle des Insectes*, Volume 7, Han-Mou. 827 pp. In Société de Gens de Lettres, de Savans et d'Artistes, *Encyclopédie Méthodique*. Panckoucke, Paris.  
[The volume is attributed to Olivier, but the article on "Mouche" is signed on p. 827 by Manuel.]
- 1811 Mouche. [Concl.] Pp. 1-45. In Olivier, G. A., *Histoire Naturelle des Insectes*, Volume 8, Mou-Pao. 722 pp. In Société de Gens de Lettres, de Savans et d'Artistes, *Encyclopédie Méthodique*. Agasse, Paris.  
[A footnote on p. 45, after "Mouche", states that Olivier was returning to the editorship of this volume after an absence of several years on Government service, during which time the work from the letter L onwards had been entrusted to several collaborators. Manuel's name is signed at the end of the first part of the article on "Mouche" (see above). The rest of this article is likewise attributed to Manuel as the footnote appears after the article. Also, see Poche 1933.]

**Mayer, J.**

- 1779 Bemerkungen über natürliche Gegenstände der Gegend um Schüttenhofen in Böhmen, und eines Theils der benachbarten Gebirge. *Abh. Privat Ges. Bohmen., Prag* 4: 132-184.

**Mayr, E.**

- 1969 *Principles of Systematic Zoology*. xi + 428 pp. New York, St. Louis, San Francisco.

**Meigen, J. W.**

- 1803 Versuch einer neuen GattungsEintheilung der europäischen zweiflüglichen Insekten. *Illiger's Mag. Ins.* 2: 259-281.

- 1804 *Klassifikation und Beschreibung der europäischen zweiflügeligen Insekten. (Diptera Linn.)* Erster Band. xxviii + 152 pp., 8 pls. Reichard, Braunschweig. (5 November)
- 1818 *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten.* Erster Theil. xxxvi + 333 pp., 11 pls. Forstmann, Aachen. (before October)  
 [Listed as received in the October issue of *Isis (Oken's.)*.]
- 1826 *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten.* Fünfter Theil. xii + 412 pp., pls. 42-54. Schultz, Hamm. (before November)
- 1830 *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten.* Sechster Theil, xi + 401 pp., pls. 55-66. Schulz, Hamm.
- 1835 [new species in Gistl, J. N. F. X. (q.v.)]
- 1838 *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten.* Siebenter Theil oder Supplementband, xii + 434 pp., pls. 67-74. Schulz, Hamm.
- 1975 [*Nomina nuda in Morge.*] Meijere —see de Meijere
- Meuschen, F. C.**
- 1787 *Museum sive index rerum naturalium continens instructissimam copiam pretiosissimorum omnis generis ex tribus regnis naturae objectorum quam dum in vivis erat magna diligentia multaque cura comparavit vir amplissimus Abrahamus Gevers olim consiliarius primusque urbis Rotterodamensis consul praefectus sylvarum Hollandiae & Westrisiae Societatis Indiae orientalis director academiae caesareae naturae curiosorum socius etc. etc. iv + 659 pp.* P & L Hosteyn, Rotterodami [=Rotterdam]. [before 12 September]  
 [This is a sale catalog, the sale being held on 12 September. The work was also reviewed in *Göt. Gel. Anz.* on 22 September (Evenhuis in litt.).]
- Michelsen, V.**
- 1977 On the identity of *Musca vagans* Panzer, 1798 (Diptera: Anthomyiidae and Muscidae). *Ent. Scand.* 8: 320. (15 December)
- 1979a Notes on the identity and type-material of the Anthomyiidae, Fanniidae, and Muscidae (Diptera) described by J. C. Fabricius. *Steenstrupia* 5: 181-196. (20 September)
- 1979b Identity of *Musca conformis* Fallén, 1825, with a new synonymy in the genus *Pegomya* Robineau-Desvoidy, 1830 (Diptera: Anthomyiidae). *Ent. Scand.* 10: 257-258. (15 December)
- 1983 The Anthomyiidae (Diptera) described by Fallén, with a review of European *Emmesomyia* Malloch. *Ent. Scand.* 14: 109-120. (15 April)
- 1985 A revision of the Anthomyiidae (Diptera) described by J. W. Zetterstedt. *Steenstrupia* 11: 37-65. (5 March)
- Mikkola, K.**
- 1983 Diagnostic insect pins: some problems of the Linnean insect collection solved. *Antenna* 7: 16-17.

**Morge, G.**

- 1973 Entomology in the Western World in antiquity and in medieval times. Pp. 37-80. In Smith, R. F., T. E. Mittler & C. N. Smith [eds]. *History of Entomology*. vii + 517 pp. Annual Reviews Inc., Palo Alto, California.
- 1975 Dipteren-Farbtafeln nach den bisher nicht veröffentlichten Original-Handzeichnungen Meigens: "Johann Wilhelm Meigen: Abbildung der europaeischen zweiflügeligen Insecten, nach der Natur". Pars I. Vorwort des Herausgebers - Faksimiles von Original-Legenden und -Index: "Erklaerung der Fig." und "Alphabetisches Verzeichniss der Gattungen und Arten" - Farbtafeln I-LXXX. *Beitr. Ent.* 25: 383-500, pls 1-80.
- 1976a Dipteren-Farbtafeln nach den bisher nicht veröffentlichten Original-Handzeichnungen Meigens: "Johann Wilhelm Meigen: Abbildung der europaeischen zweiflügeligen Insecten, nach der Natur". Pars II: Farbtafeln LXXXI-CLX. *Beitr. Ent.* 26: 441, pls 81-160.
- 1976b Dipteren-Farbtafeln nach den bisher nicht veröffentlichten Original-Handzeichnungen Meigens: "Johann Wilhelm Meigen: Abbildung der europaeischen zweiflügeligen Insecten, nach der Natur". Pars III: Farbtafeln CLXI-CCCV. *Beitr. Ent.* 26: 543, pls 161-305.

**Morris, F. O.**

- 1865 *A catalogue British Insects, in all the Orders*. 125 pp. Longmans, Green, Reader & Dyer, London.

**Moufet, T.**

- 1634 *Insectorvm sive minimorum animalium theatrvm: olim ab Edoardo Wottono. Conrado Gesnero. Thomaqve Pennio inchoatum: tandem Tho. Movfeti Londinatis operâ sumptibusq; maximis concinnatum, auctum, perfectum: et ad vivum expressis iconibus supra quingentis illustratum*. [18+] 326 pp. Cotes, Londoni [=London].

**Müller, O. F.**

- 1764 *Favna insectorum Fridrichsdalina, sive methodica descriptio insectorum agri Fridrichsdalensis, cum characteribus genericis et specificis, nominibus trivialibus, locis natalibus, iconibus allegatis, novisq; pluribus speciebus additis*. xxiv + 96 pp. Io. Frid. Gleditschii, Hafniae et Lipsiae [=Copenhagen and Leipzig]. (26 March)
- 1766 [New species in Allioni 1766 (q.v.)]  
[See Bezzu (1908a: 74) for details on authorship.]
- 1776 *Zoologiae Danicae prodromus, seu Animalium Daniae et Norvegiae indigenarum characteres, nomina, et synonyma imprimis popularium*. xxxii + 274 [+8] pp. Hallageriis, Havniae [=Copenhagen]. (20 May)  
[Anker (1950: 28) gave 20 May for publication date. The book was also listed for sale at the Leipzig Easter Book Fair, which would mean a publication date before 21 April.]

**Müller, P. L. S.**

- 1775 *Des Ritters Carl von Linné Königlich Schwedischen Leibarztes &c. &c. vollständiges Natursystem nach der zwölften lateinischen Ausgabe und nach Anleitung des holländischen Houttuynischen Werks mit einer ausführlichen Erklärung ausgefertiget.* Vol. 5, pt. 2, pp. 761-1066 + register (114 unnumbered pages), pls. 23-26. Gabriel Nicolaus Raspe, Nürnberg.

**Nash, R. & P. J. Chandler**

- 1978 The Irish species of two-winged flies (Diptera) belonging to the families Fanniidae and Muscidae. *Proc. R. Ir. Acad. (B)* 78: 13-43.

**Neboiss, A.**

- 1963 The Trichoptera types of species described by J. Curtis. *Beitr. Ent.* 13: 582-635.

**Osten Sacken, C. R.**

- 1881a A brief notice of Carl Ludwig Doleschall, the dipterologist. *Ent. mon. Mag.* 18: 114-116. (October)

- 1881b An essay of comparative chaetotaxy, or the arrangement of characteristic bristles of Diptera. *Mitt. münchen. ent. Ver.* 5: 121-138.

- 1884 An essay of comparative chaetotaxy, or the arrangement of characteristic bristles of Diptera. *Trans. ent. Soc. London* 1884: 497-517. (3 December)

- 1887 On Mr. Portchinski's publications on the larvae of Muscidae, including a detailed abstract of his last paper: Comparative biology of the necrophagous and coprophagous larvae. *Berl. ent. Z.* 31: 17-28. (August)

**Ôuchi, Y.**

- 1938a Diptera Sinica. Muscidae-Muscinae. I. On some muscid flies from eastern China. (Entomological Report No. 5). *J. Shanghai Sci. Inst.* (3) 4: 1-14, pl. 1. (February)

- 1938b Diptera Sinica. Muscidae-Muscinae. II. A new murcid [sic] fly from southern China. (Entomological Report No. 10). *J. Shanghai Sci. Inst.* (3) 4: 31-32. (February)

**Pamplona, D.**

- 1986 Revalidação de *Biopyrellia* Townsend, 1932 com redescrição de *B. bipuncta* (Wiedemann, 1830) (Diptera, Muscidae, Muscinae). *Revta bras. Biol.* 46: 49-54. (February)

- 1987 Revalidação de *Morellia nitida* Wiedemann, 1830 (Diptera, Muscidae, Muscinae). *Revta bras. Biol.* 47: 217-218. (30 May)

**Pandellé, L.**

- 1894 Études sur les Muscides de France. II<sup>e</sup> partie. *Revue Ent.* 13: 1-113.

- 1898a Études sur les Muscides de France. III<sup>e</sup>-partie. [Part.] *Revue Ent.* 17 (7-8) (Suppl.): 1-32. (?August)

- 1898b Études sur les Muscides de France. III<sup>e</sup>-partie. [Part.] *Revue Ent.* 17 (9-10) (Suppl.): 33-64. (?October)

**Panzer, G. W. F.**

- 1794a *Favnae insectorum germanicae initia oder Deutschlands Insecten*. H. 17, 2 pp., 24 pls. Felsecker, Nürnberg. (January)
- 1794b *Favnae insectorum germanicae initia oder Deutschlands Insecten*. H. 20, 2 pp., 24 pls. Felsecker, Nürnberg. (April)
- 1794c *Favnae insectorum germanicae initia oder Deutschlands Insecten*. H. 24, 2 pp., 24 pls. Felsecker, Nürnberg. (after 21 August)
- 1798a *Favnae insectorum germanicae initia oder Deutschlands Insecten*. H. 54, 2 pp., 24 pls. Felsecker, Nürnberg.
- 1798b *Favnae insectorum germanicae initia oder Deutschlands Insecten*. H. 59, 2 pp., 24 pls. Felsecker, Nürnberg.
- 1798c *Favnae insectorum germanicae initia oder Deutschlands Insecten*. H. 60, 2 pp., 24 pls. Felsecker, Nürnberg. (after 1 May)
- 1800 *Favnae insectorum germanicae initia oder Deutschlands Insecten*. H. 73, 2 pp., 24 pls. Felsecker, Nürnberg.
- 1801 *Favnae insectorum germanicae initia oder Deutschlands Insecten*. H. 82, 2 pp., 24 pls. Felsecker, Nürnberg.
- 1804 D. Jacobi Christiani Schaefferi Iconum Insectorum circa Ratisbonam indigenorum Enumeratio Systematica. xvi + 260 pp. Palmii, Erlangae [=Erlangen].
- 1805 *Favnae insectorum germanicae initia oder Deutschlands Insecten*. H. 97, 2 pp., 24 pls. Felsecker, Nürnberg.
- 1806 *Favnae insectorum germanicae initia oder Deutschlands Insecten*. H. 104, 2 pp., 24 pls. Felsecker, Nürnberg.

**Pape, T.**

- 1986 A revision of the Sarcophagidae (Diptera) described by J. C. Fabricius, C. F. Fallén, and J. W. Zetterstedt. *Ent. Scand.* 17: 301-312. (30 December)
- 1988 A revision of the Palaearctic Sarcophagidae (Diptera) described by C. Rondani. *Stuttgart. Beitr. Naturk.* (A) 416, 22 pp. (1 September)

**Paterson, H. E.**

- 1956 East-African Muscidae (Diptera). (*Ergebnisse der Deutschen Zoologischen Ostafrika-Expedition 1951/1952*, Gruppe Lindner, Stuttgart, Nr. 20). *Beitr. Ent.* 6: 154-179. (April)
- 1957 Three new species of *Musca* from southern Africa (Dipt., Muscidae). *J. entomol. Soc. sth. Africa* 20: 106-113. (31 May)

**Patton, W. S.**

- 1920 Some notes on the arthropods of medical and veterinary importance in Mesopotamia, and on their relation to disease. Part II. Mesopotamian house flies and their allies. *Indian J. Med. Res.* 7: 751-777, pls. 68-71. (April)
- 1922a Notes on the species of the genus *Musca* Linnaeus - Part I. *Bull. ent. Res.* 12: 411-426. (February)

- 1922b New Indian species of the genus *Musca*. *Indian J. Med. Res.* 10: 69-77. (July)
- 1923a Some Philippine species of the genus *Musca* Linnaeus. *Philipp. J. Sci.* 23: 309-322. (October)
- 1923b A new Oriental species of the genus *Musca*, with a note on the occurrence of *Musca dasyops* Stein in China and a revised list of the Oriental species of the genus *Musca* Linnaeus. *Philipp. J. Sci.* 23: 323-335. (October)
- 1925 Diptera of medical and veterinary importance, I. Types of older authors in continental museums. *Philipp. J. Sci.* 27: 177-200. (13 June)
- 1926 The Ethiopian species of the genus *Musca* Linnaeus. *Rec. Indian Mus.* 28: 29-52, pls 1-4. (31 March)
- 1932 Studies on the higher Diptera of medical and veterinary importance. A revision of the species of the genus *Musca* based on a comparative study of the male terminalia. I. The natural grouping of the species and their relationship to each other. *Ann. Trop. Med. Parasit.* 26: 347-405. (29 October)
- 1933a Studies on the higher Diptera of medical and veterinary importance. A revision of the genera of the tribe *Muscini*, subfamily *Muscinae*, based on a comparative study of the male terminalia. 1. The genus *Musca* Linnaeus. *Ann. trop. Med. Parasit.* 27: 135-156. (10 April)
- 1933b Studies on the higher Diptera of medical and veterinary importance. A revision of the species of the genus *Musca*, based on a comparative study of the male terminalia. II. A practical guide to the Palaearctic species. [Part.] *Ann. trop. Med. Parasit.* 27: 327-345. (7 July)
- 1933c Studies on the higher Diptera of medical and veterinary importance. A revision of the species of the genus *Musca*, based on a comparative study of the male terminalia. II. A practical guide to the Palaearctic species. [Concl.] *Ann. trop. Med. Parasit.* 27: 397-430. (21 October)
- 1933d A new haematophagous species of *Musca* from Malaya. *Ann. trop. Med. Parasit.* 27: 477-480. (21 October)
- 1934 The blood-sucking species of the genus *Musca* and the evolution of the blood-drawing proboscis in the genus. *Proc. Pacif. Sci. Congr.* 1933 5: 3361-3366.
- 1936 Studies on the higher Diptera of medical and veterinary importance. A revision of the species of the genus *Musca*, based on a comparative study of the male terminalia. III. A practical guide to the Ethiopian species. *Ann. Trop. Med. Parasit.* 30: 469-490. (23 December)
- 1937a Studies on the higher Diptera of medical and veterinary importance. A revision of the species of the genus *Musca*, based on a comparative study of the male terminalia. IV. A practical guide to the Oriental species. [Part.] *Ann. trop. Med. Parasit.* 31: 127-140. (8 April)
- 1937b Studies on the higher Diptera of medical and veterinary importance. A revision of the species of the genus *Musca*, based on a comparative

study of the male terminalia. IV. A practical guide to the Oriental species. [Concl.] *Ann. trop. Med. Parasit.* 31: 195-213. (13 July)

**Patton, W. S. & F. W. Cragg**

- 1912 The genus *Pristirhychomyia* Brunetti, 1910. *Ann. trop. Med. Parasit.* 5: 509-514, pl. 25. (26 February)

- 1913a On certain haematophagous species of the genus *Musca*, with descriptions of two new species. *Indian J. Med. Res.* 1: 11-25, pls. 1-5. (July)

- 1913b A new species of *Philaematomyia*, with some remarks on the genus. *Indian J. Med. Res.* 1: 26-33, pls. 6-8. (July)

**Patton, W. S. & E. MacGill.**

- 1925 Comparative studies of the antennae of some of the higher Diptera. *Indian J. med. Res.* 13: 275-285, pls 7-13. (October)

**Patton, W. S. & R. Senior-White.**

- 1924 The Oriental species of the genus *Musca* Linnaeus. *Rec. Indian Mus.* 26: 553-577, pls. 29-33. (29 November)

**Percheron, A.**

- 1837 *Bibliographie Entomologique, comprenant l'indication par ordre alphabétique de noms d'Auteurs. I° des ouvrages entomologiques publiés en France et à l'Étranger, depuis les temps les plus réculés jusques et y compris l'Année 1834; 2° des monographies et mémoires contenus dans les recueils, journaux et collections académiques Françaises et Étrangères; accompagnée de notices sur les ouvrages périodiques, les Dictionnaires et les Mémoires des Sociétés savantes; suivie d'une table méthodique et chronologique des matières.* Vol. 1, xii + 326 pp.; vol. 2, 376 pp. J. B. Baillièvre, Paris & London.

**Persson, P. I.**

- 1964 Entomologiska museet och entomologer i Lund fram till 1800-talets slut. *Opusc. ent.* 29: 57-80. (31 March)

- 1972 "Eugenies resa." Localities, dates and labels of the insects collected during the voyage around the world by the Swedish frigate "Eugenie" in the years 1851-1853. *Ent. Tidskr.* 92 [1971]: 164-172.

**Persson, P. I., A. C. Pont & V. Michelsen**

- 1984 Notes on the insect collection of Charles De Geer, with a revision of his species of Fanniidae, Anthomyiidae and Muscidae (Diptera). *Ent. scand.* 15: 89-95. (7 May)

**Petagna, V.**

- 1792 *Institutiones entomologicae.* 12 + 718 + 10 pp., 10 pls. C. Raymundi, Neapoli [=Naples].

**Peus, F.**

- 1960 Zur Kenntnis der ornithoparasitischen Phormiinen (Diptera, Calliphoridae). *Dt. Ent. Ztschr. (N.F.)* 7: 193-235. (1 June)

**Poche, F.**

- 1933 Die Autorschaft einiger Abschnitte der Encyclopédie Méthodique, Histoire Naturelle, Insectes. *Wien. Ent. Zeitung* 53: 142-147. (30 November)

**Poda von Neuhaus, N.**

- 1761 *Insecta musei Graecensis, quae in ordines, genera et species juxta Systema Naturae Caroli Linnaei digessit.* 127 + [xii] pp., 2 pls. Widmanstadii, Graecii [=Graz] (3 September)

[There are two versions of this work: one issued as a university program with a longer and more formal title and 8 extra pages of various university matters; and the other contains only Poda's dissertation. The university program was for 3 September, so the separate work was issued on or before that date. Also, a facsimile edition by W. Junk, Berlin, 1915.]

**Pont, A. C.**

- 1966 Notes on the Muscidae of New Guinea (Diptera). I. The types of Francis Walker. *Ann. Mag. nat. Hist.* (13) 9: 87-99. (1 September)
- 1967 Notes on some Australasian Muscidae (Diptera) described by J. Macquart. *Annls Soc. ent. Fr.* (n.s.) 3: 181-190.
- 1968 The Diptera described by W. J. Rainbow from Funafuti Atoll, Ellice Islands. *Proc. R. ent. Soc. Lond.* (B) 37: 89-90. (4 October)
- 1970a The type-material of Oriental and Australasian Muscidae (Diptera) in the Zoological Museum, Amsterdam. *Beaufortia* 18: 77-111. (28 May)
- 1970b *Myospila hennigi* Gregor and Povolny, 1959 (Dipt., Muscidae), new to Britain, and notes on the European species of *Myospila* Rondani, 1856. *Ent. mon. Mag.* 106: 111-113. (22 December)
- 1972 97. Family Muscidae. 111 pp. In [Papavero, N. (ed.)] *A Catalogue of the Diptera of the Americas South of the United States*. Museu de Zoologia, Universidade de São Paulo. (12 January)
- 1973 Studies on Australian Muscidae (Diptera). IV. A revision of the subfamilies Muscinae and Stomoxyinae. *Australian J. Zool. Suppl.* 21: 129-296. (30 August)
- 1976 Hough's African Muscidae (Diptera). *J. ent. Soc. sth. Africa* 39: 267-270. (September)
- 1977a La faune terrestre de l'Ile de Saint-Hélène, III. 29. Fam. Muscidae (including Anthomyiidae). *Annls Mus. r. Afr. cent. Sér. 8vo (Sci. zool.)* 215: 121-141. (July)
- 1977b Family Muscidae. Pp. 451-523. In Delfinado, M. D. & D. E. Hardy (eds.), *A Catalog of the Diptera of the Oriental region*. Vol. 3. Suborder Cyclorrhapha (excluding Division Aschiza). x + 854 pp. University of Hawaii Press, Honolulu. (19 December)
- 1980 85. Family Muscidae. Pp. 721-761. In Crosskey, R. W. (ed.), *Catalogue of the Diptera of the Afrotropical Region*. 1437 pp. British Museum (Natural History), London. (10 July)

- 1981 The Linnaean species of the families Fanniidae, Anthomyiidae and Muscidae (Insecta: Diptera). *Biol. J. Linnean Soc.* 15: 165-175. (23 February)
- 1984 A revision of the Fanniidae and Muscidae (Diptera) described by Fallén. *Ent. scand.* 15: 277-297. (27 September)
- 1986a Family Muscidae. Pp. 57-215. In Soós, A. & L. Papp (eds.), *Catalogue of Palaearctic Diptera*. Volume 11. Scathophagidae-Hypodermatidae. 346 pp. Akadémiai Kiadó, Budapest. (15 February)
- 1986b A revision of the Fanniidae and Muscidae described by J. W. Meigen (Insecta: Diptera). *Ann. Naturhist. Mus. Wien* 87(B): 197-253, 1 pl. (May)
- 1989 107. Family Muscidae. Pp. 675-699. In Evenhuis, N. L. (ed.), *Catalog of the Diptera of the Australasian and Oceanian Regions*. 1155 pp. Bishop Museum Press & E. J. Brill, Honolulu & Leiden. (23 August)  
[Bishop Museum Special Publication 86.]
- 1990 Muscidae. Pp. 163-170. In Townsend, B. C. (ed.), A Catalogue of the types of bloodsucking flies in the British Museum (Natural History). *Occ. Papers Syst. Ent.* 7, 371 pp. (25 January)
- In Family Muscidae. In Thompson, F. C. (ed), Systematic database of Press Nearctic Diptera. A catalog of names of flies, mosquitoes, gnats, midges and maggots, with information on their classification, distribution and documentation. *USDA Agric. Handbk*
- Pont, A. C. & L. Matile**
- 1976 The identity of *Musca albomaculata* Macquart, 1843, and *Musca dorsomaculata* Macquart, 1843 (Diptera, Muscidae). *Ann. Natal Mus.* 22: 743-746. (30 November)
- Pont, A. C. & V. Michelsen**
- 1982 The Muscoidea described by Moses Harris (Diptera: Fanniidae, Scathophagidae, Anthomyiidae, Muscidae). *Steenstrupia* 8: 25-46. (5 March)
- Pontoppidan, E. L.**
- 1765 *Kurzgefasste Nachrichten, die Naturhistorie in Dänemark betreffend. Aus dem Dänischen übersetzt.* 232 pp., 17 pls. Rothen, Profft & Bock, Kopenhagen & Hamburg. (before 19 December)
- Portschinsky, J. A.**
- 1885 *Muscarum cadaverinarum stercorariarumque biologia comparata.* [In Russian.] *Trudy russk. ent. Obshch.* 19: 210-244. (1 December)
- 1892 Biologie des mouches coprophages et nécrophages. Première partie. [In Russian.] *Trudy russk. ent. Obshch.* 26: 63-131. (14 February)
- 1910 Recherches biologiques sur le *Stomoxys calcitrans* L. et biologie comparée des mouches coprophages. [In Russian.] *Trudy Byuro Ent.* 8(8), 63 + 2 + 90 pp., 1 pl.
- 1911 *Hydrotaea dentipes* F. sa biologie et la destruction par ses larves celles de *Musca domestica* L. [In Russian.] *Trudy Byuro Ent.* 9 (5): 30 pp.

**Presl, J. S.**

- 1822 Additamenta ad Faunam protogaeam, sistens descriptiones aliquot animalium in succino inclusorum. Pp. 191-207. In Presl, J. S. & Presl, C. B. (eds.), *Deliciae Pragenses historiam naturalem spectantes*. viii + 244 pp. Calvae, Pragae [=Prague]. (July)

[Date from *Flora* 5: 576 (Evenhuis in litt.).]

**Preyssler, J. D. E.**

- 1791 Beschreibungen und Abbildungen derjenigen Insekten, welche in Sammlungen nicht aufzubewahren sind, dann aller, die noch ganz neu, und solcher, von denen wir noch keine oder doch sehr schlechte Abbildung besitzen. *Mayer, Samm. Phys. Aufsätze, Dresden* 1: 55-151, 3 pls

1793 [new species in Preyssler, Lindacker & Hofer 1793 (q.v.)]

**Preyssler, J. D. E., J. D. Lindacker & J. K. Hofer**

- 1793 Beobachtungen über Gegenstände der Natur auf einer Reise durch den Böhmerwald im Sommer 1791. *Mayer, Samm. Phys. Aufsätze, Dresden* 3: 135-378, pl. 1.

**Rainbow, W. J.**

- 1897 The atoll of Funafuti, Ellice Group: its zoology, botany, ethnology, and general structure, based on collections made by Mr. Charles Hedley, of the Australian Museum, Sydney, N. S. W. IV. The insect fauna. *Mem. Aust. Mus.* 3: 89-102, pl 1. (25 February)

**Rathvon, S. S.**

- 1855 [new species in Reigert 1855 (q.v.)]

**Ratzeburg, J. T. C.**

- 1840 *Die Forst-Insecten oder Abbildung und Beschreibung der in den Waldern Preussens und der Nachbarstaaten als schädlich oder nützlich bekannt gewordenen Insecten; in systematischer Folge und mit besonderer Rücksicht auf die Vertilgung der Schädlichen*. Vol. 2: Die Falter. viii + 252 pp., 4 tables, 17 pls. Nicolai, Berlin. (14 September)

- 1844 *Die Forst-Insecten oder Abbildung und Beschreibung der in den Waldern Preussens und der Nachbarstaaten als schädlich oder nützlich bekannt gewordenen Insecten; in systematischer Folge und mit besonderer Rücksicht auf die Vertilgung der Schädlichen*. Vol. 3: Die Ader-, Zwei-, Halb-, Netz- und Geradflügler. viii + 314 pp., 5 tables, 16 pls. Nicolai, Berlin. (4 June)

- 1872 *Forstwissenschaftliches Schriftsteller-Lexikon*. xiii + 1-274 [part 1], [ii +] 275-516 pp. [part 2] Fr. Nicolai, Berlin.

**Rautenberg, L. E.**

- 1957 Beiträge zur Kenntnis alter naturwissenschaftlicher Werke. Beitrag 12. *Beitr. Ent.* 7: 609-631. (November)

**Ray, J.**

- 1710 *Historia insectorum. Opus posthumum jussu regiae societas Londinensis editum.* xv + 400 pp. Churchill, Londini [=London].

Razoumowsky—see de Razoumowsky

**Réaumur, R. A. F. de**

- 1738 *Mémoires pour servir à l'histoire des insectes.* Tome quatrième. [ii] + xxxiv + 636 pp., 44 pls. L'Imprimerie Royale, Paris.
- 1740 *Mémoires pour servir à l'histoire des insectes.* Tome cinquième. xliv + 728 pp., 38 pls. L'Imprimerie Royale, Paris.
- [Species redescribed following Linnaean system by Vallot (1802).]

**Reigert, J. F.**

- 1855 *A treatise on the cause of cholera. An interesting discovery.* 15 pp. E. J. Pinkerton, [Philadelphia?].

**Redi, F.**

- 1668 *Esperienze intorno alla generazione degl'insetti.* 228 pp., 29 pls. Stella, Firenze [=Florence].
- 1778 *Opere di Francesco Redi gentilvomo aretino e accademico della crvsca seconda edizone Napoletana corrette e migliorata.* Tomo 1. viii + 180 pp., 16 pls. Michele Stasi, Napoli [=Naples].

**Retzius, A. I.**

- 1783 *Caroli Lib. Bar. De Geer regiae avlæ maresch. r. ord. Wasiaci commend. crvcig. r. ord. de Stella Bor. eqvit. avrat. r. Acad. Scient. Svec. Membr. et Parisinae correspond. Genera et species insectorum e generosissimi avctoris scriptis extraxit, digessit, latine qvoad partem reddit, et terminologiam insectorum linneanam addidit.* vi + 220 + 32 pp. Lebrecht, Lipsiae [=Leipzig]. (8 November)

**Richet, R.**

- 1987 L'identité de la "Mouche à damier", *Sarcophaga carnaria* (Linné, 1758) [Dipt. Sarcophagidae]. *Bull. Soc. ent. France* 91: 131-135 (1986). (March)

**Robineau-Desvoidy, A. J. B.**

- 1830 Essai sur les Myodaires. *Mém. présentés Acad. R. Sci. Inst. France* [Paris] (2)2: 1-813. (5 June)  
[Recorded in the 5 June issue of *Bibliogr. Fr.*.]

- 1863 *Histoire naturelle des Diptères des environs de Paris. Oeuvre posthume du Dr Robineau-Desvoidy publiée par les soins de sa famille, sous la direction de M. H. Monceaux.* Vol. 2, 920 pp. V. Masson et fils, Paris. (early 1863)

**Roemer, J. J.**

- 1789 *Genera Insectorum Linnaei et Fabricii iconibus illustrata.* 86 [+ 3] pp., 37 pls. Henric. Steiner et Socias, Vitoduri [=Winterthur].

**Rognes, K.**

- 1986 The identity of *Sarcophaga alpina* Zetterstedt, 1838 and the interpretation of *Acrophaga* Brauer & Bergenstamm, 1891 (Diptera: Calliphoridae). *Ent. Scand.* 16: 351-354. (10 March)
- 1987a A new species in the *intermedia*-group and a new synonymy in the genus *Pollenia* Robineau-Desvoidy, 1830 (Diptera: Calliphoridae). *Syst. Ent.* 12: 381-388.
- 1987b The taxonomy of the *Pollenia rufa* species-group in the Holarctic region (Diptera: Calliphoridae). *Syst. Ent.* 12: 475-502.
- 1988 The taxonomy and phylogenetic relationships of the *Pollenia semicinerarea* species-group (Diptera: Calliphoridae). *Syst. Ent.* 13: 315-345.
- 1991a Blowflies (Diptera, Calliphoridae) of Fennoscandia and Denmark. *Fauna Ent. Scand.* 24, 272 pp. (January)
- 1991b Revision of the cluster-flies of the *Pollenia viatica* species-group (Diptera: Calliphoridae). *Syst. Ent.* 16: 439-498.
- 1992 Revision of the cluster-flies of the *Pollenia vagabunda* species-group (Diptera: Calliphoridae). *Ent. scand.* 23: 95-114. (April)

**Rondani, C.**

- 1848 Esame di varie specie d'insetti ditteri brasiliiani. In Baudi & Truqui, *Studi Ent. (Turin)* 1: 63-112, pl. 3 bis.
- 1856 *Dipterologiae Italicae Prodromus*. Vol 1. Genera italica ordinis Dipterorum ordinatim disposita et distincta et in familias et stirpes aggregata. 226 [+ 2 pp.]. Stocchi, Parmae [=Parma].
- 1862 *Dipterologiae Italicae prodromus*. Vol. 5: Species Italicae ordinis dipterorum in genera characteribus definita, ordinatim collectae, methodo analitica distinctae, et novis vel minus cognitis descriptis. Pars quarta: Muscidae, Phasiinae-Dexinae-Muscinae-Stomoxidinae. 239 pp. Grazioli, Parmae [=Parma].
- 1868 Diptera Italica non vel minus cognita descripta vel annotata observationibus nonnullis additis. Fasc. III. *Atti Soc. Ital. Sci. Nat.* 11: 21-54.
- 1875 Muscaria exotica musei civici januensis observata et distincta. Fragmentum III. Species in insula Bonae Fortunae (Borneo), Provincia Sarawak, annis 1865-1868, lectae a March. J. Doria et Doct. O. Beccari. *Ann. Mus. civ. Stor. Nat. Genoa* 7: 421-464.

**Roser, C. L. F. von**

- 1840 Erster Nachtrag zu dem im Jahre 1834 bekannt gemachten Verzeichnisse in Württemberg vorkommender zweiflügiger Insekten. *CorrespBl. k. württ. landw. Ver. (n.F.)* 1: 49-64.

**Rossi, P.**

- 1788 Osservazioni insectologiche. *Memorie mat. fis. Soc. ital.* 4: 122-149, pl. 1.
- 1790 *Fauna Etrusca. Sistens insecta quae in provinciis Florentina et Pisana praesertim collegit*. Vol. 1, 272 pp.; vol. 2, 348 pp., 10 pls. Masi, Liburni [=Livorno].

- 1792- *Mantissa insectorum, exhibens species nuper in Etruria collectas,*  
 1794. *adiectis Faunae Etruscae illustrationibus, ac emendationibus.* 2 vols,  
 148 pp., 154 pp., 8 pls. Polloni, Pisis [=Pisa].

**Royer, M.**

- 1931 La collection de Diptères de Robineau-Desvoidy retrouvée et trans-  
 portée au Muséum National d'Histoire Naturelle. *Bull. Ass. Nat. Vall.*  
*Loing* 14: 44-48, portrait. (17 September)

**Rozkosny R., M. Chvála & A. C. Pont**

- 1982 Diptera described by Johann Daniel Preyssler, 1790-1793. *Scr. Fac.*  
*Sci. nat. Univ. purkyn. brun.* 12: 349-356, 4 pls.

Rydbeck, E. O.—see Linnaeus 1758b

**Sabrosky, C. W.**

- 1956 The nomenclature of *Protocalliphora* (Diptera: Calliphoridae). *Proc.*  
*R. ent. Soc. London* (B) 25: 175-179. (October)

- 1958 An overlooked name in "Musca" (Diptera). *Proc. Ent. Soc. Washing-*  
*ton* 60: 134-135. (18 April)

- 1961 Rondani's "Dipterologiae Italicae Prodromus." *Ann. Ent. Soc. Amer.*  
 54: 827-831. (15 November)

- 1974 In defence of Robineau-Desvoidy. *Mosquito Systematics* 6: 220-221.

**Saccà, G.**

- 1947 Sull' esistenza di mosche domestiche resistenti al DDT. *Riv. Parassi-*  
*tol.* 8: 127-128.

**Saccà, G. & L. Rivosecchi**

- 1956a Una nuova sottospecie di 'Musca domestica' L. della regione etiopica.  
*Atti Accad. naz. Lincei Rc.* (8) 19(1955): 497-498.

- 1956b Ricerche sulla speciazione nelle mosche domestiche. III. — Diagnosi  
 differenziale di *M. domestica curviforceps* subsp. nova. *Boll. Zool.*  
 32(1955): 215-224.

**Sachtleben, H.**

- 1944 Ueber einen Rest der Sammlung Bouché's und die in ihm enthaltenen  
 Cocciden (Diptera, Hymenoptera und Hemiptera-Homoptera: Cocco-  
 dae). *Arb. morph. taxon. ent. Berlin* 11: 65-76. (30 September)

1961. Nachträge zu "Walther Horn & Ilse Kahle: Über entomologische  
 Sammlungen." *Beitr. Ent.* 11: 482-540. (September)

**Salem, H. H. & A. F. El Sherif**

- 1960 A new muscoid fly from Gebel Elba (Diptera: Muscidae). *Bull. Soc.*  
*Ent. Egypte* 44: 171-174. (1 July)

**Samouelle, G.**

- 1819 *The Entomologist's useful compendium; or an introduction to the*  
*knowledge of British insects, comprising the best means of obtaining*  
*and preserving them, and a description of the apparatus generally*  
*used; together with the genera of Linné, and the modern method of*  
*arranging the classes Crustacea, Myriapoda, Spiders, Mites and*  
*Insects, from their affinities and structure, according to the views of*

- Dr. Leach. Also an explanation of the terms used in entomology; a calendar of the times of appearance and usual situations of near 3,000 species of British insects, with instructions for collecting and fitting up objects for the microscope.* 496 pp., 12 pls. Boys, London. (June)
- Santos Abreu, E.**
- 1976 *Monografia de los Anthomyidos de las Islas Canarias (Dipteros).* 175 pp., 24 col. figs. Servicio del Aula de Cultura "Elias Santos Abreu", Excmo. Cabildo Insular de la Palma, Canary Is.
- Saunders, E.**
- 1888 *Index to Panzer's Fauna Insectorum Germaniae.* 48 pp. Gurney & Jackson, London.
- Sayre, G.**
- 1959 *Dates of publications describing Musci, 1801-1821.* [vi] + 102 pp. Troy, New York. (April).
- Schäffer [Schaeffer], J. C.**
- 1753 *Die Sattelfliege.* [6 +] 20 pp., 1 color pl. Emanuel Adam Weiss, Regensburg. (1 September)  
[Reviewed in *Göt. Gel. Anz.* 1753: 954 (Evenhuis in litt.)]
- 1764 V. *Die Sattelfliege.* Pp. 241-256, 1 pl. In his *Abhandlungen von Insecten.* Volume 2. 344 pp., 18 pls. Johann Leopold Montag, Regensburg.
- 1766a-*Icones Insectorum circa Ratisbonam indigenorum coloribus naturam referentibus expressae. Natürlich ausgemahlte Abbildungen Regensburgischer Insecten.* Vol. 1, pt. 1, vi + 50 pp., pls. 1-50 (1766a); vol. 1, pt. 2, vi + 50 + xii pp., pls. 51-100 (no date); vol. 2, pt. 1, iv + 50 pp., pls. 101-150 (no date); vol. 2, pt. 2, vi + 50 + vi pp., pls. 151-200 (no date); vol. 3, iv + 80 + vi pp., pls. 201-280 [1779]. Breitfeld, Regensburg.
- 1766b *Elementa Entomologica. CXXXV. tabvlae aere excvsae floridisqve coloribvs distinctae. Einleitung in die Insectenkenntnis. CXXXV. ausgemahlte Kupfertafeln.* [7 + 8 + 17 pp.,] + 135 plates with facing legends. Weissianis, Ratisbonae [=Regensburg].
- Scharfenberg, G. L.**
- 1805 [new species in Bechstein & Scharfenberg (q. v.)]  
[According to the introduction, new species are to be credited to Scharfenberg alone.]
- Schellenberg, J. R.**
- 1803 *Genres des Mouches Diptères représentés en XLII. planches projetées et dessinées par Mr. J. R. Schellenberg, et expliquées par deux amateurs de l'Entomologie. Gattungen der Fliegen in XLII. Kupfer-tafeln entworfen und gezeichnet von J. R. Schellenberg, und erklärt durch zwey Liebhaber der Insektenkunde.* 95 pp., 42 col. pls. Orell and Fuesli, Zürich.

[As clearly indicated in the title and introduction, Schellenberg only provided the plates, while the text and names were provided by two anonymous amateurs. However, as these anonymous authors are unknown, we follow tradition and attribute authorship of the whole work to Schellenberg.]

**Schiner, I. R.**

- 1856 Scriptores austriaci rerum dipterologicarum. Eine Revision der von österreichischen Entomologen aufgeführten Dipteren. *Verhandl. K.-k. Zool.-Bot. Ver.*, Wien 6: 399-424.
- 1857 Diptera Austriaca. Aufzählung aller im Kaiserthume Oesterreich bisher aufgefundenen Zweiflügler. III. Die österreichischen Syrphiden. *Verhandl. K.-k. Zool.-bot. Ges.* Wien 7: 279-506.
- 1860- *Fauna Austriaca. Die Fliegen (Diptera). Nach der analytischen Methode bearbeitet, mit der Characteristik sämmtlicher europäischer Gattungen, der Beschreibung aller in Deutschland vorkommenden Arten und der Aufzählung aller bisher beschriebenen europäischen Arten.* Vol. 1, lxxx + 674 pp. H. 1, pp. 1-72; H. 2, pp. 73-184, pl. 1 (1860). H 3/4, pp. 185-368; H. 5, pp. 369-440 (1861). H 6/7, pp. 441-656; H. 8 (part), pp. 657-674 + i-lxxx, pl. 2 (1862). Gerold, Wien [=Vienna].
- 1868 Diptera. In [Wüllerstorf-Urbair, B. von], *Reise der österreichischen Fregatte Novara um die Erde in den Jahren 1857, 1858, 1859, unter den Befehlen des Commodore B. von Wüllerstorf-Urbair. Zoologischer Theil* 2, 1 (B). vi + 388 pp., 4 pls. Gerold, Wien [= Vienna]. (30 April)

**Schnabl, J. & H. Dziedzicki**

- 1911 Die Anthomyiden. *Nova Acta Acad. Caesar. Leop. Carol.* 95: 53-358. pl. 3-37.

**Schrantz, F. von P.**

- 1776 Verzeichniss einiger Insekten, derer im Linneeanischen Natursysteme nicht gedacht wird. 7th Abh., pp. 59-98. In his *Beyträge zur Naturgeschichte*. 137 pp., 7 pls. Fritsch, Leipzig. (21 April)
- 1781 *Envmeratio insectorum Avstriae indigenorum.* xxiv + 548 + [4] pp., 4 pls. Eberhardi Klett et Franck, Augustae Vindelicorum [=Augsburg] (June)  
[Reviewed in *Allg. Verz. Neuer Bücher* 6: (Evenhuis in litt.).]
- 1796 Naturhistorische Beobachtungen um Pöttmes, Neuburg und Weihering. Pp. 97-226. In his *Sammlung naturhistorischer und physikalischer Aufsätze*. xvi + 456 pp., 7 pls. Raspe, Nürnberg.
- 1803 *Favna Boica. Durchgedachte Geschichte der in Baiern einheimischen und zahmen Thiere.* Vol. 3, Pt. 1. viii + 272 pp. Krüll, Landshut.
- 1837 [Names and *nomina nuda* in Gistl 1837a]

**Schumann, H.**

- 1974 Revision der palaearktischen *Bellardia*-Arten (Diptera, Calliphoridae). *Dt. ent. Ztschr.* (n.F.) 21: 231-299. (10 December)

**Schummel, T. E.**

- 1832 [Notes on order Diptera]. In Gravenhorst, J. C. Bericht der entomologischen Section in der Schlesischen Gesellschaft für vaterländische

Kultur, am Ende des Jahres 1831. *Uebers. Arb. Schles. Ges.* 1831: 72-77.

**Schwenckfeld, C.**

- 1603 *Theriotrophevm Silesiae, in quo animalium, hoc est, quadrupedum, reptilium, avium, piscium, insectorum natura, vis & usus sex libris perstringuntur.* [23 +] 563 pp. Davidis Alberti, Lignicili [=Legnica].

**Scopoli, J. A.**

- 1763 *Entomologia carniolica exhibens insecta carnioliae indigena et distributa in ordines, genera, species, varietates, methodo Linnaeana.* 421 pp. Trattner, Vindobonae [=Vienna]. (before 23 June)
- 1772 *Annus historico naturalis.* Vol. 5. 128 pp. Hilscheri, Lipsiae [=Leipzig]. (21 November)
- 1777 *Introvdvctio ad historiam natvralem sistens genera lapidvm, plantarvm, et animalivm hactenvs detecta, caracteribvs essentialibvs donata, in tribvs divisa, svbinde ad leges Natvrae.* 506 pp. Gerle, Pragae [=Prague]. (13 April)

[Appeared at the Leipzig Easter Book Fair (*cf. Schröter's J. 4: 482*) (Evenhuis in litt.).]

- 1780 Zwölfter Brief. Ueber eine Gattung schädlicher Insekten, die man die kolombachischen Mücken nennet, welche unter den Ochsen, Schafen, Ziegen, Schweinen und Pferden in Serviert, dem Temeswarer Bannat, und der abendländischen Wallachen grossen Schaden anrichten. Pp. 123-135 in Griselini, F. (ed.), *Versuch einer politischen und natürlichen Geschichte des temeswarer Banats in Briefen an Standespersonen und Gelehrte.* Zweyter Theil. 135 pp., 9 pls. Johann Paul Krauss, Wien

- 1972 *Entomologia Carniola.* xxxiii + 7 pls. + 421 pp., 43 pls. Akademische Druck-u. Sanstalt, Graz.

[This reprint consists of an biography of Scopoli by Otto Guuglia followed by a photo facsimile of Scopoli (1763) and the unpublished plates.]

**Scudder, S. H.**

- 1877 The first discovered traces of fossil insects in the American Tertiaries. *Bull. U. S. geol. geogr. Surv. Territ.* 3: 741-762.
- 1891 Index to the known fossil insects of the World including myriapods and arachnids. *Bull. U. S. Geol. Surv.* 71: 3-744.

**Séguy, E.**

- 1928 Mouches parasites. I. Conopides, Œstrides et Calliphorines de l'Europe occidentale. *Encycl. Ent.* (A)9: 3-251. (21 February)
- 1937 Diptera Fam. Muscidae. In Wytsman, P. (ed.), *Genera Insectorum.* 205: 604 pp., 9 pls. Desmet-Verteneuil, Bruxelles. (23 October)

[Recorded in 23 October issue of *Naturae Novitates*.]

**Shaw, G.**

- 1806 *General zoology or systematic natural history.* Volume 6, part 2, Insecta. Pp. i-x, 241-509, pls 80-137. Kearsley, London.

**Sherborn, C. D.**

- 1902 *Index Animalium sive index nominum quae ab A. D. MDCCCLVIII generibus et speciebus animalium imposita sunt. Sectio Prima. A Kalendis Ianuariis, MDCCCLVIII usque ad finem Decembris, MDCCC. lix + 1195 pp.* Cantabridgiae [=Cambridge, England]
- 1922 *Sectio Secunda. A kalendis Ianuariis, MDCCCI usque ad finem Decembris, MDCCCL. cxxxi + 7056 pp.* British Museum, London.
- 1932 *Sectio Secunda. A kalendis Ianuariis, MDCCCI usque ad finem Decembris, MDCCCL. Epilogue, additions to Bibliography, additions and corrections, and index to trivialia. cxlviii + 1098 pp.* British Museum, London.

**Shinonaga, S. & R. Kano**

- 1971 *Muscidae (Insecta: Diptera). Vol. I. 242 pp., 28 pls. In Fauna Japonica.* Academic Press of Japan, Tokyo.  
[The publication date of "March" is certainly incorrect. Most probable date is late June.]
- 1977 A new species of the genus *Musca* L. from the Oriental Region. *Japan. J. Sanit. Zool.* 28: 111-113. (15 June)

**Skidmore, P.**

- 1985 The biology of the Muscidae of the world. *Series Ent.* 29. xiv + 550 pp. W. Junk, The Hague. (February)

**Smith, A. Z.**

- 1986 *A history of the Hope Entomological Collections in the University Museum Oxford with lists of archives and collections.* xiv + 172 pp., Clarendon Press, Oxford

**Smith, K. G. V., R. W. Crosskey & A. C. Pont**

- 1980 Bibliography of cited literature. Pp. 889-1196. In Crosskey, R. W. (ed.), *Catalogue of the Diptera of the Afrotropical Region.* 1437 pp. British Museum (Natural History), London. (10 July)

**Smith, K. G. V. & E. Taylor**

- 1964 On the location and recognition of some Walker types of Diptera from the W. W. Saunders collection. *Ent. mon. Mag.* 100: 21-33. (8 September)

**Snyder, F. M.**

- 1951 Some old and new species of Muscinae from the Ethiopian region (Diptera, Muscidae). *Amer. Mus. Novit.* 1533, 42 pp. (1 October)
- 1965 Diptera: Muscidae. *Insects Micronesia* 13: 191-327, 2 tables. (22 March)

Söderberg, D. H.—see Linnaeus 1771

**Sommerfelt, S. C.**

- 1824- Physisk-oeconomisk Beskrivelse over Saltdalen i Nordlandene. K.
- 1827 *nor. Vidensk. Selsk. Skr.* [2]2(1): 1-148.

**Speiser, P.**

- 1924 Eine übersicht über die Dipterenfauna Deutsch-Ostafrikas. Pp. 90-156. In Braun, M., *Beiträge aus der Tierkunde. Herrn Geh.*

*Regierungsrat Prof. Dr. med. et phil. M. Braun aus Anlass seines goldenen medizinischen Doktor-Jubiläums als Festgabe dargebracht von Schülern und Freunden.* Gräfe und Unzer, Königsberg 156 p., 3 pls. (September)

**Spilman, T. J.**

- 1967 Gmelin's 13th Edition of the *Systema Naturae*: A case of neglect. *Ent. News* 78: 169-172.

**Stafleu, F. A. & R. S. Cowan**

- 1976- *Taxonomic Literature. A selective guide to botanical publications and collections with dates, commentaries and types.* 2nd Edition. Vol. 1: A-G, xl + 1136 pp. (1976); II: H-Le, xviii + 991 pp. (1979); III: Lh-O, xii + 980 pp. (1981); IV: P-Sak, ix + 1214 pp. (1983); V: Sal-Ste, [vi +] 1066 pp. (1985); VI: Sti-Vuy, [vi +] 926 pp. (1986); VII: W-Z, lvi + 653 pp. (1988). Bohn, Scheltema & Holkema, Utrecht.

**Stein, J. R. Am**

- 1857 Bündner Dipteren. Verzeichniss der Sammlung des Hrn. Major Am Stein in Malans. *Jber. naturf. Ges. Graubündens* (n. F.) 2: 89-111.

**Stein, P.**

- 1903 Die Bestimmung und Beschreibung der Muscinen und Anthomyinen mit Aushnahme der Gattung *Lispa* Latr. Pp. 99-122. In Becker, T., Aegyptische Dipteren. *Mitt. zool. Mus. Berlin* 2: 1-66 (12 August 1902); 67-195, pl. 4b-4e (23 September 1903)

- 1909 Neue javanische Anthomyiden. *Tijdschr. Ent.* 52: 205-271. (31 December)

- 1910 The Percy Sladen Trust expedition to the Indian Ocean in 1905, under the leadership of Mr. J. Stanley Gardiner. Vol. 3. No. IX. - Diptera, Anthomyidae, mit den Gattungen *Rhinia* und *Idiella*. *Trans. Linn. Soc. London* (2, Zool.) 14: 149-163. (November)

- 1913 Neue afrikanische Anthomyiden. *Annls hist.-nat. Mus. natn. hung.* 11: 457-583. (24 December)

- 1916 Die Anthomyiden Europas. Tabellen zur Bestimmung der Gattungen und aller mir bekannten Arten, nebst mehr oder weniger ausführlichen Beschreibungen. *Arch. Naturgesch.* 81 A 10 [1915]: 1-224. (June)

- 1919 Die Anthomyidengattungen der Welt, analytisch bearbeitet, nebst einem kritisch-systematischen Verzeichnis aller aussereuropäischen Arten. *Arch. Naturgesch.* 83 A 1 [1917]: 85-178. (February)

**Stephens, J. F.**

- 1829a *The Nomenclature of British Insects; being a Compendious List of such Species as are contained in the Systematic Catalogue of British Insects, and forming a guide to their classification, &c. &c.* 68 pp. Baldwin & Cradock, London. (1 June)

- 1829b *A systematic catalogue of British insects: being an attempt to arrange all the hitherto discovered indigenous insects in accordance with their natural affinities. Containing also the references to every English*

writer on entomology, and to the principal foreign authors. With all the published British genera to the present time. Part II. *Insecta Haustellata*. 388 pp. Baldwin & Cradock, London. (15 July)

**Steyskal, G. C.**

- 1968 48. Family Micropezidae. 33 pp. In [Papavero, N. (ed.)], *A Catalogue of the Diptera of the Americas south of the United States*. Museu de Zoologia, Universidade de São Paulo
- 1974a On the life and influence of J. W. Meigen by J. A. Förster, Stettin. Entomol. Ztg. 1846. 66-74; 130-140. [A translation.] *Mosquito Systematics* 6: 79-87.
- 1974b Notice nécrologique sur Meigen by J. Macquart, Ann. Soc. Ent. Fr. 16: 323-334, 1847. [A translation, in part.] *Mosquito Systematics* 6: 87-88.

**Sulzer, J. H.**

- 1761 *Die Kennzeichen der Insekten, nach Anleitung des Königl. Schwed. Ritters und Leibarzts Karl Linnaeus, durch XXIV. Kupfertafeln erläutert und mit derselben natürlichen Geschichte begleitet.* xxviii + 203 + 67 p., 24 pls. Heidegger, Zürich.
- 1776 *Abgekürzte Geschichte der Insecten nach dem Linaeischen System.* Vol. 1, xxviii + 274 + 71 pp., 32 pls. Steiner, Winterthur. (21 April)  
[Shown at Leipzig Easter book fair (Cobres 1782: 380).]

**Swammerdam, J. J.**

- 1685 *Historia Insectorum generalis, in qua quaecunque ad insecta eorumque mutationes spectant, dilucide ex sanioris philosophiae & experientiae principiis explicantur, cum figuris & indicibus necessariis.* [16+] 212 [+ 17] pp. Luchtmans, Lugd. Batavorum [= Leiden]

**Swederus, N. S.**

- 1787 Et nytt genus och femtio nya species af insecter. *K. Vetenskapsakad. Nya Handl.* (2) 8: 181-201, pl. 8, 276-290.

**Thompson, F. C.**

- 1973 De Geer's exotic *Musca* species (Diptera: Syrphidae and Calliphoridae). *Proc. ent. Soc. Washington* 75: 354-356. (5 October)
- 1980 The problem of old names as illustrated by *Brachyopa "conica* Panzer," with a synopsis of the Palaearctic species of *Brachyopa*. *Ent. Scand.* 11: 209-216. (7 July)
- 1981a The flower flies of the West Indies (Diptera: Syrphidae). *Mem. ent. Soc. Washington* 9, 200 pp. (2 September)
- 1981b Nomenclature of the European species of *Neoascia* Meigen (Diptera: Syrphidae). *Ent. Scand.* 12: 470-478. (15 September)
- 1988 Syrphidae (Diptera) described from unknown localities. *J. New York ent. Soc.* 96: 200-226. (28 April)
- 1990 Standard data elements for classification and nomenclature. 7 pp. In Thompson, F. C. (coordinator), R. A. Hellenthal, J. Louton, G. R. Noonan, R. T. Schuh & M. K. Thayer, *Automatic Data Processing for*

*Systematic Entomology: Promises and Problems. A report for the Entomological Collection Network.* 48 pp. Entomological Collections Network, Baton Rouge

**Thompson, F. C., J. R. Vockeroth & M. C. D. Speight**

- 1982 The Linnaean species of flower flies (Diptera: Syrphidae). *Mem. ent. Soc. Washington* 10: 150-165 (30 September)

**Thomson, C. G.**

- 1868 6. Diptera. Species novas descriptis. Pp. 443-614, pl. 9. In K. Svenska Vetenskaps-Akademien, *Kongliga svenska fregatten Eugenies resa omkring jorden under befäl af C. A. Virgin, åren 1851-1853.* Pt. 2: Zoologie, 1; Insekter, 617 pp., 9 pls. P. A. Norstedt & Söner, Stockholm. (December)

**Thomson, R. C. M.**

- 1937 Observations on the biology and larvae of the Anthomyidae. *Parasitology* 29: 273-358. (5 July)
- 1947 Notes on the breeding habits and early stages of some Muscids associated with cattle in Assam. *Proc. R. ent. Soc. Lond. (A)* 22: 89-100. (20 December)

**Thunberg, C. P.**

- 1789 *D. D. Museum Naturalium Academiae Upsaliensis. Cujus Partem Septiman.* [ii] + pp. 85-94. Upsaliae [=Uppsala]. (7 March)
- 1791 *D. D. Museum Naturalium Academiae Upsaliensis. Appendix II.* [i] + pp. 123-130. Upsaliae [=Uppsala]. (April)

**Topsel, E.**

- 1656 *The history of four-footed beasts and serpents: describing at large their true and lively figure, their several names, conditions, kinds, virtues (both natural and medicinal), countries of their breed, their love and hatred to mankind, and the wonderful work of God in their creation, preservation and destruction. Interwoven with curious variety of historical narrations out of scriptures, fathers, philosophers, physicians, and poets: illustrated with divers hieroglyphicks and emblems, &c. both pleasant and profitable for students in all faculties and professions. Collected out of the writings of Conradus Gesner and other authors. Whereunto is now added, The Theater of Insects; or, lesser living creatures: as bees, flies, caterpillars, spiders, worms, &c. A most elaborate work: by T. Muffet, Dr. of Physick. The whole revised, corrected, and enlarged with the addition of two useful physical tables, by J. R. M. D.* 1130 pp. Cotes, London.

[Muffet's "Theater of insects" is a self-contained section of the book, pp. 889-1130, with its own title page on which the author's name is rendered "Mouffet." "J. R." is presumably Dr. John Ray.]

**Townsend, C. H. T.**

- 1911 Review of work by Pantel and Portchinski on reproductive and early stage characters of muscoid flies. *Proc. ent. Soc. Wash.* 13: 151-170. (30 September)

- 1913 Inquiry into the relationships and taxonomy of the muscoid flies. *Can. Ent.* 45: 37-57. (10 February)
- 1915 Correction of the misuse of the generic name *Musca*, with description of two new genera. *J. Wash. Acad. Sci.* 5: 433-436. (19 June)
- 1921 Some new muscoid genera ancient and recent. *Insecutor Inscit. menstr.* 9: 132-134. (3 October)
- 1931a Notes on American Oestromuscoid types. *Revta Ent.* 1: 65-104 (25 April)
- 1931b Notes on American Oestromuscoid types. *Revta Ent.* 1: 157-183 (July)
- 1931c Notes on Old-World Oestromuscoid types. — Part I. *Ann. Mag. nat. Hist.* (10) 8: 369-391. (October)
- 1931d New genera and species of American Oestromuscoid flies. [conclusion]. *Revta Ent.* 1: 437-479. (14 November)
- 1932 Notes on Old-World Oestromuscoid types. — Part II. *Ann. Mag. nat. Hist.* (10) 9: 33-57. (January)
- 1937 Muscoid generic diagnoses and data. Glossinini to Agriini. Part V. 234 pp. In his *Manual of Myiology in twelve parts*. Townsend, Itaquaquecetuba, São Paulo.

**Tumrasvin, W. & S. Shinonaga.**

- 1977 Studies on medically important flies in Thailand. III. Report of species belonging to the genus *Musca* Linné, including the taxonomic key (Diptera: Muscidae). *Bull. Tokyo med. dent. Univ.* 24: 209-218. (September)

**Turton, W.**

- 1801 A general system of nature, through the three grand kingdoms, of animals, vegetables, and minerals: Systematically divided into their several classes, orders, genera, species, and varieties, with their habitations, manners, economy, structure and peculiarities. Translated from Gmelin's last edition of the celebrated *Systema Naturae*, by Sir Charles Linné: Amended and enlarged by the improvements and discoveries of later naturalists and Societies, with appropriate copper-plates. "1800" Vol. III, 784 pp. David Williams, Swansea.  
[The date is taken from that on the colophon.]

**Tuxen, S. L.**

- 1967a The entomologist, J. C. Fabricius. *Ann. Rev. Ent.* 12: 1-14, portrait.
- 1967b Bibliographie von I. C. Fabricius. *Zool. Anzeiger* 178: 174-185.
- 1973 Entomology systematizes and describes: 1700-1815. Pp. 95-118. In Smith, R. F., T. E. Mittler & C. N. Smith [eds], *History of Entomology*. vii + 517 pp. Annual Reviews Inc., Palo Alto, California.

**Vallot, J. N.**

- 1802 *Concordance systématique, servant de table des matières à l'ouvrage de Réaumur intitulé: Mémoires pour servir à l'Histoire des Insectes*. 196 + 11 pp. Gregoire & Thouvenin, Paris ("An X") (before 22 September)

**Vane-Wright, R. I.**

- 1975 The butterflies named by J. F. Gmelin (Lepidoptera: Rhopalocera). *Bull. Br. Mus. nat. Hist. (Ent.)* 32: 17-64, 6 pls. (11 April)

**Verrall, G. H.**

- 1888 *A list of British Diptera*. 34 + 4 + 12 pp. Pratt, London.
- 1901a *British flies*. Vol. VIII. Platypezidae, Pipunculidae and Syrphidae of Great Britain. [ii] + 691 pp. Gurney & Jackson, London. (1 January)
- 1901b *A list of British Diptera*. 2nd edition. 47 pp. Cambridge University Press, Cambridge. (December)

**Villeneuve, J.**

- 1916 A contribution to the study of the South African higher Myodarii (Diptera Calypratae) based mostly on the material in the South African Museum. *Ann. S. Afr. Mus.* 15: 469-515. (8 December)
- 1922 Descriptions d'espèces nouvelles du genre "Musca." *Annls Sci. Nat. (Zool., 10)* 5: 335-336. (October)
- 1936 Description d'une nouvelle espèce africaine du genre *Musca*. *Bull. Annls Soc. Ent. Belg.* 76: 414. (31 December)
- 1937 Myodaires supérieurs africains. *Bull. Mus. R. Hist. Nat. Belg.* 13 (35), 4 pp. (August)

Villers, C. J. de—see de Villers, C. J.

**Vockeroth, J. R.**

- 1953 *Musca autumnalis* Deg. in North America (Diptera: Muscidae). *Can. Ent.* 85: 422-423. (4 December)
- 1969 A revision of the genera of the Syrphini (Diptera: Syrphidae). *Mem. Ent. Soc. Canada* 62, 176 pp. (15 July)
- 1971 The identity of some Holarctic and Old World species of *Sphaerophoria* (Diptera: Syrphidae). *Can. Ent.* 103: 1627-1634. (November)

**Voss, W.**

- 1881 Ioannes Antonius Scopoli. Lebensbild eines österreichischen Naturforschers und dessen Kenntnisse der Pilze Krains. *Verh. zool.-bot. Ges. Wien* 31: 17-66, portrait.

**Walckenaer, C. A.**

- 1802 *Faune Parisienne, Insectes. Ou histoire abrégée des Insectes des environs de Paris, classés d'après le Système de Fabricius, précédée d'un Discours sur les Insectes en général, pour servir d'introduction à l'étude de l'Entomologie*. Vol. 2, xxii + 438 pp. Dentu, Paris. ["an XI"]. (after 22 September)

**Walker, F.**

- 1836 Descriptions, & c. of the Diptera. Pp. 331-359. In Curtis, J., A. H. Haliday and F. Walker, Descriptions, & c. of the Insects collected by Captain P. P. King, R. N., F. R. S., in the Survey of the Straits of Magellan. *Trans. Linn. Soc. London* 17: 315-359. (9 July)

- 1849 *List of the specimens of dipterous insects in the collection of the British Museum.* Vol. 4, pp. 689-1172. British Museum, London. (8 December)
- 1851 Diptera. Vol. 1. 314 pp., pls. 1-10. In [Walker, F., H. T. Stainton, and S. J. Wilkinson], *Insecta Britannica*. Reeve and Benham, London. (2 February)
- 1853a Diptera. Vol. 2, vi + 298 pp., pls. 11-20. In [Walker, F., H. T. Stainton, and S. J. Wilkinson], *Insecta Britannica*. Lovell Reeve, London.
- 1853b *Insecta saundersiana: or characters of undescribed insects in the collection of William Wilson Saunders, Esq., F.R.S., F.L.S., &c.* Vol. 1. Diptera. [Part 4] Pp. 253-414, pls. 7-8. van Voorst, London. (before 7 March)
- 1856a Catalogue of the dipterous insects collected at Singapore and Malacca by Mr. A. R. Wallace, with descriptions of new species. *J. Proc. Linn. Soc. London (Zool.)* 1: 4-39, pls. 1-2. (1 March)
- 1856b Catalogue of the dipterous insects collected at Sarawak, Borneo, by Mr. A. R. Wallace, with descriptions of new species. *J. Proc. Linn. Soc. London (Zool.)* 1: 105-136, pl. 6. (1 November)
- 1858a Characters of undescribed Diptera in the collection of W. W. Saunders, Esq., F.R.S., &c. [part.] *Trans. Ent. Soc. London (N.S.)* 4: 190-235. (January)
- 1858b Catalogue of the dipterous insects collected in the Aru Islands by Mr. A. R. Wallace, with descriptions of new species. [part] *J. Proc. Linn. Soc. London (Zool.)* 3: 77-110. (1 November)
- 1859a Catalogue of the dipterous insects collected in the Aru Islands by Mr. A. R. Wallace, with descriptions of new species. [concl.] *J. Proc. Linn. Soc. London (Zool.)* 3: 111-131. (1 February)
- 1859b Catalogue of the dipterous insects collected at Makessar in Celebes, by Mr. A. R. Wallace, with descriptions of new species. [part] *J. Proc. Linn. Soc. London (Zool.)* 4: 97-144. (8 December)
- 1860 Catalogue of the dipterous insects collected in Amboyna by Mr. A. R. Wallace, with descriptions of new species. *J. Proc. Linn. Soc. London (Zool.)* 5: 144-168. (18 July)
- 1861a Characters of undescribed Diptera in the collection of W. W. Saunders, Esq., F.R.S., & c. [Part] *Trans. Ent. Soc. London (2)* 5: 297-334. (February)
- 1861b Catalogue of the dipterous insects collected at Dorey, New Guinea, by Mr. A. R. Wallace, with descriptions of new species. *J. Proc. Linn. Soc. London (Zool.)* 5: 229-254. (27 March)
- 1861c Catalogue of the dipterous insects collected at Manado in Celebes and in Tond, by Mr. A. R. Wallace, with descriptions of new species. [Part] *J. Proc. Linn. Soc. London (Zool.)* 5: 258-264. (27 March)
- 1861d Catalogue of the dipterous insects collected in Batchian, Kaisaa and Makian, and at Tidon in Celebes, by Mr. A. R. Wallace, with descrip-

- tions of new species. *J. Proc. Linn. Soc. London (Zool.)* 5: 270-303. (24 May)
- 1864 Catalogue of the dipterous insects collected in Waigou, Mysol, and North Ceram, by Mr. A. R. Wallace, with descriptions of new species. *J. Proc. Linn. Soc. London (Zool.)* 7: 202-238. (5 April)
- Walker, K. L.**
- 1985 Mystery of Francis Walker specimens solved. *Antenna* 9: 78-80. (April)
- Weiss, H. B.**
- 1926 Two entomologists of the eighteenth Century -- Eleazar Albin and Moses Harris. *Science Mon., New York* 23: 558-564.
- 1927 Dru Drury, silversmith and entomologist of the eighteenth century. *Ent. News* 38: 208-214 (1 July)
- West, L. S.**
- 1951 *The housefly. Its natural history, medical importance and control.* xi + 584 pp. Comstock, Ithaca, New York
- West, L. S. & O. B. Peters**
- 1973 *An annotated bibliography of Musca domestica Linnaeus.* xiii + 743 pp. Dawsons, Pall Mall, London.
- Westring, J. P.**
- 1814 Om ett slags Flug-larver i sår, och deras förvandling; anmärkte och beskrifne. *K. svenska Vetenskapsakad. Handl.* 181: 51-56, pl. 4b.
- Westwood, J. O.**
- 1840 Order XIII. Diptera Aristotle. (Antliata Fabricius. Halteriptera Clairv.). Pp. 125-154. in his *An introduction to the modern classification of insects; founded on the natural habits and corresponding organisation of the different families. Synopsis of the genera of British insects.* 158 pp. Longman, Orme, Brown, Green & Longmans, London. (6 June)  
[Reviewed in *Athenaeum* 658: 458 (Evenhuis *in litt.*)]
- White, A.**
- 1843 [new species in White & Doubleday, 1843.]
- White, A. & E. Doubleday**
- 1843 List of the annulose animals hitherto recorded as found in New Zealand, with the descriptions of some new species. Pp. 265-295. In Gray, J. E., Fauna of New Zealand. Materials towards a fauna of New Zealand, Auckland Island, and Chatham Islands. Pp. 177-296. In Dieffenbach, E., *Travels in New Zealand; with contributions to the geography, geology, botany, and natural history of that country.* Vol. II, iv + 396 pp. John Murray, London. (21 January)  
[Reviewed in *Athenaeum* 795: 61 (Evenhuis *in litt.*)]
- White, I. M.**
- 1987 The Linnaean species of the family Tephritidae (Insecta: Diptera). *Zool. J. Linn. Soc.* 90: 99-107.

**Wiedemann, C. R. W.**

- 1817 Neue Zweiflügler (Diptera Linn.) aus der Gegend um Kiel. *Zool. Mag. (Wiedemann's)* 1(1): 61-86. (July)  
 [Recorded in the July issue of *Isis (Oken's)*.]
- 1818 Neue Insecten vom Vorgebirge der guten Hoffnung. *Zool. Mag. (Wiedemann's)* 1(2): 40-48.
- 1819a Beschreibung neuer Zweiflügler aus Ostindien und Afrika. *Zool. Mag. (Wiedemann's)* 1(3): 1-39.
- 1819b Brasilianische Zweiflügler. *Zool. Mag. (Wiedemann's)* 1(3): 40-56.
- 1824 *Munus rectoris in Academia Christiana Albertina aditurus Analecta entomologica ex Museo Regio Havniensi maxime congesta profert iconibusque illustrat.* 60 pp. E regio typographeo scholarum, Kiliae [=Kiel].
- 1825 [new species in Germar, E. F. (q.v.)]
- 1830 *Aussereuropäische zweiflügelige Insekten.* Vol. 2, xii + 684 + xi pp., 5 pls. Schulz, Hamm. (14 December)  
 [Recorded in the 14 December issue of *Isis (Oken's)*.]

**Wilkinson, R. S.**

- 1980 The Thadeus[sic] W. Harris and Samuel H. Scudder Papers. *Ent. J. Var.* 92: 225-226. (September)

**Wilson, J.**

- 1835 *A treatise on insects, general and systematic; being the article "Entomology," from the seventh edition of the Encyclopaedia Britannica.* 327 pp., pls. 220-239. Black, Edinburgh. (November)

**Wotton, E.**

- 1552 *Edoardi Wottoni Oxoniensis de differentiis animalium libri decem.* [22 +] 220 double-spread [+ 26] pp. Lutetiae Parisiorum [= Paris] apud Vascosanum.

**Yarom, I.**

- 1990 A review of the genus *Sapromyza* Fallén in Israel with remarks on *S. (Sapromyzosoma) quadripunctata* (Linnaeus)(Diptera: Lauxaniidae). *Ent. Scand.* 21: 289-304. (October)

**Yeats, T. P.**

- 1773 *Institutions of Entomology: being a translation of Linnaeus's ordines et genera insectorum; or, systematic arrangement of insects. Collated with the different systems of Geoffroy, Schaeffer and Scopoli; together with observations of the translator.* viii + 272 pp. Horsfield, London.

**Zetterstedt, J. W.**

- 1837 *Conspectus familiarum, generum et specierum dipterorum, in Fauna Insectorum Lapponica descriptorum.* *Isis, Jena* 21: 28-67. (January)
- 1838 *Sectio Tertia: Diptera,* pp. 477-868. *In his Insecta Lapponica.* vi + 1,140 (pp.). L. Voss, Lipsiae [=Leipzig]. (before September)
- 1842- *Diptera Scandinaviae. Disposita et descripta.* Vol. 1, pp. iii-xvi + 1860 1-440 (1842); vol. 2, pp. 441-894 (1843); vol. 3, pp. 895-1280 (1844);

vol. 4, pp. 1281-1738 (1845); vol. 5, pp. 1739-2162 (1846); vol. 6, pp. 2163-2580 (1847); vol. 7, pp. 2581-2934 (1848); vol. 8, pp. 2935-3366 (1849); vol. 9, pp. 3367-3710 (1850); vol. 10, pp. 3711-4090 (1851); vol. 11, pp. v-xii + 4091-4546 (1852); vol. 12: Sup. 3, pp. v-xx + 4547-4942 (1855); vol. 13: Sup. 4, pp. v-xvi + 4943-6190 (1859); vol. 14, pp. 6191-6609 (1860). *Lundbergiana*, Lundae [=Lund].

#### Zielke, E.

- 1970 Fundorte afrikanischer Muscinae (Diptera, Muscidae). *Z. angew. Zool.* 57: 499-510.
- 1971a Description of six new species of Muscidae (Diptera) from the Ethiopian region. *Proc. Biol. Soc. Washington* 84: 103-111. [30 June]
- 1971b Muscinae vom südlichen Afrika. *Zool. Anz.* 187: 214-218.
- 1971c Revision der Muscinae der äthiopischen Region. *Series Ent.* 7, viii + 199 pp. W. Junk, The Hague.
- 1971d Beitrag zur Kenntnis der Verbreitung afrikanischer Musciden (Muscidae; Diptera). *Ent. Mitt. zool. Mus. Hamburg* 4: 173-181. (15 August)
- 1973 Beitrag zur Kenntnis der Verbreitung afrikanischer Musciden (Muscidae; Diptera). Teil II: Canadian National Collection of Insects. *Z. angew. Zool.* 59: 361-375.
- 1974 Beitrag zur Kenntnis der Verbreitung afrikanischer Musciden (Muscidae; Diptera). Teil III: Sammlung der California Academy of Sciences. *Z. angew. Zool.* 61: 1-44.

#### Zimin, L. S.

- 1948 Key to the larvae of synanthropic flies of Tadzhikistan (in the third stage). [In Russian] *Opred. Faune SSSR* 28, 114 pp., 61 figs.
- 1951 Insects-Diptera, 18, No. 4. Family Muscidae, true flies (Tribes Muscini and Stomoxydini). [In Russian] *Fauna SSSR* (n. s.) 45, 286 pp.

#### Zimsen, E.

- 1954 The insect types of C. R. W. Wiedemann in the Zoological Museum in Copenhagen. *Spolia zool. Mus. haun.* 14, 43 pp.
- 1964 *The type material of I. C. Fabricius*. 656 pp., 2 pl. Munksgaard, Copenhagen.

#### Zschach, J. J.

- 1788 *Museum N. G. Leskeanum. Pars entomologica ad Systema Entomologiae Cl. Fabricii ordinata cura I. I. Zschachii, M. Bacc.* [iv +] 136 pp., 3 col. pls. Müller, Lipsiae [=Leipzig]. (14 July)

[Reprinted in 1789 as pp. 1-136 in Karsten, G., *Musevm Leskeanvm. Regnum Animale. Qvod ordine systematico disposuit atque descripsit D. L. Gustavvs Karsten*. Vol. 1, [42 +] xiv + 320 pp., frontispiece, 8 col. pls. Müller, Lipsiae [=Leipzig]. Reviewed in *Gött. Gel. Anz.* 1788: 1123 (Evenhuis *in litt.*.)]

## Postscript

This project began with a real problem. I continually discovered situations where there were identical names which applied to different flies. These situations required that one name had to be changed, and in most cases, it was the name that workers were using that had to be changed. Naturally, my colleagues complained with the familiar refrain, "Why don't you make up your mind?" So, I wanted to examine sources of all names to gather the data necessary to resolve nomenclatural matters so names would no longer be changed for purely legalistic reasons (as opposed to changes resulting from an increase of our knowledge of species). Alternative courses of action were available: Rather than making the changes mandated by the rules, one could apply for an exception to the rules, or one could simply ignore the rules. However, both the alternatives are unethical, being contrary to the principle of priority and respect for history. The promise of Automatic Data Processing (ADP) made the task seem simple and guaranteed that the data could be shared by all. Thus, with the desire to resolve the problem of old names once and for all times, I started my odyssey through ancient history and modern ADP. Now that I am at the end of this odyssey, I offer a postscript.

Data standards (including editorial) are an often forgotten essential element. The lack of complete and comprehensive data standards haunted the project from the start and was the cause of much unproductive effort. ADP and desktop publishing are wonderful tools as enormous amounts of data can be quickly processed into eloquent publications, but they can just as easily turn the data into a mountain of garbage. What determines the difference is good standards and rigid adherence to them. While we followed the community standards, such as those of ICZN and BIOSIS, I discovered that many details are not covered by them. Hence, I evolved my own supplemental standards (Thompson *et alia* 1990). These are available to all. Beyond standards and ADP, xerox was essential. Much of the older literature is now almost inaccessible, locked away in rare book rooms. While we felt we got most things right the first time, we were always amazed by what we missed that first time. Xerox provided working copies that were readily available for re-examination. Sherborn's *Index Animalium* done without xerox and computers is in retrospect truly a Herculean accomplishment.

The *Musca* database was developed and is maintained on a Wang VS minicomputer using the Data Management System (DMS) supplied with the VS operating system. The text was done in Wang Word-Processing. VENTURA PUBLISHER was used to generate this book. COBOL programs were written to format the data for VENTURA. The word-processing documents

## Systematic Database of *Musca* Names

were converted to standard PC ASCII text files. Then these files were merged and printed with VENTURA. This database is now being converted from the Wang DMS environment to the PC environment.

From the *Musca* data file evolved the *Biosystematic Database of World Diptera*, which now contains virtually all the family-group (4,296) and genus-group (14,413) names that apply to flies and about 1/3 of the species-group names (30,658). From these data various publications are planned, such as *Systematic Database of Nearctic Diptera*, and eventually all the data will be available on CD-ROM disks. A number of systematists are already working with me so my journey is no longer solitary and I hope more dipterists will join us. So the *Musca* database must be viewed as only the first part of an international cooperative effort to master the names of flies that we know. From there we will be able to name more easily the ones we do not know. And that is a large part of the first task in understanding the biodiversity that man's world and welfare depend upon. So, I conclude this work by dedicating it to all my colleagues who have helped and continue to help me master the names for flies.

F. CHRISTIAN THOMPSON

1 October 1992  
Washington, D. C.













39088007650385



\* 3 9 0 8 8 0 0 7 6 5 0 3 8 5 \*